

APP-003197

**IRRIGATION BASED AGRICULTURAL ACTIVITIES AND CULTIVATION OF
GENETICALLY MODIFIED MAIZE ON FARM CUXHAVEN OOS, OSHIKOTO
REGION**

ENVIRONMENTAL ASSESSMENT SCOPING REPORT




Assessed by:



Assessed for:

L M Potgieter

December 2023

| | | |
|----------------------------------|---|---|
| Project: | IRRIGATION BASED AGRICULTURAL ACTIVITIES AND CULTIVATION OF GENETICALLY MODIFIED MAIZE ON FARM CUXHAVEN OOS, OSHIKOTO REGION: ENVIRONMENTAL ASSESSMENT SCOPING REPORT | |
| Report Version/Date: | Final December 2023 | |
| Prepared for: (Proponent) | L M Potgieter P O Box 761 Tsumeb | |
| Lead Consultant | Geo Pollution Technologies (Pty) Ltd PO Box 11073 Windhoek Namibia | TEL.: (+264-61) 257411 FAX.: (+264) 88626368 |
| Main Project Team: | Quzette Bosman (BA. Geography/Sociology); (BA Environmental Management) André Faul (B.Sc. Zoology/Biochemistry); (B.Sc. (Hons) Zoology); (M.Sc. Conservation Ecology); (Ph.D. Medical Bioscience) Pierre Botha (B.Sc. Geology/Geography); (B.Sc. (Hons) Hydrology/Hydrogeology) Johann Strauss (BA. Geography/Psychology/Environmental Management) Stefan Short Health and Safety Supervisor/GIS Technician | |
| Cite this document as: | Bosman Q, Botha P, Faul A, Short S; Strauss J. 2023 December, Irrigation Based Agricultural Activities and Cultivation of Genetically Modified Maize on Farm Cuxhaven Oos, Oshikoto Region: Environmental Assessment Scoping Report. | |
| Copyright | Copyright on this document is reserved. No part of this document may be utilised without the written permission of Geo Pollution Technologies (Pty) Ltd. | |
| Report Approval |  André Faul Conservation Ecologist | |

I _____, acting as representative of (L M Potgieter), hereby confirm that the project description contained in this report is a true reflection of the information which the Proponent provided to Geo Pollution Technologies. All material information in the possession of the Proponent that reasonably has or may have the potential of influencing any decision or the objectivity of this assessment is fairly represented in this report and the report is hereby approved.

Signed at _____ on the _____ day of _____ 2024
2023.

L M Potgieter

70071400370
ID Number

EXECUTIVE SUMMARY

Geo Pollution Technologies (Pty) Ltd was appointed by L M Potgieter (the Proponent) to undertake an environmental assessment for irrigation activities and cultivation of genetically modified maize on the farm Cuxhaven Oos FMB/01278 in the Oshikoto Region. Existing activities on the farm are focussed on irrigated crop cultivation and livestock farming. The Proponent currently utilizes an area of approximately 260 ha for cultivation, of which 76 ha is irrigated by means of centre pivot systems utilising abstracted groundwater, and the remaining 184 ha is used for dry land crop production. In order to optimize cultivation of maize, the Proponent wish to apply for the necessary permits to cultivate genetically modified maize. The genetically modified maize events (strains) earmarked for cultivation are insect resistant, herbicide tolerant and a combination of insect resistant and herbicide tolerant maize.

The main operational activities related to agriculture include:

- ◆ land preparation,
- ◆ planting,
- ◆ water abstraction and irrigation,
- ◆ fertilizer application and pest control,
- ◆ harvesting, and transporting activities specific to each crop, and
- ◆ cattle, sheep and potentially other livestock farming, and
- ◆ bush clearing and charcoal production (only as part of rangeland management).

All historically cleared areas for crop cultivation and rangeland improvement across the farm, including the existing and potential irrigation areas amount to approximately 349 ha. Pending the outcome of a hydrogeological specialist study, the total hectares of land to be irrigated simultaneously, may be increased. For irrigation, water is abstracted from two registered production boreholes on Cuxhaven Oos. The boreholes are registered with the Ministry of Agriculture, Water and Land Reform and the Proponent has a valid water permit for water abstraction. The main produce cultivated are vegetables and maize for local and international markets.

The environmental assessment determines all environmental, safety, health and socio-economic impacts associated with the continued and planned agricultural activities on the farm. Relevant environmental data was compiled by making use of primary data (hydrogeological specialist study), secondary data and from a reconnaissance site visit. Potential environmental impacts and associated social impacts were identified and are addressed in this report.

The project area is located amidst other farms and due to the nature and location of the Proponent's agricultural activities, limited impacts can be expected on the surrounding environment. Regular environmental performance monitoring is thus recommended to ensure regulatory compliance and the implementation of corrective measures when necessary, especially with regards to water abstraction and the planting of genetically modified maize. The Proponent's operations play a role in contributing to the Namibian agricultural sectors and provide valuable employment opportunities in the region.

The main concerns related to the operations are potential groundwater, surface water and soil contamination, decreased groundwater availability, ecological and social impacts. The addition of genetically modified maize on the farm, if not implemented responsibly, has the potential of aggravating existing impacts or causing additional impacts, while also being contentious issue for some people. A safety, health, environmental and quality policy coupled to an environmental management plan will contribute to effective management procedures, to prevent and mitigate impacts. All regulations relating to agriculture, genetically modified organisms, labour, and health and safety relevant legislation should be adhered to. Groundwater and soil pollution must be prevented at all times. Restrictions and prescriptions pertaining to the environmental release and handling of genetically modified maize should be strictly adhered to. This include, but is not limited to, planting of refuges, maintaining adequate buffers between genetically modified and traditional maize fields, correct pesticide application and vigilance and reporting of any signs of insect or weed resistance onset. All staff must be made aware of the importance of biodiversity and poaching or illegal harvesting of animal and plant products prohibited. Groundwater abstraction permits must be strictly adhered to. Any waste produced must be burned or removed from site and disposed of at an appropriate facility or re-used or recycled where

possible. Hazardous waste must be disposed of at an approved hazardous waste disposal site. By appointing local employees and by implementing monitoring and training programs, the positive socio-economic impacts can be maximised while preventing mitigating negative impacts.

The environmental management plan included in Section 10 of this document should be used as an on-site reference document during all phases (planning, operations (including maintenance) and decommissioning) of the development. All monitoring and records kept should be included in six monthly reports to ensure compliance with the environmental management plan and the Ministry of Environment, Forestry and Tourism's requirements. Parties responsible for transgression of the environmental management plan should be held responsible for any rehabilitation that may need to be undertaken. A safety, health, environmental and quality policy should be used in conjunction with the environmental management plan. Operators and responsible personnel must be taught the contents of these documents. Local or national regulations and guidelines must be adhered to and monitored regularly as outlined in the environmental management plan.

TABLE OF CONTENTS

| | | |
|----|---|----|
| 1 | BACKGROUND AND INTRODUCTION..... | 1 |
| 2 | SCOPE | 2 |
| 3 | METHODOLOGY..... | 2 |
| 4 | OPERATIONS AND RELATED ACTIVITIES | 3 |
| | 4.1 LAND CLEARING | 3 |
| | 4.1 ARABLE FARMING..... | 4 |
| | 4.2 CULTIVATION OF GENETICALLY MODIFIED MAIZE | 6 |
| | 4.3 LIVESTOCK | 7 |
| | 4.4 SUPPORT INFRASTRUCTURE | 7 |
| | 4.5 IRRIGATION AND WATER SUPPLY | 10 |
| | 4.6 EMPLOYMENT | 13 |
| 5 | ALTERNATIVES..... | 14 |
| | 5.1 LOCATION ALTERNATIVES..... | 14 |
| | 5.2 PROJECT IMPLEMENTATION AND DESIGN ALTERNATIVES..... | 14 |
| | 5.2.1 Irrigation Methods..... | 14 |
| | 5.2.2 Soil Preparation..... | 15 |
| | 5.2.3 Crop Selection (Maize)..... | 16 |
| | 5.3 No Go ALTERNATIVE..... | 17 |
| 6 | ADMINISTRATIVE, LEGAL AND POLICY REQUIREMENTS..... | 18 |
| 7 | ENVIRONMENTAL CHARACTERISTICS | 22 |
| | 7.1 LOCALITY AND SURROUNDING LAND USE..... | 22 |
| | 7.2 CLIMATE..... | 23 |
| | 7.3 TOPOGRAPHY, DRAINAGE..... | 26 |
| | 7.4 SOIL..... | 27 |
| | 7.5 GEOLOGY AND HYDROGEOLOGY | 29 |
| | 7.6 PUBLIC WATER SUPPLY | 31 |
| | 7.7 ECOLOGY..... | 31 |
| | 7.8 LOCAL ECONOMY..... | 34 |
| | 7.9 DEMOGRAPHIC PROFILE | 35 |
| | 7.10 CULTURAL, HERITAGE AND ARCHAEOLOGICAL ASPECTS..... | 36 |
| 8 | PUBLIC CONSULTATION..... | 37 |
| 9 | MAJOR IDENTIFIED IMPACTS..... | 37 |
| | 9.1 SOIL AND GROUNDWATER CONTAMINATION | 37 |
| | 9.2 GROUNDWATER ABSTRACTION | 37 |
| | 9.3 FIRE..... | 38 |
| | 9.4 DUST AND AIR QUALITY..... | 39 |
| | 9.5 TRAFFIC..... | 39 |
| | 9.6 HEALTH AND SAFETY | 39 |
| | 9.7 ECOSYSTEM AND BIODIVERSITY IMPACT | 39 |
| | 9.8 PESTICIDE RESISTANCE..... | 39 |
| | 9.9 HORIZONTAL GENE TRANSFER..... | 39 |
| | 9.10 SOCIO-ECONOMIC IMPACTS | 39 |
| 10 | ASSESSMENT AND MANAGEMENT OF IMPACTS | 40 |
| | 10.1 RISK ASSESSMENT AND ENVIRONMENTAL MANAGEMENT PLAN | 41 |
| | 10.1.1 Planning..... | 43 |
| | 10.1.2 Revenue Generation in the Professional Sector | 44 |
| | 10.1.3 National Development Goals: Water, Agriculture and Land Use Planning | 45 |
| | 10.1.4 Skills and Development..... | 46 |
| | 10.1.5 Revenue Generation and Employment..... | 47 |
| | 10.1.6 Demographic Profile and Community Health..... | 48 |
| | 10.1.7 Agricultural Produce..... | 49 |

| | | |
|-----------|---|-----------|
| 10.1.8 | Traffic | 50 |
| 10.1.9 | Health, Safety and Security | 51 |
| 10.1.10 | Fire | 52 |
| 10.1.11 | Noise | 53 |
| 10.1.12 | Waste Production | 54 |
| 10.1.13 | Ecosystem and Biodiversity Impact | 56 |
| 10.1.14 | Groundwater, Surface Water and Soil Contamination | 58 |
| 10.1.15 | Groundwater Abstraction | 59 |
| 10.1.16 | Visual Impact | 60 |
| 10.1.17 | Cumulative Impact | 61 |
| 10.2 | DECOMMISSIONING AND REHABILITATION | 62 |
| 10.3 | ENVIRONMENTAL MANAGEMENT SYSTEM | 62 |
| 11 | CONCLUSION | 62 |
| 12 | BIBLIORGAPHY | 64 |

LIST OF APPENDICES

| | | |
|-------------|--|-----|
| APPENDIX A: | CERTIFICATES | 66 |
| APPENDIX B: | HYDROGEOLOGICAL SPECIALIST STUDY | 68 |
| APPENDIX C: | GM MAIZE AND COTTON IN NAMIBIA SPECIALIST REPORT | 94 |
| APPENDIX D: | TREE INFORMATION | 173 |
| APPENDIX E: | PROOF OF PUBLIC CONSULTATION | 179 |
| APPENDIX F: | CONSULTANTS' CURRICULUM VITAE | 194 |

LIST OF FIGURES

| | | |
|------------|---|----|
| FIGURE 1-1 | PROJECT LOCATION | 1 |
| FIGURE 4-1 | CLEARED AREAS ON THE FARM | 3 |
| FIGURE 4-2 | CULTIVATED AND CLEARED AREAS | 4 |
| FIGURE 4-3 | MAP WITH INFRASTRUCTURE COMPONENTS | 10 |
| FIGURE 4-4 | LOCATIONS OF BOREHOLES | 13 |
| FIGURE 7-1 | PROPERTIES ADJACENT TO THE PROJECT AREA | 23 |
| FIGURE 7-2 | DAILY AND SEASONAL RAINFALL (FUNK ET AL., 2015) | 25 |
| FIGURE 7-3 | ASPECT SLOPE AND SURFACE DRAINAGE | 27 |
| FIGURE 7-4 | ROCK TYPE AND AGRO ECOLOGICAL ZONE (ATLAS OF NAMIBIA PROJECT, 2002) | 28 |
| FIGURE 7-5 | GROUNDWATER CATCHMENTS AND WATER CONTROL AREAS | 30 |
| FIGURE 9-1 | CONCEPTUAL GROUNDWATER BALANCE WITH OVER ABSTRACTION SCENARIO | 38 |

LIST OF TABLES

| | | |
|-----------|--|----|
| TABLE 4-1 | GM MAIZE EVENTS EARMARKED FOR CULTIVATION BY THE PROPONENT | 6 |
| TABLE 4-2 | SUMMARY OF INFRASTRUCTURE COMPONENTS RELATED TO AGRICULTURAL OPERATIONS | 9 |
| TABLE 4-3 | SUMMARY OF BOREHOLE INFORMATION OBTAINED FROM THE PROPONENT | 11 |
| TABLE 5-1 | IRRIGATION SYSTEM EFFICIENCY (IWRM PLAN JOINT VENTURE NAMIBIA, 2010) | 15 |
| TABLE 5-2 | ADVANTAGES AND DISADVANTAGES OF LAND PREPARATION SYSTEMS (ADAPTED FROM HTTPS://CROPWATCH.UNL.EDU/TILLAGE/ADVDISADV) | 15 |
| TABLE 5-3 | ALTERNATIVE COMPARISON OF MAIZE TYPES FOR CULTIVATION | 16 |
| TABLE 6-1 | NAMIBIAN LAW APPLICABLE TO THE DEVELOPMENT | 18 |
| TABLE 6-2 | GUIDING DOCUMENTS, DIRECTIVES AND STANDARDS | 20 |
| TABLE 6-3 | RELEVANT MULTILATERAL ENVIRONMENTAL AGREEMENTS | 20 |
| TABLE 7-1 | ADJACENT PROPERTIES | 22 |
| TABLE 7-2 | SUMMARY OF CLIMATE CLIMATIC CONDITIONS (ADAPTED ATLAS OF NAMIBIA PROJECT, 2002) | 24 |
| TABLE 7-3 | RAINFALL STATISTICS (FUNK ET AL., 2015) | 25 |
| TABLE 7-4 | GROUNDWATER STATISTICS | 31 |
| TABLE 7-5 | GENERAL PLANT DATA (ATLAS OF NAMIBIA PROJECT, 2002) | 32 |

| | | |
|------------|---|----|
| TABLE 7-6 | TREES WITH CONSERVATION CONCERNS IN QUARTER DEGREE SQUARES 1918AA (CURTIS & MANNHEIMER, 2005) | 32 |
| TABLE 7-7 | IUNC RED LISTED SPECIES WHICH MAY OCCUR IN THE AREA..... | 33 |
| TABLE 7-8 | MAIN INDUSTRY OF EMPLOYED POPULATION AGED 15 YEARS AND ABOVE FOR THE TSUMEB CONSTITUENCY AND OSHIKOTO REGION | 34 |
| TABLE 7-9 | DEMOGRAPHIC CHARACTERISTICS OF THE TSUMEB CONSTITUENCY, THE OSHIKOTO REGION AND NATIONALLY (NAMIBIA STATISTICS AGENCY, 2011)..... | 36 |
| TABLE 10-1 | ASSESSMENT CRITERIA..... | 40 |
| TABLE 10-2 | ENVIRONMENTAL CLASSIFICATION (PASTAKIA 1998) | 41 |

LIST OF PHOTOS

| | | |
|------------|--|----|
| PHOTO 4-1 | CLEARED AREAS BETWEEN CULTIVATED AREAS | 4 |
| PHOTO 4-2 | CLEARED AREAS ADJACENT TO IRRIGATED CENTRE PIVOT FIELD | 4 |
| PHOTO 4-3 | ONION FIELD | 5 |
| PHOTO 4-4 | POTATO FIELD | 5 |
| PHOTO 4-3 | DRY LAND MAIZE CULTIVATION | 5 |
| PHOTO 4-4 | IRRIGATED MAIZE FIELD | 5 |
| PHOTO 4-5 | FERTILISER MIXING TANK..... | 5 |
| PHOTO 4-6 | PESTICIDE MIXING UNIT | 5 |
| PHOTO 4-3 | PREPARED ONION FIELD..... | 5 |
| PHOTO 4-4 | IRRIGATED MAIZE FIELD DAMAGED BY WARTHOG AND / OR PORCUPINE..... | 5 |
| PHOTO 4-5 | MAIZE DAM..... | 6 |
| PHOTO 4-6 | POTATO WASHING AND SORTING EQUIPMENT | 6 |
| PHOTO 4-11 | SHEEP HOLDING PEN..... | 7 |
| PHOTO 4-12 | CATTLE GRAZING ON CROP FIELDS | 7 |
| PHOTO 4-17 | SOLAR PANEL USED AT BOREHOLES | 8 |
| PHOTO 4-18 | FUEL STORAGE TANKS | 8 |
| PHOTO 4-21 | RAISED WATER TANKS FOR DOMESTIC USE | 8 |
| PHOTO 4-22 | STOCK WATERING RESERVOIR | 8 |
| PHOTO 4-23 | EQUIPMENT STORAGE AND MAINTENANCE SHED..... | 9 |
| PHOTO 4-24 | ANIMAL FEED AND SUPPLEMENT STORAGE | 9 |
| PHOTO 4-25 | CHEMICAL STORAGE ROOM ENTRANCE AND RELATED SAFETY SIGNS..... | 9 |
| PHOTO 4-26 | FIRE HYDRANT AND SAFETY SIGNS AT THE ENTRANCE OF THE SHED | 9 |
| PHOTO 4-31 | CENTRE PIVOT SYSTEM ON ONION FIELDS | 11 |
| PHOTO 4-32 | CENTRE PIVOT IRRIGATION ON ESTABLISHED MAIZE FIELD | 11 |
| PHOTO 4-31 | BOREHOLE PUMP INSTALLATION USED FOR IRRIGATION..... | 12 |
| PHOTO 4-32 | BOREHOLE PUMP INSTALLATION USED FOR IRRIGATION (WWW10069)..... | 12 |
| PHOTO 4-33 | STOCK WATERING BOREHOLE..... | 12 |
| PHOTO 4-34 | DWA BOREHOLE WW100071 | 12 |
| PHOTO 4-35 | GENERAL EMPLOYEE ABLUTION | 14 |
| PHOTO 4-36 | MOBILE TOILET IN OPERATIONAL AREAS | 14 |
| PHOTO 4-35 | EMPLOYEE FOOTBALL FIELD..... | 14 |
| PHOTO 4-36 | EMPLOYEE VEGETABLE GARDEN | 14 |
| PHOTO 7-5 | CLAYEY LOAMY SOILS..... | 28 |
| PHOTO 7-6 | ONIONS IN CLAYEY LOAMY SOILS | 28 |
| PHOTO 7-9 | WILDLIFE SCAT | 33 |
| PHOTO 7-10 | TERMITE MOUND ON SITE | 33 |

LIST OF ABBREVIATIONS

| | |
|-----------------|---|
| AEZ | Agro-Ecological Zone |
| AIDS | Acquired Immune Deficiency Syndrome |
| BE | Biological/Ecological |
| Bt | <i>Bacillus thuringiensis</i> |
| BH | Borehole |
| CHIRPS-2 | Climate Hazards Group Infra-Red Precipitation with Station data |
| DWA | Department of Water Affairs |
| EA | Environmental Assessment |
| EIA | Environmental Impact Assessment |
| EMA | Environmental Management Act No 7 of 2007 |
| EMP | Environmental Management Plan |
| EMS | Environmental Management System |
| EO | Economic/Operational |
| ES | Environmental Classification |
| GDP | Gross Domestic Product |
| GM | Genetically Modified |
| GMO | Genetically Modified Organism |
| GPS | Global Positioning System |
| GPT | Geo Pollution Technologies |
| HGT | Horizontal Gene Transfer |
| HIV | Human Immunodeficiency Virus |
| IAPs | Interested and Affected Parties |
| IUCN | International Union for Conservation of Nature |
| LNAPL | Light Non-Aqueous Phase Liquids |
| mamsl | Meters Above Mean Sea Level |
| m/s | Metre per second |
| mbs | Metres below surface |
| MEFT | Ministry of Environment, Forestry and Tourism |
| MAWLR | Ministry of Agriculture, Water and Land Reform |
| mm/a | Millimetres per annum |
| MSDS | Material Safety Data Sheet |
| NCRST | National Commission on Research, Science and Technology |
| NDP | National Development Plan |
| PC | Physical/Chemical |
| PPE | Personal Protective Equipment |
| ppm | Parts per million |
| SANS | South African National Standards |
| SC | Sociological/Cultural |
| SHEQ | Safety, Health, Environment and Quality |
| SRTM | Shuttle Radar Topography Mission |
| UNFCCC | United Nations Framework Convention on Climate Change |
| WHO | World Health Organization |

GLOSSARY OF TERMS

Alternatives - A possible course of action, in place of another, that would meet the same purpose and need but which would avoid or minimize negative impacts or enhance project benefits. These can include alternative locations/sites, routes, layouts, processes, designs, schedules and/or inputs. The “no-go” alternative constitutes the ‘without project’ option and provides a benchmark against which to evaluate changes; development should result in net benefit to society and should avoid undesirable negative impacts.

Assessment - The process of collecting, organising, analysing, interpreting and communicating information relevant to decision making.

Competent Authority - means a body or person empowered under the local authorities act or Environmental Management Act to enforce the rule of law.

Construction - means the building, erection or modification of a facility, structure or infrastructure that is necessary for the undertaking of an activity, including the modification, alteration, upgrading or decommissioning of such facility, structure or infrastructure.

Cumulative Impacts - in relation to an activity, means the impact of an activity that in itself may not be significant but may become significant when added to the existing and potential impacts eventuating from similar or diverse activities or undertakings in the area.

Environment - As defined in the Environmental Assessment Policy and Environmental Management Act - “land, water and air; all organic and inorganic matter and living organisms as well as biological diversity; the interacting natural systems that include components referred to in sub-paragraphs, the human environment insofar as it represents archaeological, aesthetic, cultural, historic, economic, palaeontological or social values”.

Environmental Impact Assessment (EIA) - process of assessment of the effects of a development on the environment.

Environmental Management Plan (EMP) - A working document on environmental and socio-economic mitigation measures, which must be implemented by several responsible parties during all the phases of the proposed project.

Environmental Management System (EMS) - An Environment Management System, or EMS, is a comprehensive approach to managing environmental issues, integrating environment-oriented thinking into every aspect of business management. An EMS ensures environmental considerations are a priority, along with other concerns such as costs, product quality, investments, PR productivity and strategic planning. An EMS generally makes a positive impact on a company’s bottom line. It increases efficiency and focuses on customer needs and marketplace conditions, improving both the company’s financial and environmental performance. By using an EMS to convert environmental problems into commercial opportunities, companies usually become more competitive.

Evaluation –The process of ascertaining the relative importance or significance of information, the light of people’s values, preference and judgements in order to make a decision.

Green Scheme - The Green Scheme is an initiative conducted by the Ministry of Agriculture, Water and Forestry to encourage the development of irrigation based agronomic production in Namibia with the aim of increasing the contribution of agriculture to the country's Gross Domestic Product. Its aim is also to simultaneously achieve the social development and upliftment of communities located within suitable irrigation areas and to also promote the human resources and skills development within the irrigation sub-sector. Such initiative could possibly enhance cross-border investment and facilitate the exchange of relevant and limited resources with neighbouring countries in this regard.

Hazard - Anything that has the potential to cause damage to life, property and/or the environment. The hazard of a particular material or installation is constant; that is, it would present the same hazard wherever it was present.

Interested and Affected Party (IAP) - any person, group of persons or organisation interested in, or affected by an activity; and any organ of state that may have jurisdiction over any aspect of the activity.

Mitigate - The implementation of practical measures to reduce adverse impacts.

Proponent (Applicant) - Any person who has submitted or intends to submit an application for an authorisation, as legislated by the Environmental Management Act no. 7 of 2007, to undertake an activity or activities identified as a listed activity or listed activities; or in any other notice published by the Minister or Ministry of Environment & Tourism.

Public - Citizens who have diverse cultural, educational, political and socio-economic characteristics. The public is not a homogeneous and unified group of people with a set of agreed common interests and aims. There is no single public. There are a number of publics, some of whom may emerge at any time during the process depending on their particular concerns and the issues involved.

Scoping Process - process of identifying: issues that will be relevant for consideration of the application; the potential environmental impacts of the proposed activity; and alternatives to the proposed activity that are feasible and reasonable.

Significant Effect/Impact - means an impact that by its magnitude, duration, intensity or probability of occurrence may have a notable effect on one or more aspects of the environment.

Stakeholder Engagement - The process of engagement between stakeholders (the proponent, authorities and IAPs) during the planning, assessment, implementation and/or management of proposals or activities. The level of stakeholder engagement varies depending on the nature of the proposal or activity as well as the level of commitment by stakeholders to the process. Stakeholder engagement can therefore be described by a spectrum or continuum of increasing levels of engagement in the decision-making process. The term is considered to be more appropriate than the term “public participation”.

Stakeholders - A sub-group of the public whose interests may be positively or negatively affected by a proposal or activity and/or who are concerned with a proposal or activity and its consequences. The term therefore includes the proponent, authorities (both the lead authority and other authorities) and all interested and affected parties (IAPs). The principle that environmental consultants and stakeholder engagement practitioners should be independent and unbiased excludes these groups from being considered stakeholders.

Sustainable Development - “Development that meets the needs of the current generation without compromising the ability of future generations to meet their own needs and aspirations” – the definition of the World Commission on Environment and Development (1987). “Improving the quality of human life while living within the carrying capacity of supporting ecosystems” – the definition given in a publication called “Caring for the Earth: A Strategy for Sustainable Living” by the International Union for Conservation of Nature (IUCN), the United Nations Environment Programme and the World Wide Fund for Nature (1991).

1 BACKGROUND AND INTRODUCTION

Geo Pollution Technologies (Pty) Ltd was appointed by L M Potgieter (the Proponent) to undertake an environmental assessment for the existing and proposed agricultural activities on the Farm Cuxhaven Oos No.1278, located in the Abenab area of the Oshikoto Region (Figure 1-1). The main commercial activities of the Proponent on the farm includes crop cultivation and livestock farming. An additional planned activity by the Proponent is the cultivation of genetically modified (GM) maize. For purposes of crop cultivation, the Proponent utilizes approximately 260 ha for irrigation and dryland farming. Pending the outcome of a hydrogeological specialist study, the total hectares of land to be irrigated simultaneously, may be increased. Irrigation are from two production boreholes by means of centre pivot irrigation systems. The main operational activities include:

- ◆ land preparation,
- ◆ planting (including proposed planting of genetically modified (GM) maize),
- ◆ water abstraction and irrigation,
- ◆ fertilizer application and pest control,
- ◆ harvesting,
- ◆ packaging and transporting activities specific to each crop,
- ◆ cattle, sheep and potentially other livestock farming, and
- ◆ bush clearing and charcoal production (only as part of rangeland management).




Figure 1-1 Project location

A detailed project description is provided in Section 4. The potential impacts of the project on the environment, resulting from various operational, maintenance and construction, and possible decommissioning activities, were determined through the risk assessment as presented in this report.

The environment, being defined in the Environmental Management Act as “land, water and air; all organic and inorganic matter and living organisms as well as biological diversity; the interacting natural systems that include components referred to in sub-paragraphs, the human environment insofar as it

represents archaeological, aesthetic, cultural, historic, economic, paleontological or social values". The environmental assessment was conducted to apply for an environmental clearance certificate in compliance with Namibia's Environmental Management Act (Act No 7 of 2007) (EMA).

Project Justification – Traditionally farms in the region were used for cattle ranching with limited dryland crop cultivation. However, in the area, including on the Proponent's farm, farming activities were diversified to include irrigation based crop cultivation. The Proponent has a well-established irrigation and agriculture development, which sees an optimisation of crop production by means of irrigation, augmented by rainwater. It is now the Proponent's intention to further diversify farming activities by cultivating GM Maize. This addition is proposed in an effort to increase resilience in food production for Namibia. Namibia aims on increasing sustainable food production and ensuring food security in the country. In addition, agriculture is an important employment sector for Namibia, adding to roughly a third of the workforce. Existing and planned agricultural activities require employment, which is required to be maintained for continued operations. Pivot irrigation systems also require significant investment costs and therefore the development of the irrigation areas, has ensured a sizeable investment into the area and the Abenab district.

Benefits of the agricultural activities conducted by the Proponent include.

- ◆ Food production and enhanced food security.
- ◆ Employment and supporting of livelihoods of both unskilled and skilled labourers.
- ◆ Technological development and investment in agricultural practices.
- ◆ Generation of income that contributes to the national treasury and a positive trade balance through the export of produce to international markets.
- ◆ Support for economic resilience in the area through diversified business activities and opportunities.

2 SCOPE

The scope of this report is to, in compliance with the requirements of EMA:

1. Present a detailed project and environmental description related to the Proponent's activities.
2. Determine the potential environmental impacts emanating from the Proponent's activities and potential future decommissioning of such activities.
3. Identify a range of management actions to mitigate the potential adverse impacts to acceptable levels.
4. Provide sufficient information to the relevant competent authority and the Ministry of Environment, Forestry and Tourism (MEFT) and related authorities to make an informed decision regarding the project and the issuing of an environmental clearance certificate.

3 METHODOLOGY

Methods employed to investigate and report on potential impacts of the Proponent's activities on the social and natural environment include:

1. Detailed infrastructure and operational procedures received from the client are presented in this report.
2. Baseline information about the site and its surroundings were obtained from primary information (hydrogeological assessment), existing secondary information as well as from a reconnaissance site visit.
3. A specialist report related to the benefits, impacts and concerns of environmental release of GM crops was compiled and the findings of this report was considered in the environmental assessment.
4. As part of the scoping process to determine potential environmental impacts, interested and affected parties (IAPs) were consulted about their views, comments and opinions, all of which are presented in this report.

5. As per the findings of this environmental assessment, a scoping report with an environmental management plan (EMP) were prepared and this will be submitted to the MEFT.

4 OPERATIONS AND RELATED ACTIVITIES

Agricultural activities, focussing on local food production, have constituted the core of the Proponent's operations. Agricultural activities have been conducted on the farm for the past 50 years with the Proponent continuing and gradually expanding agricultural activities for the past ten years. Various vegetables and maize are produced throughout the year, some of which are packaged on site before transport to local markets. In an attempt to increase resilience of maize and thereby increasing local maize yields, the Proponent wish to, in the future, cultivate GM maize. Livestock farming, mainly cattle and sheep, is conducted while there is also some game on the farm. However, game farming and related fencing is not an active pursuit of the Proponent. Existing and planned operations are reliant on support infrastructure and resources, all of which are described below.

4.1 LAND CLEARING

The farm is a known agricultural unit for, more than 50 years as indicated on the 1975 topographic map indicating crop cultivation areas on the farm. Initial land clearing was conducted to accommodate dryland cropping. More recent agricultural activities have seen an expansion of the cleared areas for irrigation based crop cultivation, while also realising rangeland improvement through bush-thinning activities. Such activities are especially required for sheep husbandry, which forms part of the farming operations. Approximately 260 ha across the farming unit has been cleared for irrigation and dryland crop production, while approximately 89 ha is managed rangelands. Additional areas have been identified for future rangeland improvement. The areas described above are presented in Figure 4-1. Vegetation was also cleared, and is maintained so, next to fences to accommodate firefighting.




Figure 4-1 Cleared areas on the farm



Photo 4-1 Cleared areas between cultivated areas



Photo 4-2 Cleared areas adjacent to irrigated centre pivot field

4.1 ARABLE FARMING

A variety of crops and vegetables are planted on a rotational basis over the farm. Crop rotation is required as pest management measure to ensure crop harvests do not fail. For example, potatoes should only be planted on the same patch of soil, every five years. Figure 4-2 depicts cultivated fields and cleared areas. Most of these vegetables and crops are irrigated by means of moveable or fixed centre pivot systems. Crops and vegetables produced on the farm include onions, potatoes, carrots and wheat. Some of these vegetables, such as the onions and potatoes, are required to be hand harvested and are therefore labour intensive. Dry land and irrigation based crop production of maize is also conducted. Each crop has a different planting and cultivation regime. The majority of produce are annual crops, which require one or two seasons to complete their life cycle. Preparation of the land and planting of annual crops entail mechanical activities like ripping, tilling and seeding of the soil, with tractors and specialised implements.




Figure 4-2 Cultivated and cleared areas



Photo 4-3 Onion field



Photo 4-4 Potato field



Photo 4-5 Dry land maize cultivation



Photo 4-6 Irrigated maize field

Fertilizers and pesticides are applied as required and according to the specifications for application. For irrigated fields, fertilisers are mixed with water in a large mixing tank. Once the desired mixing ratio is achieved, the fertiliser is fed into the irrigation system for administration onto the crops. The Proponent utilises a low-soluble-nutrient fertilizer which may be readily absorbed by crops and requires less water. Pesticides are administered as per the specified application procedures for the corresponding pest by means of tractor spraying. To ensure correct and safe application of pesticides, a pesticide plan is implemented and regularly updated. The Proponent requires a minimum amount of pesticides as compared to conventional agricultural production. Some crops, such as onions are sensitive to chemical treatment and therefore the majority of weed control is conducted manually. Larger mammals, such as warthog, porcupine and antelope, can cause considerable damage to maize fields.



Photo 4-7 Fertiliser mixing tank



Photo 4-8 Pesticide mixing unit



Photo 4-9 Prepared onion field



Photo 4-10 Irrigated maize field damaged by warthog and / or porcupine

Various harvesting techniques are employed for the variety of crops. For example, although onions and potatoes are hand harvested, they still have to be mechanically loosened from the soil, which require specialist implements. After the potatoes and onions have been harvested, they are washed sorted and packed on the farm. To facilitate this process, the Proponent has invested in related sorting and packing machinery. Any produce not fit for human consumption is provided as feed to livestock and/or game on the farm.



Photo 4-11 Maize dam



Photo 4-12 Potato washing and sorting equipment

4.2 CULTIVATION OF GENETICALLY MODIFIED MAIZE

The Proponent plans to cultivate GM maize. Applications for the environmental release of GM maize for cultivation, based on existing procedures, policies and plans, will be submitted to the National Commission on Research Science and Technology (NCRST) under the Ministry of Higher Education, Technology and Innovation for approval. Such applications must be accompanied by a completed application form, this environmental assessment and its accompanying EMP, the ECC, and emergency response plans for its cultivation and transport. The GM maize events earmarked for cultivation are listed in Table 4-1. Each type of maize is referred to as an “event”.

Table 4-1 GM maize events earmarked for cultivation by the Proponent

| Event | Commonly Referred/Trade Name | Trait |
|--------------------|-------------------------------------|---|
| MON 810 | Bt Maize/ YieldGard™ | Resistant to lepidopteran* larvae like African maize stalk borer and fall armyworm |
| MON 89034 | Bt Maize/ YieldGard™ VT Pro™ | Resistant to lepidopteran larvae African maize stalk borer and fall armyworm |
| NK 603 | Roundup Ready™ 2 Maize | Resistant (tolerant) to glyphosate herbicide (RoundUp™) |
| MON 89034 × NK 603 | Roundup Ready® Maize 2 | Resistant (tolerant) to glyphosate herbicide (RoundUp™) and resistant to lepidopteran larvae like African maize stalk borer and fall armyworm |
| NK 603 × MON 810 | YieldGard™ CB + RR | Resistant (tolerant) to glyphosate herbicide (RoundUp™) and resistant to lepidopteran larvae like African maize stalk borer and fall armyworm |

*Lepidopterans are the order Lepidoptera comprising moths and butterflies

The insect resistant events are protected during an outbreak of pests like the African maize stalk borer and fall armyworm. These are the larvae (caterpillar) of moths. Due to a specific protein the plant produce as a result of the genetic modification, the larvae of the moths die when eating the maize plants, thus minimizing crop loss without the need for applying pesticides. Herbicide resistant events have been modified to be tolerant to RoundUp™ which is a broad spectrum herbicide with the active ingredient glyphosate. Post-emergent Roundup Ready™ maize can thus be sprayed with RoundUp™ to kill weeds without harming the maize plant itself. This eliminates the need for pre-planting weed control regimes and manual weed removal post-emergence.

Applications for the environmental release of GM maize for cultivation, based on existing procedures, policies and plans, will be submitted to the National Commission on Research Science and Technology (NCRST) under the Ministry of Higher Education, Technology and

Innovation for approval. Such applications must be accompanied by a completed application form, this environmental assessment and its accompanying EMP, the ECC, and emergency response plans for its cultivation and transport.

The specialist report in Appendix C provides a detailed description and assessment of GMOs in general and then specifically also the GM maize events to be planted by the Proponent. Note that the report also includes GM cotton events, although the Proponent is only interested in GM maize. The report addresses myths, truths and concerns regarding GMOs and provide prevention and mitigation measures required for GM maize cultivation. The cultivation of GM maize has received conditional approval by MEFT for its cultivation in Namibia. This approval was based on a strategic environmental assessment conducted in 2019/2020 (Faul et al. 2020). The conditional approval requires that individual assessments must be undertaken for each farming unit, as is being done in this report.

In the interim, until GM maize cultivation is approved, conventional maize cultivation continues. Once harvested, maize is collected in an on-site maize silo until it is collected or dispatched to clients or mills. Once GM maize is cultivated, harvested GM and conventional maize will be kept separate, should both be planted on the farm. If not kept separate, all maize will be considered as GM maize.

4.3 LIVESTOCK

The less suitable areas for crop cultivation is used for livestock rearing. Cattle and sheep are herded and managed as a separate business unit. A dedicated workforce manage all operations related to the sheep and cattle, which includes predator protection, watering equipment, lambing or calving support, flock vaccinations, hoof care, pasture management and meat marketing. At times, such as during nights or during lambing season, some of the sheep are kept in holding pens closer to the Proponent's main operations. A feedlot is employed at times to provide for additional support during droughts. Cattle are used to fertilise crop fields after harvesting, when they are allowed to graze on the maize stover or on resting / fallow crop fields.



Photo 4-13 Sheep holding pen



Photo 4-14 Cattle grazing on crop fields

4.4 SUPPORT INFRASTRUCTURE

Operations as outlined above, require support infrastructure or resources. The most crucial of these relate to water required for irrigation and potable use. Related support infrastructure are detailed below. Water and related irrigation systems are discussed in Section 4.5 while labour and related aspects are detailed in Section 4.6.

All operations on the farm are provided with **electricity** from a 33 kV Cenerod line. A 200 kVA photovoltaic solar system is planned for future development. Solar systems are only currently used at boreholes. Employee houses are serviced with electricity. The power line has a 9 m wide servitude which is kept clean by the Proponent for the portions of the power line over their farm. **Fuel** is stored in two above ground tanks of 220 l each for the use by mainly tractors and farming related operations.



Photo 4-15 Solar panel used at boreholes



Photo 4-16 Fuel storage tanks

Water is pumped from various boreholes for irrigation, stock watering and domestic use. Storage of water is determined by its use. While irrigation boreholes have no storage structures, stock watering rely on reservoirs while domestic use employ raised water storage tanks. All offices and employees' houses are provided with septic tank and french drain systems to accommodate waste water. **Waste disposal** mainly comprise an excavated pit where waste is regularly burned. Due to a lack of any recyclers in the area, recycling of certain wastes are not possible. However, where possible, certain waste items are not discarded, but rather re-used for alternative purposes. This includes the re-use of old oil when not collected by oil recycling companies. All spoiled produce are made available as animal feed. Any hazardous waste is stored in suitable bunded areas.



Photo 4-17 Raised water tanks for domestic use



Photo 4-18 Stock watering reservoir

A **storage and maintenance area** is located on the farm and comprise of a shed and storerooms where implements and other maintenance material are stored under roof and on impermeable surfaces. Any maintenance and or minor repairs are conducted on site and within these areas. Unused equipment and related materials are stored in an access controlled area. Offices and employee houses are all located on the farm as well. All pesticides and herbicides are stored in a dedicated **chemical store**. The chemical store is access controlled. Fertilisers are stored in an open area within the storage shed, separate from all other chemicals or materials, on an impermeable layer. All areas have firefighting equipment and safety signs where required. A summary of the support infrastructure components are presented in Table 4-2.



Photo 4-19 Equipment storage and maintenance shed



Photo 4-20 Animal feed and supplement storage



Photo 4-21 Chemical storage room entrance and related safety signs



Photo 4-22 Fire hydrant and safety signs at the entrance of the shed

Table 4-2 Summary of infrastructure components related to agricultural operations

| Project Component | Current Provision | Future Provision |
|--------------------------------------|---|---|
| Electricity Provision | Estimated 33 kVA mainly sourced from Cenored | No significant increase expected |
| Photovoltaic Solar System | Limited to borehole pump installations | 200 kVA |
| Water Provision | Groundwater abstraction from various boreholes | An increase in water allocation may be applied for |
| Water Storage | Various stock watering reservoirs and water tanks | No storage reservoirs planned for irrigation related activities |
| Equipment and General Storage | One existing storage complex | No additional storage proposed |
| Sanitation | Current septic tank and french drain systems catering for existing staff compliment | Additional septic tank and french drains may be required for planned expansions |
| Landfill | One landfill site | No additional sites planed |
| Fuel Storage | Two diesel tanks with a combined capacity of 4,200 l located in a bund wall | No additional tanks will be erected for the foreseeable future |
| Chemical Storage Area | One chemical storage unit | No additional chemical storage unit planned |





Figure 4-3 Map with infrastructure components

4.5 IRRIGATION AND WATER SUPPLY

The Proponent utilises groundwater for all aspects of operations. Existing and proposed irrigation on crops, make up the bulk of the water use and is the determining factor in terms of water use and related permitting. The irrigation systems employed on the farm is centre pivots. Moveable centre pivot systems allow the Proponent to move the systems from irrigation field (pivot circle) to irrigation field. Although the Proponent has several irrigation fields, he only has a couple of centre pivot systems which are moved among the fields, seasonally. For example, one pivot may serve a maize field in the summer while moving to another field during the fall and winter.

Phocaides (2007) provides a description of the centre pivot, being a low to medium pressure, fully mechanised, automated irrigation of permanent assemble. It basically comprise a sprinkler pipeline (usually of high tensile galvanized light steel or aluminium pipes) supported above ground by mobile A-frame towers, long spans, steel trusses and/or cables (Photo 4-23). The

pipeline is connected to a central tower with the “pivot mechanism” and main control panel. Moveable systems are mounted on wheels which allows it to be dragged from one field and fixed water supply point, to the next. The entire active irrigation system remains self-propelled to slowly rotate around the central tower while dispensing water through sprinklers (emitters) connected to the pipeline (Photo 4-24). An automatic alignment systems ensures the irrigation pipeline remains straight while a drive system enables the system movement. Small variations to the emitter sequence may be done when moving between different crops which may have different irrigation requirements. The Proponent has approximately 76 ha of pivot related irrigation fields on the farm and would like to further develop another 14 ha.



During the recognisance site visit all known boreholes on the farm were documented. Twelve boreholes were visited and data gathered about their status, use and physical description. A number of the boreholes are known DWA boreholes which are sealed. GPS locations of all boreholes were recorded and mapped, as presented in Figure 4-4. Of the boreholes surveyed, two are used for irrigation purposes.

The Proponent has a water abstraction permit for 120,000 m³ per year. Renewal of the permit is planned with no immediate increase in the allocation.

Table 4-3 Summary of borehole information obtained from the Proponent

| Map Ref. | Farm Name | Borehole Name(s) | Use | Borehole Depth (m) | Yield (m ³ /h) | Water Level (mbs) |
|----------|----------------------|-------------------------|----------------|--------------------|---------------------------|-------------------|
| EB1 | Cuxhaven Oos No.1278 | WW38010 (Klein pos gat) | Stock watering | 80 | 15 | |
| EB 2 | Cuxhaven Oos No.1278 | WW100065 | DWA BH | 200 | | 31 |
| EB 3 | Cuxhaven Oos No.1278 | | Not used | | 6 | 30 |
| EB 4 | Cuxhaven Oos No.1278 | | Not used | | | 31 |
| EB 5 | Cuxhaven Oos No.1278 | | Stock watering | 80 | 2 | |
| EB 6 | Cuxhaven Oos No.1278 | WW100071 | DWA BH | 200 | 8 | 45 |
| EB 7 | Cuxhaven Oos No.1278 | WW100069 | Irrigation | 200 | 300 | 48 |
| EB 8 | Cuxhaven Oos No.1278 | | Not used | 140 | 60 | 50 |
| EB 9 | Cuxhaven Oos No.1278 | WW 38011 (Huis gat) | Domestic | 100 | 20 | |
| EB 10 | Cuxhaven Oos No.1278 | WW39436 | Not used | 100 | | 42 |
| EB 11 | Cuxhaven Oos No.1278 | WW41033 | Irrigation | 180 | 50 | 73 |
| EB 12 | Cuxhaven Oos No.1278 | | Domestic | 100 | 30 | 49 |



Photo 4-25 Borehole pump installation used for irrigation (EB11)



Photo 4-26 Borehole pump installation used for irrigation (WW100069 / EB7)



Photo 4-27 Stock watering borehole (EB5)



Photo 4-28 DWA Borehole (WW100071 / EB13)




Figure 4-4 Locations of boreholes

4.6 EMPLOYMENT

All operations on the farm are reliant on labour. Operations currently require 10 permanent employees and up to 80 seasonal employees (during the various harvesting periods throughout the year). All permanent employees are provided with housing, running warm water and electricity. There are dedicated permanent housing units. All employees are further provided with personal protective equipment when appropriate, while support is provided in terms of education, etc. Limited contractors are used as the Proponent's focus is to provide employment as well as to build and equip their own workforce with knowledge and skills related to the various components of operations.



Photo 4-29 General employee ablution



Photo 4-30 Mobile toilet in operational areas



Photo 4-31 Employee football field



Photo 4-32 Employee vegetable garden

5 ALTERNATIVES

The Proponent has incorporated various possible revenue generating activities on the property to ensure a robust and sustainable operational unit. A combination of agriculture and related activities are implemented, thereby significantly reducing possible feasible alternatives. Alternatives considered and described below, relate mostly to the implementation of the various project components but also include:

- ◆ Location alternatives;
- ◆ Project implementation and design alternatives;
- ◆ No-go alternative.

5.1 LOCATION ALTERNATIVES

The location of the irrigation areas is well suited for crop production due to the availability of water and suitability of soils. Boreholes are already in place and land clearing and field establishment have already been completed for existing operations. The central and northern portions of the site are clay rich area which are not suited for crop cultivation. It should also be noted that the area used for cultivation has been indicated as crop fields in topographic maps as far back as 1975. Therefore, the area is a known crop producing unit for almost 50 years. No location alternatives are therefore considered feasible, as the Proponent owns the property on which operations are conducted and proposed.

5.2 PROJECT IMPLEMENTATION AND DESIGN ALTERNATIVES

Various alternatives are continually considered to optimise crop production an irrigation. Boreholes are already in place and no surface water is available. Therefore, there are no alternative water sources for the irrigation operations. However, there are a number of alternatives with regards to the application of the water used. The most pertinent relates to crop irrigation methods.

5.2.1 Irrigation Methods

When considering alternative irrigations systems, the most viable irrigation option is not only based on the irrigation system's design efficiency, but should include environmental constrains and operating costs. Some systems are simply not viable due to climatic and topographical features as well as cost implications. For example, flood irrigation is not viable on steeper gradients and are more expensive due to water pumping costs.

The type of produce cultivated also plays a determining role. It will not be feasible to install highly efficient yet expensive irrigation systems (such as drip irrigation) for crops with lower economic yields. In turn, some crops will not produce such high yields when cultivated under less efficient systems. Table 5-1 depicts different types of irrigation systems as per the South African Irrigation Institute's suggested efficiencies (IWRM Plan Joint Venture Namibia, 2010). The estimated average costs are based on 35 ha units and although outdated estimates are still useful for comparisons purposes. Although flood systems are not viable irrigation methods, these have been included for comparison with regards to capital cost and design efficiency.

Table 5-1 Irrigation system efficiency (IWRM Plan Joint Venture Namibia, 2010)

| Irrigation System | Design Efficiency | Capital Costs (R /ha) |
|-----------------------------------|--------------------------|------------------------------|
| Flood: Furrow | 65% | 13,000 |
| Flood: Border | 60% | 17,600 |
| Flood: Basin | 75% | 18,800 |
| Sprinkler: Dragline | 75% | 24,800 |
| Sprinkler: Quick-coupling | 75% | 22,500 |
| Sprinkler: Permanent | 85% | 34,500 |
| Sprinkler: Travelling boom | 80% | 23,200 |
| Sprinkler: Centre pivot | 85% | 43,300 |
| Sprinkler: Linear | 85% | 69,400 |
| Sprinkler: Micro sprinkler | 85% | 36,300 |
| Micro: Spray | 90% | 53,200 |
| Micro: Drip | 95% | 46,300 |

In the Abenab district, climatic and soil conditions necessitate an irrigation system with a high rate of water deposition (due to evaporation and soil salinization). For purposes of irrigation, centre pivot and sprinkler systems are suitable. All irrigation is adjusted and implemented according to rainfall. During higher rainfall periods, less water is irrigated.

5.2.2 Soil Preparation

Traditionally, soil is prepared for planting by tilling and ploughing. These processes break the top layer of soil at varying depths and mix residual plant material into the soil. It also uproots weeds and provide for loose soil. There is nowadays however a shift in the approach to soil preparation that has some advantageous over traditional tilling. Conservation tillage practises aim at less disturbance of the soil and have advantages of less erosion, less evaporation and save on time and costs of traditional tilling. Conservation tillage can either be just partial tillage as is the case with strip-tilling or no tilling at all. With strip-tillage, only narrow strips are tilled in the area where planting will take place. The areas, between planted rows, are left untilled and with residual plant material from the previous harvest. With no-tillage, seeds are planted on the field with no soil preparation at all. The Proponent should investigate the applicability and potential advantages of conservation tillage within the constraints of optimal vegetable growth requirements.

Table 5-2 Advantages and disadvantages of land preparation systems (adapted from <https://cropwatch.unl.edu/tillage/advdisadv>)

| System | Major advantages | Major disadvantages |
|---------------|---|---|
| Plough | Suited for poorly drained soils. Excellent incorporation (mixing of soil for easy combination with chemical and organic elements). Well-tilled seedbed. | Major soil erosion. High soil moisture loss. Timeliness considerations. Highest fuel and labour costs. Reducing soil organic matter (micro flora and fauna), reduced soil structural stability. Increased surface runoff and water or wind erosion. |

| System | Major advantages | Major disadvantages |
|-------------------|---|--|
| Disk | Less erosion with more residue. Well adapted for well-drained soils. Good incorporation. | Little erosion control with more operations. High soil moisture loss. Destroys soil structure. Compacts wet soil. |
| Strip-till | Tilled residue-free strip warms quickly. Injection of nutrients into row area. Well suited for poorly drained soils. Less wear on machinery, less use of fuel or animal power, less time devoted to soil preparation by the farmer thus a possible overall improvement in gross returns for the farm. Heavy rain, is more likely to concentrate in the seeder slots and thereby penetrate directly to the crop's root zone. Improve general water use efficiency by the crop. | Cost of preplant operation. Strips may dry too much, crust, or erode without residue. Not suited for drilled crops (mechanised seeding). Timeliness in wet falls. It also disturbs the soil but limits that disturbance to rows or slots in which the crop seeds and fertilizer are placed. |
| No-till | Excellent erosion control. Soil moisture conservation. Minimum fuel and labour costs. Builds soil structure and health. | No incorporation. Increased dependence on herbicides. Slow soil warming on poorly drained soils. Problems of disease and residue handling. Herbicides have long-term impacts on the environment. Some weeds have developed resistance to some herbicides, leading to a need to rotate both crops and herbicide groups in order to keep crops weed-free, or to plant GMO crops. |

5.2.3 Crop Selection (Maize)

Conventional crop cultivation in Namibia excludes GM maize. Major advantages and disadvantages of traditional non-GM maize and various strains of GM maize are presented in Table 5-3.

Table 5-3 Alternative comparison of maize types for cultivation

| Alternative | Advantages | Disadvantages | Preferred Option |
|--|---|--|--|
| Maize type | | | |
| Traditional non-GM maize and cotton | <ul style="list-style-type: none"> ◆ Long established crops of which the positive and negative properties are well known ◆ Cheaper seeds ◆ Seeds easily available ◆ Can keep some harvested maize seed for next planting season | <ul style="list-style-type: none"> ◆ Highly susceptible to crop damage by insects ◆ Reduced crop yields when significant pest outbreaks occur ◆ Maize is only broad leaf herbicide tolerant ◆ Cotton is only grass herbicide tolerant ◆ More labour intensive ◆ More spraying result in more fuel use and thus greenhouse gasses ◆ Increased water use due to need for dilution of insecticides | <ul style="list-style-type: none"> ◆ Cultivation of GM maize and cotton with traditional maize and cotton as refuges. Planting a combination of GM maize and cotton events, or varying GM maize and cotton events between planting seasons, will contribute to delaying the onset of insect resistance. |
| MON 810 | <ul style="list-style-type: none"> ◆ Resistant to main pests like fall armyworm and African stalk borer ◆ Increased actual yields ◆ Reduced insecticide use ◆ Less labour intensive | <ul style="list-style-type: none"> ◆ Only one BT toxin can potentially lead to more rapid insect resistance to <i>Bacillus thuringiensis</i> ◆ Seed is more expensive ◆ Seed is less easily obtainable | |

| Alternative | Advantages | Disadvantages | Preferred Option |
|--------------------------|--|--|------------------|
| | <ul style="list-style-type: none"> ◆ Less greenhouse gas emissions due to reduced fuel use for spraying ◆ Reduced water use due to less need for dilution of insecticides | <ul style="list-style-type: none"> ◆ Requires special knowledge and proper management to prevent potential negative impacts | |
| MON 89034 (Maize) | <ul style="list-style-type: none"> ◆ Resistant to main pests like fall armyworm and African stalk borer ◆ Two <i>Bacillus thuringiensis</i> toxins has high efficiency and delay insect resistance ◆ Increased actual yields ◆ Reduced insecticide use ◆ Less labour intensive ◆ Less greenhouse gas emissions due to reduced fuel use for spraying ◆ Reduced water use due to less need for dilution of insecticides | <ul style="list-style-type: none"> ◆ Seed is more expensive ◆ Seed is less easily obtainable ◆ Requires special knowledge and proper management to prevent potential negative impacts | |
| NK 603 (Maize) | <ul style="list-style-type: none"> ◆ Easier weed control ◆ Increased actual yields | <ul style="list-style-type: none"> ◆ Weeds can become resistant to glyphosate ◆ Requires special knowledge and proper management to prevent potential negative impacts | |
| Stacked events | <ul style="list-style-type: none"> ◆ Both insect resistance and easier weed control ◆ Increased actual yields ◆ Reduced insecticide use ◆ Less labour intensive ◆ Less greenhouse gas emissions due to reduced fuel use for spraying ◆ Reduced water use due to less need for dilution of insecticides | <ul style="list-style-type: none"> ◆ Pests and weeds can become resistant to Bt proteins and glyphosate ◆ Requires special knowledge and proper management to prevent potential negative impacts | |

5.3 NO GO ALTERNATIVE

Agriculture has been a core activity in the region for years. Maize is supplied to Namibian mills and the stover used for fodder. Staple vegetable production conducted on the farm is of utmost importance to Namibia. Should the project not receive an environmental clearance certificate, there would be a loss in capital investment and a significant loss in employment. This will lead to a decrease in the spending power of the local community. Finally, less revenue will be

generated for Namibia and more money will be required for importing of feed and food. However, the most important aspect of the no go alternatives will be the lack of staple food production for the local market.

6 ADMINISTRATIVE, LEGAL AND POLICY REQUIREMENTS

All projects, plans, programmes and policies with potential adverse impacts on the environment require an environmental assessment, as per the Namibian legislation. This promotes protection of the environment as well as sustainable development. The legislation and standards provided in Table 6-1 to Table 6-3 govern the environmental assessment process in Namibia, and are relevant to the assessed development.

Table 6-1 Namibian law applicable to the development

| Law | Key Aspects |
|--|---|
| The Namibian Constitution | <ul style="list-style-type: none"> ◆ Promotes the welfare of people ◆ Incorporates a high level of environmental protection ◆ Incorporates international agreements as part of Namibian law |
| Environmental Management Act Act No. 7 of 2007, Government Notice No. 232 of 2007 | <ul style="list-style-type: none"> ◆ Defines the environment ◆ Promotes sustainable management of the environment and the use of natural resources ◆ Provides a process of assessment and control of activities with possible significant effects on the environment |
| Environmental Management Act Regulations Government Notice No. 28-30 of 2012 | <ul style="list-style-type: none"> ◆ Commencement of the Environmental Management Act ◆ List activities that requires an environmental clearance certificate ◆ Provides Environmental Impact Assessment Regulations |
| Fertilizers, Farm Feeds, Agricultural Remedies and Stock Remedies Act Act No. 36 of 1947; Government Notice No. 1239 of 1947 | <ul style="list-style-type: none"> ◆ Governs the registration, importation, sale and use of fertilizers, farm feeds, agricultural remedies and stock remedies ◆ Various amendments and regulations |
| Seed and Seed Varieties Act 23 of 2018 Act No. 23 of 2018, Government Notice No. 368 of 2018 | <ul style="list-style-type: none"> ◆ Provides for restrictions on the importation of seed ◆ Not in force yet |
| Water Resources Management Act Act No. 11 of 2013 | <ul style="list-style-type: none"> ◆ Provides for management, protection, development, use and conservation of water resources ◆ Prevention of water pollution and assignment of liability |
| Forest Act (Act 12 of 2001, Government Notice No. 248 of 2001) | <ul style="list-style-type: none"> ◆ Makes provision for the protection of the environment and the control and management of forest fires ◆ Provides for the licencing and permit conditions for the removal of woody and other vegetation as well as the disturbance and removal of soil from forested areas |
| Forest Regulations: Forest Act, 2001 Government Notice No. 170 of 2015 | <ul style="list-style-type: none"> ◆ Declares protected trees or plants ◆ Issuing of permits to remove protected tree and plant species ◆ Issuing of permits for harvesting of trees for wood and charcoal production and transport |

| Law | Key Aspects |
|---|--|
| Soil Conservation Act Act No. 76 of 1969 | <ul style="list-style-type: none"> ◆ Laws relating to the combating and prevention of soil erosion, the conservation, improvement and manner of use of the soil and vegetation and the protection of the water sources in Namibia |
| Biosafety Act Act No. 7 of 2006 | <ul style="list-style-type: none"> ◆ Regulates activities involving the research, development, production, marketing, transport, application and other uses of genetically modified organisms and specified products derived from genetically modified organisms ◆ Prohibits planting of genetically modified organisms without registration |
| Petroleum Products and Energy Act Act No. 13 of 1990, Government Notice No. 45 of 1990 | <ul style="list-style-type: none"> ◆ Regulates petroleum industry ◆ Makes provision for impact assessment ◆ Petroleum Products Regulations (Government Notice No. 155 of 2000) ◆ Prescribes South African National Standards (SANS) or equivalents for construction, operation and decommissioning of petroleum facilities (refer to Government Notice No. 21 of 2002) |
| Local Authorities Act Act No. 23 of 1992, Government Notice No. 116 of 1992 | <ul style="list-style-type: none"> ◆ Defines the powers, duties and functions of local authority councils |
| Public and Environmental Health Act Act No. 1 of 2015, Government Notice No. 86 of 2015 | <ul style="list-style-type: none"> ◆ Provides a framework for a structured more uniform public and environmental health system, and for incidental matters ◆ Deals with Integrated Waste Management including waste collection disposal and recycling, waste generation and storage, and sanitation |
| Labour Act Act No 11 of 2007, Government Notice No. 236 of 2007 | <ul style="list-style-type: none"> ◆ Provides for Labour Law and the protection and safety of employees ◆ Labour Act, 1992: Regulations relating to the health and safety of employees at work (Government Notice No. 156 of 1997) |
| Hazardous Substances Ordinance Ordinance No. 14 of 1974 | <ul style="list-style-type: none"> ◆ Applies to the manufacture, sale, use, disposal and dumping of hazardous substances as well as their import and export ◆ Aims to prevent hazardous substances from causing injury, ill-health or the death of human beings |
| Pollution Control and Waste Management Bill (draft document) | <ul style="list-style-type: none"> ◆ Not in force yet ◆ Provides for prevention and control of pollution and waste ◆ Provides for procedures to be followed for licence applications |

Table 6-2 Guiding documents, directives and standards

| Standard or Code | Key Aspects |
|--|---|
| South African National Standards (SANS) | <ul style="list-style-type: none"> ◆ The Petroleum Products and Energy Act prescribes SANS standards for the construction, operations and demolition of petroleum facilities ◆ SANS 10089-3:2010 is specifically aimed at storage and distribution of petroleum products at fuel retail facilities and consumer installations ◆ SANS 10131 (2004) is aimed at above-ground storage tanks for petroleum products ◆ Provide requirements for spill control infrastructure |
| Department of Water Affairs and Forestry Code of Practice: Volume 1 Septic Tank Guidelines (General Guidelines July 2008) | <ul style="list-style-type: none"> ◆ It defines french drains and septic tanks ◆ Gives location consideration and tank design guidance ◆ Septic tanks are- not allowed between two and five meters from a building and or a boundary ◆ It specifically states that in rocky areas secondary treatment must be provided for soak aways |

Table 6-3 Relevant multilateral environmental agreements

| Agreement | Key Aspects |
|---|---|
| Stockholm Declaration on the Human Environment, Stockholm 1972 | <ul style="list-style-type: none"> ◆ Recognizes the need for a common outlook and common principles to inspire and guide the people of the world in the preservation and enhancement of the human environment |
| United Nations Framework Convention on Climate Change (UNFCCC) | <ul style="list-style-type: none"> ◆ The Convention recognises that developing countries should be accorded appropriate assistance to enable them to fulfil the terms of the Convention |
| Convention on Biological Diversity, Rio de Janeiro, 1992 | <ul style="list-style-type: none"> ◆ Under article 14 of The Convention, EIAs must be conducted for projects that may negatively affect biological diversity |
| International Treaty on Plant Genetic Resources for Food and Agriculture, 2001 | <ul style="list-style-type: none"> ◆ Promotes conservation, exploration, collection, characterization, evaluation and documentation of plant genetic resources for food and agriculture ◆ Promote the sustainable use of plant genetic resources for food and agriculture |

Listed activities, which require an ECC application (Government Regulation No 29 of 2012) related to this project, include the following:

Section 4: Forestry Activities

- ◆ 4 The clearance of forest areas, deforestation, afforestation, timber harvesting or any other related activity that requires authorisation in terms of the Forest Act, 2001 (Act No 12 of 2001) or any other law. Various portions of the farm have previously been cleared (spanning a timeframe of 50 years).

Section 7: Agriculture and Aquaculture Activities

- ◆ 7.4 The import, processing and transit of genetically modified organisms; The Proponent plans to plant GM maize.
- ◆ 7.5 Pest control: The Proponent use conventional pest control products as approved by the Namibian government for some of the produce. These may include herbicides and pesticides and will vary according to season and pests encountered during a year.

Section 8 of Government Notice No. 29 of 2012: Water Resource Developments

- ◆ 8.1. The abstraction of ground or surface water for industrial or commercial purposes: Groundwater is abstracted for current and proposed commercial operations.
- ◆ 8.7 Irrigation schemes for agriculture excluding domestic irrigation: No *irrigation scheme* was developed, however, *irrigation systems* are used on the farm. Irrigation on the farm does not contribute to, or is part of any irrigation scheme, as proclaimed by the Namibian Government.

Section 9 of Government Notice No. 29 of 2012: Hazardous Substance Treatment, Handling and Storage

- ◆ 9.1 The manufacturing, storage, handling or processing of a hazardous substance defined in the Hazardous Substances Ordinance, 1974. Fuel is stored on site for daily operations.
- ◆ 9.2 Any process or activity which requires a permit, licence or other form of authorisation, or the modification of or changes to existing facilities for any process or activity which requires an amendment of an existing permit, licence or authorisation or which requires a new permit, licence or authorisation in terms of a law governing the generation or release of emissions, pollution, effluent or waste. The Proponent store more than 600 l in aboveground storage tanks and has obtained a consumer fuel certificate
- ◆ 9.5 Construction of filling stations or any other facility for the underground and aboveground storage of dangerous goods, including petrol, diesel, liquid petroleum gas or paraffin. Fuel is stored on site, in aboveground storage tanks for daily operations

Additional national planning legislation considered include:

- ◆ National Development Plans (NDPs).
- ◆ Namibia's Climate Change Adaptation

The rationale behind the NDPs is to introduce an element of flexibility within the Ministry planning system by fast tracking development in areas where progress is insufficient. It also incorporates new development opportunities and aims to address challenges that have emerged after the formulation of various NDPs. In the Strategic Plan, the amount of hectares developed for irrigation, is a key performance indicator for the plan's second pillar's strategic objectives, which are aimed:

“to increase productivity during the strategic period through the implementation of appropriate technologies e.g. Comprehensive Conservation Agriculture (CCA) and mechanization in order to ensure food security at both household and national level.”

The above ties in with NDPs which purposes to set out a roadmap for achieving envisioned rapid industrialization while adhering to the four integrated pillars of sustainable development as identified in the plan. Irrigation activities contribute primary to the “Economic Progression” pillar by increasing the volumes of locally produced goods. One of the focus areas of the economic progression pillar of NDPs is agriculture and food security. The NDPs aims to decrease the amount of food insecure individuals, increase food production and increase the share of value addition in crop and livestock farming. Development and operations of irrigation activities on the farm are in line with all of these strategies as identified in the NDPs as well as for the Strategic Plan. The operation contributes to the amount of productive, irrigated land in Namibia, provides employment, and most crucially, produces crops for local markets.

Namibia's Climate Change Adaptation Communication to the United Nations Framework Convention on Climate Change, identifies adaptation actions (amongst others) for the agriculture and water sectors. The Proponent has specifically considered the following actions:

- ◆ Develop improved crop varieties that adapt to climate change (Climate-Resilient Agriculture);
- ◆ Promote the diversification of crops to hedge against erratic rainfall and shorter seasons (Climate-Smart Agriculture); and
- ◆ Improve water demand management, particularly at the local level and in the agricultural sectors.

7 ENVIRONMENTAL CHARACTERISTICS

This section lists pertinent environmental characteristics of the study area and provides a statement on the potential environmental impacts on each.

7.1 LOCALITY AND SURROUNDING LAND USE

The Project is located in the Abenab area of the Tsumeb Constituency, Oshikoto Region in Namibia. The farm is located approximately 32 km east of Tsumeb and 12 km southwest of the small mining settlement of Abenab, centred at 19.210375°S and 18.001444°E. Although there are no active exclusive prospecting licenses (EPLs) registered across the property, EPL applications have been submitted for the base and rare metals, industrial minerals, precious metals and semi-precious stones. There is also an active Petroleum Licence PEL 103 registered across only the eastern portion of the farm.

Surrounding properties are all similar in nature and used for crop cultivation and livestock rearing (commercial farming). No national or proclaimed conservation areas, protected areas or communal conservancies, are located close to the project. The adjacent properties are listed in the table below and their locations are depicted in Figure 7 1.

Table 7-1 Adjacent properties

| Number on Map | Farm Name and/or Number |
|---------------|--------------------------------|
| 1 | Cork FMB/01777 |
| 2 | Detroit FMB/00700 |
| 3 | Deven Port FMB/00702 |
| 4 | Christiana FMB/00705 |
| 5 | Mooindraai FMB/00676 (Buffalo) |
| 6 | Brisbane FMB/01342 |
| 7 | Bombay FMB/00670 |
| 8 | Cooktown FMB/00695 |



Figure 7-1 Properties adjacent to the project area

Implications and Impacts

The location is well suited for the agricultural activities. It is already zoned for agricultural use and is located in an area suitable for irrigation. All buffer zones as required for the cultivation of GM maize should be maintained between the Proponent and neighbours cultivation traditional maize. Consideration should be provided toward prospecting activities proposed across plantations which are not allowed as per the section 1 of the Minerals (Prospecting and Mining) Act 33 of 1992 as amended by the Minerals (Prospecting and Mining) Amendment Act 8 of 2008.

7.2 CLIMATE

The lack of functioning weather stations in Namibia, in especially rural areas, limits the availability of long term, true weather data. In the absence of weather station in the area, the best possible workaround is to use long term climate data obtained from the Atlas of Namibia Project (2002) and the CHIRPS-2 database (Funk et al., 2015), see Table 7-3 and Figure 7-2.

The CHIRPS-2 dataset (Climate Hazards Group Infra-Red Precipitation with Station data version 2) consists of long term rainfall data (1981 to near-present) obtained from satellite imagery and in-situ station data. The remote sensing data provides average rainfall over a 25 km² area and such readings will be lower than some localised actual high volume rain events (thunderstorms). The resultant dataset provides a reasonably well represented overview of the climatic conditions and historic weather conditions of a general area. True values for single, site specific meteorological events may however differ to some degree.

According to the Köppen-Geiger Climate Classification system the project is located in a hot semi-arid climate (BSh) (<http://koeppen-geiger.vu-wien.ac.at/present.htm>). This means that the area receives precipitation below potential evapotranspiration, but not as low as a desert climate and has a mean annual temperature of at least 18 °C. Average rainfall received is 450-500 mm/a with a variation of 30-40 %. Monthly rainfall peaks in January. The potential evapotranspiration is 2,400 – 2,500 mm/a. By dividing the mean annual potential evapotranspiration into the mean

annual precipitation, an aridity index value for the area was computed as 0.2, which indicates the area to be Semi-Arid. The average annual minimum temperature is 6-8 °C, while the average annual maximum temperature is 32-34 °C, with an average annual temperature range of 26-28 °C. An average diurnal temperature (difference between daily minimum and maximum temperature) for this area is around 16-18 °C. Direct normal solar irradiance for the area is 6.574 kWh/m²/day. (Atlas of Namibia Team, 2022)

The rain season normally starts in October and last until April, peaking in January and February. Heavier rainfall (single day events) occur between February and April, with a single event of 67 mm in Apr (last 40 years data) being the highest (Table 7-3). Most of the single day maximums are less than 50 mm. The average annual rainfall for the last 40 years was calculated as 464 mm/a, with a coefficient of variance of 27 %. Daily and seasonal rainfall data (Funk et al., 2015) is presented in Figure 7-2. Seasonal (July to June) total rainfall, centred on the average line for the last 40 years, is presented, with the daily total rainfall and the seasonal cumulative rainfall. From the figure it is clear that six out of the last ten seasons were much drier than usual.


Table 7-2 Summary of climate climatic conditions (Adapted Atlas of Namibia Project, 2002)

| | |
|--|--|
| Köppen-Geiger climate classification | According to the Köppen-Geiger Climate Classification system the project is located in a hot semi-arid climate (BSh) (http://koepen-geiger.vu-wien.ac.at/present.htm). This means that the area receives precipitation below potential evapotranspiration, but not as low as a desert climate and have a mean annual temperature of at least 18°C |
| Average annual rainfall (mm/a) | 450-500 |
| Variation in annual rainfall (%) | 30-40 |
| Average annual evaporation (mm/a) | 2,800-3,000 |
| Water deficit (mm/a) | 1,501-1,700 |
| Average annual temperatures (°C) | 20-21 |
| Average solar radiation (kWh/m²/day) | 6-6.2 |
| Modelled Wind Rose for Abenab (Meteoblue 2023) | |

Table 7-3 Rainfall statistics (Funk et al., 2015)

| Month | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|--------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Minimum (mm) | 12 | 30 | 20 | 5 | 0 | 0 | 0 | 0 | 0 | 0 | 8 | 17 |
| Maximum (mm) | 276 | 211 | 148 | 109 | 5 | 1 | 0 | 0 | 7 | 40 | 100 | 169 |
| Average (mm) | 112 | 109 | 80 | 29 | 1 | 0 | 0 | 0 | 1 | 15 | 41 | 76 |
| Variability (%) | 57 | 44 | 41 | 82 | 245 | 325 | 648 | 0 | 199 | 199 | 52 | 51 |
| Daily maximum (mm) | 44 | 53 | 44 | 67 | 5 | 1 | 0 | 0 | 6 | 19 | 31 | 46 |
| Average rain days | 14 | 12 | 8 | 3 | 0 | 0 | 0 | 0 | 1 | 4 | 7 | 11 |

Season July - June average: 464 mm | Season coefficient of variation: 27 %
Date range: 1981-July-1 to 2023-June-30 | Lat: 19.21037°S; Long: 18.00144°E

**Figure 7-2 Daily and seasonal rainfall (Funk et al., 2015)**

The average annual rainfall for the last 40 years was calculated as 464 mm/a, with a coefficient of variance of 27 %. Daily and seasonal rainfall data (Funk et al., 2015) is presented in Table 7-3. Seasonal (July to June) total rainfall, centred on the average line for the last 40 years, is presented, with the daily total rainfall and the seasonal cumulative rainfall. From the figure it is clear that six out of the last ten seasons were much drier than usual.

Monthly temperature data was retrieved from the Modern-Era Retrospective analysis for Research and Applications version 2 (MERRA-2) data set for a height of 2 m above surface (Ronald Gelaro, et al., 2017). This data set is a NASA atmospheric reanalysis, incorporating satellite data integration and aims at historical climate analyses at 0.5 ° x 0.625 ° spatial resolution. Table 7-4 presents statistics of daily data abstracted from the data set for the last 42 years. Lowest temperature (0.2°C) over the data period was recorded in July, with on average no days in the year being below freezing point. A maximum temperature of the data period of 42.05°C was measured in November.

Table 7-4 Temperature statistics based on Merra-2 data

| Month | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|--------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Minimum (°C) | 11 | 11 | 11 | 8 | 4 | 0 | 0 | 4 | 7 | 8 | 9 | 12 |
| Maximum (°C) | 42 | 41 | 40 | 38 | 35 | 32 | 32 | 36 | 39 | 42 | 42 | 42 |
| Average (°C) | 26 | 26 | 25 | 24 | 21 | 18 | 18 | 21 | 25 | 28 | 28 | 27 |
| Diurnal (°C) | 13 | 12 | 13 | 15 | 17 | 18 | 18 | 19 | 19 | 17 | 15 | 14 |
| Average days < 0°C | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

Implications and Impacts

Rainfall events are often thunderstorms with heavy rainfall that can occur in short periods of time (“cloud bursts”). Rainfall in the area is above the Namibian average, but varies significantly year on year. Heavy rainfall can lead to soil erosion when improper agricultural practises are employed while dry seasons will necessitate greater reliance on groundwater resources. Recurring drought conditions may impact on groundwater availability due to reduced aquifer recharge.

Hot dry winds increase the risk of crop damages as well as fire risks and related severity. General winds may carry chemicals and pollen of crops in mainly a western direction while the solar radiation values are high enough to reliably support future construction of photovoltaic solar panels. Occasional frost necessitate frost management measures. Climate change contributors are largely related to the mechanised systems and synthetic fertilisers used as part of operations. Effects of climate change to consider during the proposed operations over the next 30 years include increased frequency of droughts (changing rainfall patterns) and higher temperatures (World Bank, 2021)

7.3 TOPOGRAPHY, DRAINAGE

Drainage and soils are influenced by topography which, for the project area, can be described as being level to gently undulating with a varying slope of between 0% and 5%. Elevation ranges from between 1,269 mamsl measured at in the most southern portions, to 1,250 mamsl measured around the most northern portions, indicating that elevation dips towards the north. The landscape of the project area is characterised as the Karstveld landscape with surface cover consisting of undifferentiated sediments of the Kalahari Group or more recent deposits. The subsurface geology that outcrop on the farm forms part of the Otavi Mountain Land massif. Prominent outcrops are visible south of the project area.

Drainage is poorly developed in the area. The site falls within the catchment of the Etosha Pan and therefore drainage is towards the north across the project area. The drainage development corresponds with the site elevation. The development of sinkholes, dolines and caves are common in the areas around Abenab, especially west and south-west of the area.




Figure 7-3 Aspect slope and surface drainage

7.4 SOIL

Dominant soil type for this area is Cambic Calcisol which refers to the soil type commonly found in arid or semi-arid regions with dry seasons. They form in calcium and magnesium rich alluvial, colluvial and aeolian deposits and are alternately dampened by rain and dried by evaporation which results in soft masses or hard layers of calcrete.

The farm is situated within the Kalk-2 Agro-Ecological Zone (AEZ) with an average growing period of 91 to 120 days. The Kalk-2 AEZ is ranked 2nd in Namibia in terms of agricultural potential and is deemed most suitable for short-maturing crops and large stock grazing. The Kalk-2 area is generally not regarded as suitable for cropping and this is true for some parts of the farm. The areas under irrigation around Abenab are however located in patches where sufficiently deep, quality soil is present for irrigation of crops.



Photo 7-1 Clayey loamy soils



Photo 7-2 Onions in clayey loamy soils




Figure 7-4 Rock type and Agro Ecological Zone (Atlas of Namibia Project, 2002)

Implications and Impacts

The project area is generally flat and well suited for pivot-based irrigation. The lack of major surface runoff and drainage may lead to pooling and even flooding of plains during heavy rainfall events which may negatively impact soil quality and crop production. However, during dry periods, clayey soils crack open and in some instances causes deep open cavities. In addition, different soil types loose heat at different rates. Loose sandy soils may cool more quickly than heavy, dense clayey soils. Sandy soils therefore have a higher risk of radiation frost.

7.5 GEOLOGY AND HYDROGEOLOGY

The geology underlying the project area formed during the Namibian, Quaternary and Tertiary Age. Locally the geology from the Quaternary and Tertiary Age comprises of the Kalahari Group deposits which consists of sand, calcrete and gravel. The Kalahari Group sediments originate mainly from fluvial deposition with some reworking through aeolian processes. Kalahari sediments at the project location form only a surface cover. The Kalahari Group sediments commonly overlie pre-Kalahari rocks of the Damara Sequence (Namibian Age). At the project location the Damara Sequence consists of dolostones belonging to the Hüttenberg Formation of the Tsumeb Subgroup, which forms part of the Otavi Group (Schneider, 2004). This group belongs to the Northern Platform of the Damara Sequence.

Moderate folding of the strata occurred during the Pan African Orogeny (680-450 Ma) and resulted in the formation of synclines and anticlines, generally trending east - west. To the north of the project area is the Aalborg Syncline, with the Adelaide Syncline to the south. The development of joints and fractures in the rocks are associated with the folding, which have an impact on the hydrogeological characterization of the area.

Various northeast striking magnetic dykes are known to be present in the subsurface, as inferred from aeromagnetic data. The dykes seems to be related to the Paresis intrusion which are situated just south of Otjiwarongo, with dykes radiating from this intrusion. These dykes are locally thought to have shattered the host rocks during its formation. Where dolomite is the host rock, it forms a zone favourable for the development of karst features and groundwater accumulation. At least one of these dykes cross the project area.

Geophysical-interpreted dykes occur in the area and strike towards the northeast. The Remnant dyke intersects the project area. The nature of these dykes tend to be mineralised faults with high hydraulic conductivity values. Both the Tsumeb (further to the west) and Remnant dykes represented a major exploration target for the NamWater exploration water supply programme to Windhoek. The dykes are thought to have shattered the host rocks during its formation (Hoad, 1992). Where dolomite is the host rock, it forms a zone favourable for the development of karst features and groundwater accumulation.

Several known karst features are present in the region. The most well-known of these are the mineralised karst chimney of Abendab Mine ((Bäumle, 2003), as well as several lakes (Otjikoto and Guinas) and caves (e.g. Ghaub).

The Abenab and Abenab West Mines are located approximately 9 km to the southeast of the project area. This hydrothermal deposit represents a highly mineralized zone of which metals like vanadium as well as lead, copper and zinc were mined until 1948 and 1958 respectively when the ore reserves were depleted and the mines were closed (von Bezing, et.al., 2014). The Tsumeb Mine is approximately 29 km to the west; mining ceased in 1994.

A number of springs are present in the Otavi Mountain land and most of these springs are related to the contact zones between relatively impermeable formations of the Grootfontein Metamorphic Complex and more permeable formations of the Damara Sequence. The nearest of these contact zone springs is present approximately 11 km to the southeast of the project farm.

The project area is situated in the Owambo Groundwater Basin. Localised groundwater flow may take place along preferred flow paths in different directions, but the larger scale groundwater flow is expected to be in a northern direction (Figure 7-5). Local flow patterns may vary due to

groundwater abstraction. Groundwater flow is expected to take place through primary porosity in the surface cover, while it is expected to flow along fractures, faults, dykes/mineralised faults or along contact zones (secondary porosity) and other geological structures present within the underlying formations (hard rock formations). Contact zones in the area occur between permeable and impermeable formations and create favourable conditions to promote groundwater flow.





Figure 7-5 Groundwater catchments and water control areas

Table 7-5 indicates the groundwater statistics for a radius of 5 km within the study area. The groundwater information was obtained from Department of Water Affairs (DWA) borehole database. The DWA database is generally outdated and more boreholes might be present. Groundwater is widely utilised in the study area, with 11 boreholes within a 5 km radius. The boreholes were drilled to an average depth of 71 m below surface and average yield of 9 m³/h.

Groundwater quality falls mainly under Group A category, which indicates that the water is of an excellent quality, based on the provided parameters.

Table 7-5 Groundwater statistics

| Query Centre: Cuxhaven; -19.2104°S; 18.0014°E | | Query Box Radius: 5.0km | | | | | | | | | | |
|---|--|---------------------------|------------|-----------|-------------|---------------------------|-------------------|--------------------|-----------|----------------|---------------|----------------|
|  | | NUMBER OF KNOWN BOREHOLES | LATITUDE | LONGITUDE | DEPTH (mbs) | YIELD (m ³ /h) | WATER LEVEL (mbs) | WATER STRIKE (mbs) | TDS (ppm) | SULPHATE (ppm) | NITRATE (ppm) | FLUORIDE (ppm) |
| Data points | | 11 | | | 8 | 8 | 8 | 4 | 10 | 10 | 10 | 10 |
| Minimum | | | -19.165404 | 17.953751 | 46 | 3 | 21 | 30 | 477 | 6 | 0 | 0 |
| Average | | | | | 71 | 9 | 35 | 42 | 586 | 10 | 6 | 0 |
| Maximum | | | -19.255396 | 18.049049 | 119 | 18 | 46 | 50 | 789 | 28 | 14 | 0 |
| Group A | | | | | 12.50% | 25.00% | 0.00% | 0.00% | 100.00% | 100.00% | 60.00% | 100.00% |
| <i>Limit</i> | | | | | 50 | >10 | 10 | 10 | 1000 | 200 | 10 | 1.5 |
| Group B | | | | | 75.00% | 37.50% | 100.00% | 75.00% | 0.00% | 0.00% | 40.00% | 0.00% |
| <i>Limit</i> | | | | | 100 | >5 | 50 | 50 | 1500 | 600 | 20 | 2.0 |
| Group C | | | | | 12.50% | 37.50% | 0.00% | 25.00% | 0.00% | 0.00% | 0.00% | 0.00% |
| <i>Limit</i> | | | | | 200 | >0.5 | 100 | 100 | 2000 | 1200 | 40 | 3.0 |
| Group D | | | | | 0.00% | 0.00% | 0.00% | 0.00% | 0.00% | 0.00% | 0.00% | 0.00% |
| <i>Limit</i> | | | | | >200 | <0.5 | >100 | >100 | >2000 | >1200 | >40 | >3 |

Statistical grouping of parameters is for ease of interpretation, except for the grouping used for sulphate, nitrate and fluoride, which follow the Namibian guidelines for the evaluation of drinking-water quality for human consumption, with regard to chemical, physical and bacteriological quality. In this case the groupings has the following meaning:

Group A: Water with an excellent quality

Group B: Water with acceptable quality

Group C: Water with low health risk

Group D: Water with a high health risk, or water unsuitable for human consumption.

Implications and Impacts

A risk to groundwater pollution is expected due to the geological sensitivity of the area. Groundwater is utilized in the area and such users would be at risk if groundwater contamination occurs. Irresponsible irrigation methods like over-irrigation may result in higher demands for fertiliser and pesticide which in turn will increase nitrates and pesticide concentration in the groundwater. Over application of the herbicide RoundUp™ on is specifically a common expressed concern when planting RoundUp™ ready maize.

Over abstraction may also impact on other users of the aquifer. The hydrogeological specialist study however indicates that water levels, under current groundwater abstraction rates, are stable.

7.6 PUBLIC WATER SUPPLY

The Proponent and surrounding farming communities are completely reliant on groundwater as a source of potable water supply. The boreholes tap into the Owambo Basin and are located within the Tsumeb-Otavi-Grootfontein Subterranean Water Control Area. The most northern portions of the NamWater Waterberg Water Supply Area are located approximately 30 km south-east of the project area and form part of the Otjituuo Scheme. The Berg Aukas Water Supply Scheme is located a further 36 km south-east.

Implications and Impacts

Groundwater is a valuable resource in the farming area and is controlled by a water abstraction permit system as regulated by the Ministry of Agriculture, Water and Land Reform. Groundwater contamination may negatively impact surrounding boreholes. No alternative water supply options exist if extensive contamination or deterioration of groundwater occur. Water abstraction schemes may affect the project which is located downstream of such schemes.

7.7 ECOLOGY

The project area falls within the Savanna Biome with a Karstveld vegetation type and Woodland structure. Namibia's biodiversity pattern is characterised by low species diversity, but high endemism, in the west and southwest of the country, while high species diversity, but low levels

of endemism, is present towards the northeast. Endemism in and around the farm is medium to low.

Plant diversity is expected to be in the vicinity of 400 to 500 species, the second highest diversity category for Namibia. Trees such as *Colophospermum mopane*, *Terminalia prunioides*, *Commiphora species*, *Combretum apiculatum*, *Acacia reficiens*, *Dichrostachys cinerea* and a variety of other trees are characteristic of the Karstveld vegetation type. (Table 7-6). The farm is located in in quarter degree square 1918AA. According to the Tree Atlas Project, 74 different tree species occur in this quarter degree (Curtis & Mannheimer 2005). A summary of those trees protected by legislation in Namibia, 12 in total, is presented in Table 7-7.

Not all the trees listed are expected to occur within the vicinity of the farm. A complete tree list is presented in Appendix D.

Table 7-6 General plant data (Atlas of Namibia Project, 2002)

| | |
|-------------------------------------|---|
| Biome | Savanna |
| Vegetation type | Karstveld |
| Vegetation structure type | Woodland |
| Diversity of higher plants | High (Diversity rank = 2 [1 to 7 representing highest to lowest diversity]) |
| Number of plant species | 400 - 500 |
| Percentage tree cover | 11-25 |
| Tree height (m) | 2-5 |
| Percentage shrub cover | 51-75 |
| Shrub height (m) | 1-2 |
| Percentage dwarf shrub cover | 2-10 |
| Dwarf shrub height (m) | < 0.5 |
| Percentage grass cover | 26-50 |
| Grass height (m) | < 0.5 |

Table 7-7 Trees with conservation concerns in quarter degree squares 1918AA (Curtis & Mannheimer, 2005)

| Name | Common Name | Notes |
|------------------------------|-----------------------|---|
| <i>Acacia erioloba</i> | Camel-thorn | Protected by forestry legislation. |
| <i>Albizia anthelmintica</i> | Worm-cure Albizia;Aru | The low numbers of young trees recorded is a concern, as is the number of dead trees in some areas. It is protected by forestry legislation. |
| <i>Aloe littoralis</i> | Windhoek Aloe | Potentially threatened by pachycaul trade. Protected by the Nature Conservation Ordinance and listed in CITES Appendix II. |
| <i>Berchemia discolor</i> | Bird Plum | This species is protected by forestry legislation, as well as by traditional Owambo cultures for its fruit and shade. |
| <i>Boscia albitrunca</i> | Shepherd's Tree | Although widespread and hardy, it is heavily utilised by people and animals. Protected by forestry legislation. |
| <i>Combretum imberbe</i> | Leadwood | Although heavily utilized by people regrowth is good and growth of young trees is vigorous. Because of its religious importance and many uses, it is protected locally. Old specimens warrant protection as monuments. Protected by forestry legislation. |
| <i>Erythrina decora</i> | Namib Coral-tree | Endemic to Namibia and very uncommon throughout its range. Worthy of protection as there are very few young trees. Protected by forestry legislation. |
| <i>Hyphaene petersiana</i> | Makalani Palm | Should be monitored due to extensive harvesting. Protected by forestry legislation. |
| <i>Maerua schinzii</i> | Ringwood Tree | Increasingly impacted by humans and giraffes. Protected by forestry legislation. |
| <i>Sclerocarya birrea</i> | Marula | Protected locally by communities that use them. Protected by forestry legislation. |

| Name | Common Name | Notes |
|------------------------------|---------------|------------------------------------|
| <i>Spirostachys africana</i> | Tambooti | Protected by forestry legislation. |
| <i>Ziziphus mucronata</i> | Buffalo-thorn | Protected by forestry legislation. |

Some caves, mainly southwest of the area, present suitable habitats for a number of bat species which have been documented to range across the project area. These bats, all of which are listed as least concerned, according to the IUCN Red list of threatened species, include the following species: Dent's Horseshoe Bat (*Rhinolophus denti*) and the Greater Long-fingered Bat (*Miniopterus inflatus*). The project area further falls within the habitat for a number of additional species of concern which may occur within the area. Some of the IUCN Red List of threatened species which are more likely to occur on site are listed in Table 7-8.

Table 7-8 IUNC Red listed species which may occur in the area

| Species Name | Common Name | IUCN Red List Status |
|---------------------------------|----------------------|----------------------------|
| <i>Sagittarius serpentarius</i> | Secretarybird | Threatened |
| <i>Torgos tracheliotos</i> | Lappet-faced Vulture | Endangered |
| <i>Falco vespertinus</i> | Red-footed Falcon | Vulnerable |
| <i>Neophron percnopterus</i> | Egyptian Vulture | Endangered (Breeding area) |
| <i>Aquila nipalensis</i> | Steppe Eagle | Endangered |
| <i>Acinonyx jubatus</i> | Cheetah | Vulnerable |
| <i>Gyps africanus</i> | White-backed Vulture | Critically Endangered |
| <i>Trigonoceps occipitalis</i> | White-headed Vulture | Critically Endangered |

The probability of some of the species in Table 7-8 occurring on site is very likely, mainly due to wildlife roaming freely and undisturbed on areas of the farm which may present preferred habitation areas. These areas are protected by the Proponent and aim at providing all and any wildlife with a secure area, away from noisy and or disturbing activities. Various antelope species, predators and large game are known to be present on the farm. Since the property borders farming operations who also have their own "conservation" (less disturbed) areas, an ecological corridor exist between them and the Proponent which see some species crossing to and fro. These include antelope species such as kudu and oryx, but also include predator species. Jackal, caracal, leopard and cheetah are known to be on and around the farm.



Photo 7-3 Wildlife scat



Photo 7-4 Termite mound on site

Implications and Impacts

Pollution of the soil and groundwater by hazardous chemicals and/or the excessive use of fertilizers and pesticides may negatively impact the local ecology. Irresponsible use of pesticides to kill vermin such as jackal may further impact on already threatened vulture populations as well as other scavengers. Pesticides may also magnify (biomagnification) in higher trophic levels, especially top predators. This may lead to reproductive and other physiological defects and ultimately declining populations. Over-abstraction of groundwater may lead to ecosystem changes as groundwater levels decrease, which may have direct impacts on especially cave habitats downstream (towards Etosha).

Planting of GM maize without implementing the necessary refuges, and not implementing monitoring programmes and preventative and mitigation measures when needed, may result in insect and weed resistance development. This may potentially impact the local ecosystem

structure. Concerns related to the killing of non-target insects as a result of planting insect resistant maize are addressed in the specialist report (Appendix C).

7.8 LOCAL ECONOMY

The Oshikoto Region's economy is a diverse representation of various sectors and industries within the region. These include (but are not limited to) mining, tourism and agriculture; all of which have shown potential to be developed. The rural parts of the Tsumeb constituency has much less economic diversity and the agricultural sector, specifically the irrigation schemes between Tsumeb and the project area, are large economic contributors, if not the largest in the constituency. Not only does it create jobs, but it has also been one of the main driving forces of infrastructure development and related capital expenditure which are on-going in planning considerations. Continued employment increases individual economic resilience and provides for increased social security benefits.

In evaluating water use in primary economic activities such as agriculture, it is useful to consider the entire value-chain, i.e. the upstream and downstream activities. Intensive irrigated production schemes are strong economic drivers, as witnessed by the influx of workers to such areas. This potentially make a significant contribution to the development of the Oshikoto Region and Namibia as a whole.

Water quality will have an effect on the productivity of operations, therefore the economic benefits of ensuring that the water quality and quantity of the groundwater reserve remains at its best, is an essential component of the agricultural process. If water treatment is required, then the cost of production will increase, resulting in a decrease in revenue and feasibility. The same can be said for the quality of the soil, as lowered quality soil will be less economically productive and contaminated soil, such as found in some areas within the constituency, not usable at all. Water and soil are paramount for the continued functioning of the agricultural project and therefore provide a vital ecosystem service to the Proponent.

Regionally, skilled agriculture and fishery provide the most employment (48%). It should be noted that although fishery falls within the agriculture sector it does not contribute to employment in the Oshikoto region. This trend follows through into the rural areas of the Tsumeb constituency within which the project lies. The economy of the area relies largely on commercial livestock farming supplemented with crop production and charcoal manufacturing. Livelihoods in the constituency are varied engaging various sectors such as construction, wholesale and retail, administrative (public and defence) and manufacturing.

Table 7-9 Main industry of employed population aged 15 years and above for the Tsumeb Constituency and Oshikoto Region

| Main Industry | Tsumeb Constituency Rural | Oshikoto Region |
|---|----------------------------------|------------------------|
| Total | 29,931 | 36,638 |
| Agriculture Forestry and Fishing | 17,535 | 17,860 |
| Mining And Quarrying | 195 | 929 |
| Manufacturing | 566 | 1 123 |
| Electricity Gas Steam and Air conditioning supply | 17 | 44 |
| Water Supply Sewerage Waste Management and Remediation activities | 25 | 58 |
| Construction | 1,156 | 1,713 |
| Wholesale and Retail trade; Repair of motor vehicles and motorcycles | 1,229 | 1,880 |
| Transportation and Storage | 619 | 997 |
| Accommodation and Food Service activities | 707 | 963 |
| Information and Communication | 86 | 134 |
| Financial Insurance Activities | 134 | 279 |
| Real estate Activities | 2 | 3 |
| Professional Scientific and Technical activities | 188 | 260 |

| Main Industry | Tsumeb Constituency Rural | Oshikoto Region |
|--|--|----------------------------|
| Administrative and Support service activities | 1,262 | 2,435 |
| Public Administration and Defence; compulsory social security | 1,202 | 1,464 |
| Education | 1,945 | 2,285 |
| Human Health and Social work activities | 762 | 975 |
| Arts Entertainment and Recreation | 69 | 88 |
| Other Services activities | 535 | 682 |
| Activities of Private Households | 1,622 | 2,229 |
| Activities of extraterritorial organisation and bodies | 5 | 8 |
| Don't Know | 70 | 229 |

The project provides significant employment opportunities. Especially during harvesting, which could see over 150 people employed. Since crops are planted on a rotational basis, employment varies throughout the year, Skills and training are required to maintain and operate the irrigation systems. All employees reside on the farm and have received varied skills training ranging from drivers licenses to skills transfer from professionals which are periodically employed at the farm.

Statistics for the demographic profile of the specific area have proven to be lacking in reliable and recent data. In the absence of recent government records, statistics for an area similar in agricultural operations were considered. During a Social Impact Assessment conducted for a similar agricultural project (Bosman, 2017), it was found that every employed person supports approximately six economically inactive persons.

Implications and Impacts

Operations on the farm sustain valuable full time as well as seasonal employment opportunities in a constituency which relies on the agricultural sector. The project contributes to the local and national agricultural sector and specifically in terms of the planned growth in the irrigation sector as envisioned by the local government. Employment and remuneration of such a large workforce within the area stimulates additional economic growth.

The addition of GM maize cultivation, will increase the knowledge of a part of the workforce in terms of the specific requirements linked to GMOs. On a national level, the potential increased yields of GM maize will increase food security during, for example, the outbreak of fall armyworm outbreaks. In considering Round-Up Ready maize, the cultivation thereof can lead to a reduced use of chemicals and tillage, contributing to preserving soil health. However, concerns have been raised about the impact GMO maize may have on on-GMO farmers as well as the export beef industry. These and additional concerns related to GM maize, are discussed in detail in a specialist report (Appendix C).

7.9 DEMOGRAPHIC PROFILE

The project area is located in the Tsumeb magisterial district in the Tsumeb Constituency of the Oshikoto Region and borders the Grootfontein Constituency of the Otjozondjupa Region. Goods and services are mainly sourced from either Tsumeb or Grootfontein. For demographic information of the 2011 population and housing census, refer to Table 7-10 (Namibia Statistics Agency, 2011) which includes the details for the Tsumeb Constituency in relation to the National and regional averages. However, although the project falls within the Tsumeb Constituency, the nature of the area is rural. Unemployment in the Tsumeb Constituency is lower at 34.9% compared to the national and regional averages while the literacy rate also is lower.

Table 7-10 Demographic characteristics of the Tsumeb Constituency, the Oshikoto Region and Nationally (Namibia Statistics Agency, 2011)

| | Tsumeb Constituency | Oshikoto Region | Namibia |
|---------------------------------|--------------------------------|----------------------------|----------------|
| Population (Males) | 11,794 | 87,066 | 1,021,912 |
| Population (Females) | 11,475 | 94,907 | 1,091,165 |
| Population (Total) | 23,269 | 181,973 | 2,113,077 |
| Unemployment (15+ years) | 34.9%* | 40.2%* | 37% |
| Literacy (15+ years) | 84% | 88% | 89% |

* Calculated as per the economically active segment of the population

Seasonal in-migration is a well-established demographic process in the area and also for the project. At times the project has experience a surplus of seasonal migrants which cannot be accommodated through existing operations. Additional persons residing on the properties increase pressure on service provision and may increase the possibility and occurrence of social ills. However, all employees are provided with housing with electricity, clean drinking water and sanitation. The Proponent is further assisting employees through the following means:

1. Recreational infrastructure establishment (football field),
2. A mobile registered clinic and vaccination services visits the farm periodically, and
3. Transportation of children during long weekends and holidays from and to schools located further away.

The Proponent regularly liaise with governing authorities to ensure clinic services for employees. Emergency services are facilitated if and when required between the Proponent and the related governmental agency.

Implications and Impacts

The project contributes mainly to demographic processes indirectly in requiring seasonal employment. Temporary migration in the area changes the demographic profile of the Project as well as the surrounding area. Employment of so many people in a rural area works against urbanisation of the surrounding sectors. Skills development, training and exposure to best practises in terms of wildlife management, irrigation and tourism, benefit employees during the operational phase over and above having access to economic resources and food. Increased access to such resources may increase the fertility of the local population. The concentration of the workforce requires planning of governmental services (such as education clinics and public services) to ensure adequate resources.

Diversifying farming activities by adding the cultivation of GM maize increases the economic resilience of the farming unit and thus provides increased job security to employees.

7.10 CULTURAL, HERITAGE AND ARCHAEOLOGICAL ASPECTS

There are no cultural or heritage aspects known to be present on the farm. The proximity of the farm to Tsumeb, allows for easy integration to cultural and related services for employees. The greater area has been cited to contain a number of caves and dolomite cavities which have been studied for, not only the unique habitats they present, but also the geological evidence related climate.

The Project is further located approximately 10 km from the small mining settlement of Abenab which has been noted to be of historic and geological significance with some highest-grade deposit of vanadate ore in the world, mined about 100 years ago. The largest vanadium crystal in the world (12cm in length) was found near the Abenab mine (Schneider, 2008).

Implications and Impacts

Existing and proposed areas of operations are not close to any caves or related features. However should any archaeological resources be found, such resources should be reported for investigation. Over abstraction of groundwater should be avoided to ensure no water bearing caves downstream of operations area impacted by dewatering.

8 PUBLIC CONSULTATION

Consultation with the public forms an integral component of an environmental assessment investigation and enables interested and affected parties (IAPs) e.g. neighbouring landowners, local authorities, environmental groups, civic associations and communities, to comment on the potential environmental impacts associated with projects and to identify additional issues that they feel should be addressed in the environmental assessment.

Public participation notices were advertised, twice in two weeks, in the national papers: The notices appeared in the Republikein and the Namibian Sun on 15 and 22 March 2022. A site notice was placed on site and notification letters were hand-delivered or e-mailed to neighbours as well as the relevant ministries. See Appendix E for proof of the public participation processes and registered IAPs.

9 MAJOR IDENTIFIED IMPACTS

A number of potential environmental impacts were identified during the environmental scoping exercise. The following section provides a brief description of the most important of these impacts.

9.1 SOIL AND GROUNDWATER CONTAMINATION

Soil and groundwater contamination are possible when large quantities of fertilizers or pesticides are applied. Excessive fertilizer use may result in increased soil nutrient levels (i.e. nitrogen, phosphorus and potassium), to a point that soil is regarded as contaminated. Similarly, pesticides can accumulate in soil at levels detrimental to biota. Fertilizers and pesticides can leach deeper into the ground and eventually reach and contaminate groundwater. Chemical spills, inclusive of fertilizers and pesticides, may result in very high but localised contamination of soil, increasing the risk of groundwater if spill clean-up is not performed.

Evidence shows that, when GMOs are cultivated responsibly, the volumes of pesticides used are lower for GM crops, especially for insecticides.

Hydrocarbon pollution resulting from the spilling of fuel, oil or hydraulic fluids is possible. Tractor and other vehicle breakdowns or incorrect refuelling and storage of fuel are the most likely causes of hydrocarbon pollution. Contamination of the environment may also occur if hazardous substances such as pesticides are stored or handled incorrectly and a spill occur. In addition, incorrectly managed or constructed french drains have the potential to contaminate groundwater resources.

9.2 GROUNDWATER ABSTRACTION

For a detailed assessment of groundwater abstraction and supply, refer to the hydrogeological assessment in Appendix B.

Groundwater abstraction is a very sensitive topic in a dry country where the value of land is drastically reduced if no or poor quality groundwater is present on the land. Abstraction of groundwater must be done in a sensible way not to impact on other groundwater users that depend on such groundwater. This includes water abstracted for human and animal use, irrigation, and also ecosystems that depend on groundwater. A typical groundwater balance was compiled to illustrate the potential consequences of over abstraction of groundwater, see Figure 9-1. Recharge to the area is considered to be high. It is considered that recharge can vary from 0% to 4% of rainfall with an average of 2% of the rainfall. In periods of drought there may be no recharge while in above average rainfall recharge could be above 4% (Hoad, 1992).

In a typical groundwater environment, a water balance would consist of inflow and outflow of the groundwater system. Over time an equilibrium (or steady state) is normally reached with rising water tables following good recharge events and declining water tables when recharge is below average.

Inflow into the system would typically be from infiltration following rainfall in the area and in upstream areas. The inflow component will further be enhanced by the high secondary porosity nature of the karst aquifer.

Outflow would be comprised of water leaving the system through springs and as outflow over the lower boundary of the groundwater system as well as evapotranspiration losses. Groundwater abstraction from boreholes is important as this is normally necessary to sustain human and animal demands where such users became essentially dependant on the abstracted groundwater as a reliable and sustainable source.

Typical consequences of over abstraction will include a lowering in the water table. This may lead to the collapse of underground cave roofs where the hydrostatic pressure, used to support the roof of a cave, decrease. The increased flow of water may enhance the dissolution of dolomitic rock, leading to an increase in karst structures. Lowering of water tables may further lead to the drying up of boreholes, springs, underground caves and the subsequent loss of organisms that lives in the subsurface and surface water. Vegetation will also be impacted where such vegetation has access to groundwater.

Based on current water level fluctuations in the area, as presented in the hydrogeological assessment (Appendix B), a short term threshold of 5 m below the long term average water level is set from where abstraction rates should be reduced. Note that this level refers to rest water levels and not pump water levels.

All boreholes should be equipped with a dipper pipe to enable safe water level measurements.




Figure 9-1 Conceptual groundwater balance with over abstraction scenario

9.3 FIRE

A risk of veld fires exist. Fires, used for example to cook food in areas not designated for this purpose, may spread to the nearby veld. Machinery can ignite dry vegetation if sufficient heat (e.g. exhaust pipes) or sparks are produced. Chemicals and fuels stored and used for general activities may be flammable. Electrical shorts on the electricity supply network can cause fires in buildings. Lightning can be a natural ignition source for veld fires which in turn can spread and damage infrastructure and crops or pose health impacts.

9.4 DUST AND AIR QUALITY

Dust may become a nuisance and health risk when land is ploughed, tilled or prepared for planting. Strong winds present during periods when fields are dry and barren, such as in-between planting cycles, may aggravate dust impacts.

9.5 TRAFFIC

Additional traffic is present on the main road as a result of the activities on the farm. This include the transport of staff, the delivery of fertilizers, seed, etc., as well as the transport of crops to markets. Since it is an existing operation with no short term plans for expansion, traffic impacts related to the activities of the Proponent will remain the same, and no additional impacts are expected.

9.6 HEALTH AND SAFETY

Injuries related to working with machinery, chemicals, pesticides, etc. can occur. Inhalation and dermal contact with pesticides are possible where pesticides are for example applied by means of tractor mounted sprayers or via the irrigation system. Spray drift in windy conditions can reach nearby workers or the tractor driver. Vehicle accidents involving staff during operations for instance when operating a tractor. Venomous animals like snakes, scorpions and spiders may be present.

9.7 ECOSYSTEM AND BIODIVERSITY IMPACT

No additional land clearing is foreseen in the near future. Poaching and illegal collection of plant and animal material by staff and / or non-staff members is possible. Pollution of the environment and groundwater, especially by fuel, pesticides and fertilizers, can deteriorate or alter the ecosystem structure and function. Over-abstraction of groundwater can impact on aquatic organisms living in the groundwater. Due to the endemic nature of such aquatic organisms, the continued pumping of water that results in the extraction of for example amphipods, may detrimentally affect population sizes and viability.

9.8 PESTICIDE RESISTANCE

In GM crop fields, pesticide resistance has been reported in insects (against Bt proteins) and weeds (against glyphosate). This is however not deferent from pesticide resistance reported in non-GM crop fields. Over reliance on the use of glyphosate, the lack of crop and herbicide rotation by farmers and the incorrect use of refuges, in some regions, contribute to the development of weed and insect resistance.

9.9 HORIZONTAL GENE TRANSFER

Horizontal gene transfer (HGT) is considered to have negligible risks to humans and the environment and no cases of adverse environmental effects as a result of HGT between GM crops and wild, related plants have been observed. Personal choice me be affected where GM maize is planted too close to fields of a farmer who does not want GM maize. Pollen from GM maize can travel over short distances which would mean HGT can occur between the GM and non-GM fields.

9.10 SOCIO-ECONOMIC IMPACTS

The project contribute to food security at a national level and meets some of the objectives of the various NDP goals in terms of agricultural productivity. Ten permanent employees and up to 80 seasonal employees work on the farm. Housing and amenities are available to permanent employees and their families. Proper sanitation facilities are present for all workers.

Income is generated with a positive impact on the Namibian GDP. The sale of high value crops to international clients contribute towards a positive trade balance for Namibia. At present, there is no difference in the potential yield between conventional maize and GM variants. However, actual yields for GM variants may be higher due to decreased insect damage, especially during a heavy infestation or plague, and competition with weeds. Coupled to this is the potential for

increased profit margins if reduced volumes of pesticides are used, which also mean less fuel and water consumption. GM seed are typically more expensive and crop producers will likely consider the financial benefits of GM maize and cotton vs. conventional maize and cotton in deciding which to plant.

Existing and planned developments typically entice jobseekers to migrate to the area. This may lead to high levels of unemployment and the social ills therewith associated. This include increased spread of HIV/AIDS and other diseases, alcohol or drug abuse, and theft or violence.

10 ASSESSMENT AND MANAGEMENT OF IMPACTS

The purpose of this section is to assess and identify the most pertinent environmental impacts that are expected from the operational, construction, care and maintenance, and potential decommissioning activities of the farming unit. An EMP based on these identified impacts is presented in this section.

For each impact, an environmental classification was determined based on an adapted version of the Rapid Impact Assessment Method (Pastakia, 1998). Assessment of impacts is based on the following categories: importance of condition (A1); magnitude of change (A2); permanence (B1); reversibility (B2); and cumulative nature (B3) (Table 10-1).

The environmental classification is calculated as follows:

Environmental classification = $A1 \times A2 \times (B1 + B2 + B3)$.

The environmental classifications of impacts and the respective classes are provided in Table 10-2.

The probability ranking refers to the probability that a specific impact will happen following a risk event. These can be improbable (low likelihood); probable (distinct possibility); highly probable (most likely); and definite (impact will occur regardless of prevention measures).

Table 10-1 Assessment criteria

| Criteria | Score |
|---|-------|
| Importance of condition (A1) – assessed against the spatial boundaries of human interest it will affect | |
| Importance to national/international interest | 4 |
| Important to regional/national interest | 3 |
| Important to areas immediately outside the local condition | 2 |
| Important only to the local condition | 1 |
| No importance | 0 |
| Magnitude of change/effect (A2) – measure of scale in terms of benefit/disbenefit of an impact or condition | |
| Major positive benefit | 3 |
| Significant improvement in status quo | 2 |
| Improvement in status quo | 1 |
| No change in status quo | 0 |
| Negative change in status quo | -1 |
| Significant negative disbenefit or change | -2 |
| Major disbenefit or change | -3 |
| Permanence (B1) – defines whether the condition is permanent or temporary | |
| No change/Not applicable | 1 |
| Temporary | 2 |
| Permanent | 3 |
| Reversibility (B2) – defines whether the condition can be changed and is a measure of the control over the condition | |
| No change/Not applicable | 1 |

| | |
|---|---|
| Reversible | 2 |
| Irreversible | 3 |
| Cumulative (B3) – reflects whether the effect will be a single direct impact or will include cumulative impacts over time, or synergistic effect with other conditions. It is a means of judging the sustainability of the condition – not to be confused with the permanence criterion. | |
| Light or No Cumulative Character/Not applicable | 1 |
| Moderate Cumulative Character | 2 |
| Strong Cumulative Character | 3 |

Table 10-2 Environmental classification (Pastakia 1998)

| Environmental Classification | Class Value | Description of Class |
|------------------------------|-------------|-------------------------------|
| 72 to 108 | 5 | Extremely positive impact |
| 36 to 71 | 4 | Significantly positive impact |
| 19 to 35 | 3 | Moderately positive impact |
| 10 to 18 | 2 | Less positive impact |
| 1 to 9 | 1 | Reduced positive impact |
| 0 | -0 | No alteration |
| -1 to -9 | -1 | Reduced negative impact |
| -10 to -18 | -2 | Less negative impact |
| -19 to -35 | -3 | Moderately negative impact |
| -36 to -71 | -4 | Significantly negative impact |
| -72 to -108 | -5 | Extremely Negative Impact |

10.1 RISK ASSESSMENT AND ENVIRONMENTAL MANAGEMENT PLAN

The EMP provides management options to ensure impacts of the agricultural and related activities on the farming unit are minimised. An EMP is a tool used to take pro-active action by addressing potential problems before they occur. This should limit corrective measures needed, although additional mitigation measures might be included if necessary. The environmental management measures are provided in the tables and descriptions below. For the GMO specific management plan please refer to Appendix C. These management measures should be adhered to during the execution of various activities on the farming unit. This section of the report is also presented as a stand-alone document for easy reference. All personnel taking part in the operations of the farm should be made aware of the contents of this section, so as to plan the operations accordingly and in an environmentally sound manner.

The objectives of the EMP are:

- ◆ to include all components related to operational and possible construction activities of the farming unit;
- ◆ to prescribe the best practicable control methods to lessen the environmental impacts associated with the farm;
- ◆ to monitor and audit the performance of operational personnel in applying such controls; and
- ◆ to ensure that appropriate environmental training is provided to responsible operational personnel.

Various potential and definite impacts will emanate from the operations, maintenance/construction and decommissioning phases. The majority of these impacts can be mitigated or prevented. The impacts, risk rating of impacts, as well as prevention and mitigation measures are listed below.

As depicted in the tables below, impacts related to the operational phase are expected to mostly be of medium to low significance and can typically be mitigated to have a low significance. The extent of impacts are largely site specific to local and are not of a permanent nature. Due to the

nature of the surrounding areas, cumulative impacts are possible and the most important of these are potential groundwater and biodiversity/ecological impacts.

10.1.1 Planning

During the phases of planning for the operations, maintenance/construction and decommissioning of the farming unit, it is the responsibility of the Proponent to ensure they are and remain compliant with all legal requirements. The Proponent must also ensure that all required management measures are in place prior to, and during all phases, to ensure potential impacts and risks are minimised. The following actions are recommended for the planning phase and should continue during all other phases of the project:

- ◆ Ensure that all the necessary permits from the various ministries, local authorities and any other bodies that governs the operations, maintenance/construction and decommissioning activities on the farm remain valid. These include the water abstraction permit, consumer installation certificate and permit for environmental release of GM maize.
- ◆ Ensure all appointed contractors and employees enter into an agreement, which includes the EMP. Ensure that contractors, sub-contractors, employees and all personnel present on site understand the contents of the EMP.
- ◆ Make provisions to have a Health, Safety and Environmental (HSE) Coordinator to implement the EMP and oversee occupational health and safety as well as general environmental related compliance.
- ◆ Make provision for a community liaison officer to deal with complaints.
- ◆ Have the following emergency plans, equipment and personnel on site, where reasonable, to deal with all potential emergencies:
 - EMP, risk management plan, emergency response plan and HSE manuals;
 - Adequate protection and indemnity insurance cover for incidents;
 - Procedures, equipment and materials required for emergencies (e.g. firefighting, first aid, etc.).
- ◆ Establish and maintain a fund for future ecological restoration, specifically for instances of environmental damage caused during operations including pollution remediation where required. Should project activities cease completely, and future land-use will not involve agriculture, the funds should be utilised to remove all redundant infrastructure and waste.
- ◆ Establish and/or maintain a reporting system to report on aspects of operations, maintenance/construction, and decommissioning as outlined in the EMP. Keep monitoring reports on file for bi-annual submission to MEFT in support of environmental clearance certificate renewal applications. This is a requirement by MEFT.
- ◆ Appoint a specialist environmental consultant to update the environmental assessment and EMP and apply for renewal of the environmental clearance certificate prior to expiry.

10.1.2 Revenue Generation in the Professional Sector

Consulting and professional services are engaged with for assistance in applications for new permits and renewal for existing permits such as the water permits, fuel storage and environmental clearance certificates. In addition, specialist irrigation systems, pumps and implements used by the agricultural project require specialist and professional services. Such services may further be extended to pest control for operations and accounting and legal services for administrative processes. All of these services are paid for and therefore the agricultural project contributes to revenue generation in the local and national sectors. In addition, during many of these processes, such as per the renewal of water permits, information is generated which informs and facilitates planning of the Proponent as well as affected parties and governmental agencies.

| Project Activity/Resource | Nature (Status) | (A1) Importance | (A2) Magnitude | (B1) Permanence | (B2) Reversibility | (B3) Cumulative | Environmental Classification | Class Value | Probability |
|---------------------------|--|-----------------|----------------|-----------------|--------------------|-----------------|------------------------------|-------------|-----------------|
| Planning | Employment and contribution to local and national economy | 3 | 2 | 3 | 3 | 2 | 54 | 4 | Definite |
| Daily Operations | Contracted services and contribution to local and national economy | 2 | 1 | 3 | 3 | 1 | 14 | 2 | Definite |
| Indirect Impacts | Increased economic resilience in the professional sector | 3 | 1 | 3 | 1 | 1 | 15 | 2 | Highly Probable |

Desired Outcome: Contribution to national treasury and increased economic resilience in the local and national professional sector.

Actions

Enhancement:

- ◆ Contract local Namibians where possible.
- ◆ Adhering to permit conditions on reporting.
- ◆ Deviations from this practice must be justified.

Responsible Body:

- ◆ Proponent

Data Sources and Monitoring:

- ◆ Service providers' contracts or agreements or records be kept.
- ◆ All reporting, monitoring and information sharing records kept on file.

10.1.3 National Development Goals: Water, Agriculture and Land Use Planning

The agricultural project pins down key development goals and challenges which were identified as part of the Namibian development goals. It may be considered as an agricultural / irrigation project which aims at generating income from foreign sectors by providing the most value per resource (water, soil and labour). In addition, the project is located in line with the regional planning initiatives which identified the location as an area for irrigation development. The project will further contribute to the national climate change combatting initiatives through crop diversification and proposed resilient crop cultivation. Developing of the agricultural sector was identified as one of the core plans within the NDPs for Namibia. The agricultural project therefore is considered to be a positive contributor to achieving national development goals.

| Project Activity/Resource | Nature (Status) | (A1) Importance | (A2) Magnitude | (B1) Permanence | (B2) Reversibility | (B3) Cumulative | Environmental Classification | Class Value | Probability |
|---------------------------|--|-----------------|----------------|-----------------|--------------------|-----------------|------------------------------|-------------|-----------------|
| Planning | Project implementation in line with the NDP and regional land use planning | 4 | 1 | 2 | 1 | 1 | 16 | 2 | Highly Probable |
| Daily Operations | Expansion of the agricultural sector in the Region. Project implementation in line with the regional land use planning | 3 | 2 | 2 | 2 | 2 | 36 | 4 | Highly Probable |
| Indirect Impacts | Contributing to achieving the goals set out in Vision 2030 for Namibia | 3 | 1 | 3 | 3 | 3 | 36 | 4 | Highly Probable |

Desired Outcome: Continued contribution to the development of the Region as well as implementation of project activities in line with NDPs and Vision 2030.

Actions

Enhancement:

- ◆ Liaison with regional and national governmental agencies through appropriate financial and social responsibility reporting.
- ◆ Increase recycling initiatives and incorporate additional greenhouse gas reduction activities such as conservation tillage and climate smart agriculture.
- ◆ Infrastructure maintenance and development such as, road servitude, water- and sanitation system developments (provision to employees) and node development. Where possible, public and private partnership regarding projects should be considered.

Responsible Body:

- ◆ Proponent

Data Sources and Monitoring:

- ◆ All project contributions towards regional development, inclusive of communications held with relevant authorities, to be kept on file.
- ◆ Monitoring of borehole water levels and water abstraction (monthly) and submit to the relevant custodian on a quarterly basis.

10.1.4 Skills and Development

During the operations and maintenance/construction phases, some training is provided to a portion of the workforce, to allow them to conduct certain tasks according to the required standard. Skills are transferred to an unskilled workforce for general tasks and irrigation-related operations. Development of people and technology are key to economic development and the success and safety of operations. The Proponent plays a role in promoting and sustaining the agricultural industry in the area.

| Project Activity/Resource | Nature (Status) | (A1) Importance | (A2) Magnitude | (B1) Permanence | (B2) Reversibility | (B3) Cumulative | Environmental Classification | Class Value | Probability |
|---------------------------|--|-----------------|----------------|-----------------|--------------------|-----------------|------------------------------|-------------|-------------|
| Construction | Employment and transfer of skills, technological advancements | 2 | 1 | 2 | 3 | 1 | 12 | 2 | Probable |
| Daily Operations | Employment and transfer of skills | 2 | 1 | 2 | 3 | 2 | 14 | 2 | Definite |
| Indirect Impacts | Employment and transfer of skills in Namibia's agricultural sector | 2 | 1 | 2 | 3 | 3 | 16 | 2 | Definite |

Desired Outcome: To see an increase in skills of local Namibians, as well as development and technological advancements in the agricultural industry.

Actions

Enhancement:

- ◆ Sourcing of employees and contractors must first be at local level and if not locally available, regional or national options should be considered. Deviations from this practice must be justified.
- ◆ Skills development and improvement programs must be made available as identified during performance assessments of employees.
- ◆ Inform employees about parameters and requirements for references upon employment.
- ◆ Provide managerial references for unofficial training or skills transfer.

Responsible Body:

- ◆ Proponent
- ◆ Contractors

Data Sources and Monitoring:

- ◆ Keep records of all training provided to employees.
- ◆ Ensure that all training is certified or managerial references provided (proof provided to the employees) inclusive of training attendance, completion and implementation.
- ◆ Include all information in a bi-annual report.

10.1.5 Revenue Generation and Employment

Skilled and unskilled labour are required for the operations and maintenance/construction activities associated with the farm. Livelihoods are thus sustained and the spending power of the local community increased. Revenue is generated through the sale of products (such as maize and vegetables) on national and international markets. Planting of certain GM crops, such as Round-Up Ready maize, can lead to reduced labour requirements to perform certain tasks (e.g. manual hoeing of weeds). This may offset possible job losses resulting from the planting of GM crops. Planting of the GMO crops will require less input in terms of pesticide application (including fuel and water) and labour, depending on the GM events planted. Therefore, producers will make time available for additional revenue generating activities to be considered. Many of the farming units in Namibia have diverse agricultural production units, which include agronomy, livestock farming, charcoal production and tourism.

| Project Activity/Resource | Nature (Status) | (A1) Importance | (A2) Magnitude | (B1) Permanence | (B2) Reversibility | (B3) Cumulative | Environmental Classification | Class Value | Probability |
|---------------------------|---|-----------------|----------------|-----------------|--------------------|-----------------|------------------------------|-------------|-------------|
| Construction | Employment and contribution to local and national economy | 2 | 1 | 2 | 2 | 2 | 12 | 2 | Definite |
| Daily Operations | Employment contribution to local and national economy | 2 | 1 | 3 | 3 | 1 | 14 | 2 | Definite |
| Indirect Impacts | Decrease in unemployment, contribution to local economy | 3 | 1 | 3 | 3 | 3 | 27 | 3 | Definite |

Desired Outcome: Contribution to national treasury and provision of employment to local Namibians.

Actions

Enhancement:

- ◆ The proponent must employ local Namibians where possible.
- ◆ If the skills exist locally, employees must first be sourced from the area, then the region and then nationally.
- ◆ Deviations from this practice must be justified.
- ◆ Opportunities for additional income generating activities to be investigated in order to sustain employment.
- ◆ Where feasible, employment of the same seasonal and/or temporary workforce year on year.

Responsible Body:

- ◆ Proponent

Data Sources and Monitoring:

- ◆ Bi-annual summary report based on employee records.

10.1.6 Demographic Profile and Community Health

Farming activities rely on labour. Jobseekers migrating to the Tsumeb area may lead to increased unemployment and expansion of informal settlements. Here, factors such as communicable disease like HIV/AIDS as well as alcoholism and drug abuse may thrive. These are typically aggravated when an influx of seasonal workers, and possible foreign construction teams and contractors, occur. An increase in foreign people in the area, linked to unemployment, may potentially increase the risk of criminal and socially/culturally deviant behaviour. However, since the farming unit is well established with an existing employee base, it is not foreseen that the project will result in significant migration to the Tsumeb area.

| Project Activity/Resource | Nature (Status) | (A1) Importance | (A2) Magnitude | (B1) Permanence | (B2) Reversibility | (B3) Cumulative | Environmental Classification | Class Value | Probability |
|---------------------------|---|-----------------|----------------|-----------------|--------------------|-----------------|------------------------------|-------------|-------------|
| Construction | In-migration and social ills related to foreign contractors temporarily on site | 2 | -1 | 1 | 1 | 2 | -8 | -1 | Probable |
| Daily Operations | Social ills possibly associated with staff and jobseekers | 2 | -1 | 1 | 2 | 2 | -10 | -2 | Probable |
| Indirect Impacts | The spread of disease | 2 | -1 | 2 | 2 | 2 | -12 | -2 | Improbable |

Desired Outcome: To prevent the occurrence of social ills and prevent the spread of diseases such as HIV/AIDS.

Actions:

Prevention:

- Appointment of reputable contractors where applicable.
- Adhere to all local authority by-laws relating to environmental health, which includes, but is not limited to, sanitation requirements for employees.
- Provide educational, awareness information for employees on various topics of social behaviour and HIV/AIDs.
- Disciplinary steps, within the legal parameters of Namibia, to be taken for socially deviant behaviour at the employee-housing compound or during working hours should be clearly stipulated in employment contracts.

Mitigation:

- Take disciplinary action against employees not adhering to contractual agreements with regard to socially deviant behaviour (e.g. alcohol or drug abuse during working hours).

Responsible Body:

- Proponent

Data Sources and Monitoring:

- Summary report based on educational programmes and training conducted.
- Employee contracts on file.
- Bi-annual report and review of employee demographics.

10.1.7 Agricultural Produce

The project is in line with the objectives of Namibia's NDPs and contributes to the economy of, and food security in, Namibia. Locally produced crops decrease the amount of crops that needs importing. GMO cultivation has the potential to safeguard crops against pests, thereby increasing the overall yield. This could (considering GMO maize), increase the amount of food available locally, if and when such crops are plagued by pests. The overall gain would be an increase in food security.

| Project Activity/Resource | Nature (Status) | (A1) Importance | (A2) Magnitude | (B1) Permanence | (B2) Reversibility | (B3) Cumulative | Environmental Classification | Class Value | Probability |
|-----------------------------------|--|-----------------|----------------|-----------------|--------------------|-----------------|------------------------------|-------------|-------------|
| Construction and Daily Operations | Contribution to economy, contribution to food security in Namibia | 1 | 2 | 3 | 3 | 2 | 16 | 2 | Definite |
| Indirect Impacts | Reduced import needs, increase in trade balance, spread of knowledge and skills, increased crop productivity | 1 | 2 | 3 | 3 | 3 | 18 | 2 | Definite |

Desired Outcome: Maximum contribution to the food security and economy of Namibia. Provide a positive contribution to the trade balance of Namibia by reducing the amount of imported produce and exporting higher value products.

Actions:

Enhancement:

- ◆ Teach employees on sustainable farming practices to enable the spread of knowledge and skills and thereby increase the productivity of small-scale farming as well.
- ◆ Diversification and continuous improvement to maximise sustainability of the farm.

Responsible Body:

- ◆ Proponent

Data Sources and Monitoring:

- ◆ Bi-annual reporting on educational programmes and training conducted.

10.1.8 Traffic

Potential traffic impacts will mostly be limited to the turnoff from the main road to the farm. Traffic is mostly related to the transport of staff, the delivery of fertilizers and seed, as well as the transport of crops to markets. As this is an existing operation, an increase in traffic impacts is expected to be unlikely. The turnoff from the C42 main road to the farming unit is a key section of concern. As this is an existing operation, an increase in traffic impacts is expected to be unlikely in the near future. The farming unit accommodates two NamWater boreholes, therefore access to the farm is also open to government officials. Additional and uncontrolled access to the farming unit will result in enhancing the potential security risk related to poaching and farm attacks.

| Project Activity / Resource | Nature (Status) | (A1) Importance | (A2) Magnitude | (B1) Permanence | (B2) Reversibility | (B3) Cumulative | Environmental Classification | Class Value | Probability |
|-----------------------------|---|-----------------|----------------|-----------------|--------------------|-----------------|------------------------------|-------------|-------------|
| Construction | Delivery of equipment and building supplies | 2 | -1 | 2 | 2 | 1 | -10 | -2 | Improbable |
| Daily Operations | Increased traffic, road wear and tear and accidents | 2 | -1 | 3 | 2 | 1 | -12 | -2 | Improbable |

Desired Outcome: Minimum impact on traffic and no transport or traffic related incidents.

Actions

Prevention:

- ◆ Erect clear signage regarding access and exit points at the farm as well as speed limits on the gravel roads within the farm where required.
- ◆ Only licenced drivers who are well trained to be allowed on the national roads.

Mitigation:

- ◆ If any traffic impacts are expected, possibly as a result of delivery of equipment or construction material, traffic management should be performed.

Responsible Body:

- ◆ Proponent

Data Sources and Monitoring:

- ◆ Record all traffic related complaints and the actions taken to prevent impacts from repeating itself.
- ◆ Compile a bi-annual report of all incidents reported, complaints received, and actions taken.

10.1.9 Health, Safety and Security

Daily operational and intermittent maintenance and construction activities on the farm are reliant on human labour. Such activities have varying degrees of health and safety risks. Examples include the operation of vehicles and machinery with moving parts, such as harvesters, and the handling of hazardous chemicals with inherent health hazards, such as pesticides and fuel, when ingested, inhaled or physical contact occur. Encounters with wild animals, and especially venomous species like snakes, may pose risks to employees. Security risks relates to unauthorized entry on the farming unit, theft and sabotage.

| Project Activity/Resource | Nature (Status) | (A1) Importance | (A2) Magnitude | (B1) Permanence | (B2) Reversibility | (B3) Cumulative | Environmental Classification | Class Value | Probability |
|---------------------------|--|-----------------|----------------|-----------------|--------------------|-----------------|------------------------------|-------------|-------------|
| Construction | Physical injuries, exposure to chemicals and criminal activities | 1 | -2 | 3 | 3 | 1 | -14 | -2 | Probable |
| Daily Operations | Physical injuries, exposure to chemicals and criminal activities | 1 | -2 | 3 | 3 | 2 | -16 | -2 | Probable |

Desired Outcome: To prevent injury, health impacts and theft.

Actions

Prevention:

- Implement and maintain an integrated health and safety management system, to act as a monitoring and mitigating tool.
- Comply with all health and safety standards as specified in the Labour Act and related legislation.
- Clearly label dangerous and restricted areas as well as dangerous equipment and products.
- Lock away or store all equipment and goods on site in a manner suitable to discourage criminal activities (e.g. theft).
- Provide all employees with required and adequate personal protective equipment (PPE) where required.
- Ensure that all personnel receive adequate training on the operational procedures of equipment and machinery and the handling of hazardous substances.
- Train selected personnel in first aid and ensure first aid kits are available on site.
- The contact details of all emergency services must be readily available.
- Implement a maintenance register for all equipment whose malfunction can lead to injury or exposure to hazardous substances.
- Apply and adhere to all industry specific health and safety procedures and regulations applicable to the handling of food produce for markets.

Mitigation:

- Treat all minor work related injuries immediately and obtain professional medical treatment if required.
- Assess any safety problems and implement corrective action to prevent future occurrences.

Responsible Body:

- Proponent
- Contractors

Data Sources and Monitoring:

- Record any incidents with the actions taken to prevent future occurrences.
- Compile a bi-annual report of all incidents reported. The report should contain dates when training was conducted and when safety equipment and structures were inspected and maintained.

10.1.10 Fire

Construction activities, failing electrical infrastructure, lightning and fires outside of designated areas may increase the risk of the occurrence of uncontrolled fires which may spread into the nearby fields and surrounding farms. Lightning may cause natural fires during the dry season. Farming operations do not present the same fire risk as operations which include charcoal production in the greater Abenab area. Uncontrolled fires which have generated in other areas will present a risk to existing and proposed operations.

| Project Activity/Resource | Nature (Status) | (A1) Importance | (A2) Magnitude | (B1) Permanence | (B2) Reversibility | (B3) Cumulative | Environmental Classification | Class Value | Probability |
|---------------------------|-----------------|-----------------|----------------|-----------------|--------------------|-----------------|------------------------------|-------------|-------------|
| Construction | Fire risk | 1 | -2 | 2 | 2 | 1 | -10 | -2 | Probable |
| Daily Operations | Fire risk | 1 | -2 | 2 | 2 | 1 | -10 | -2 | Probable |

Desired Outcome: To prevent property damage, veld fires, possible injury and impacts caused by uncontrolled fires.

Actions:

Prevention:

- ◆ Prepare a holistic fire protection and prevention plan. This plan must include evacuation plans and signage, an emergency response plan and a firefighting plan.
- ◆ Personnel training (safe operational procedures, firefighting, fire prevention and responsible housekeeping practices).
- ◆ Ensure all flammable chemicals are stored according to material safety data sheet (MSDS) and SANS instructions and all spills or leaks are cleaned immediately.
- ◆ Maintain regular site, mechanical and electrical inspections and maintenance.
- ◆ Maintain firefighting equipment and promote good housekeeping.
- ◆ Clean and maintain firebreaks at strategic locations on the properties, especially where vulnerable to external fire.
- ◆ Notify the farmers' association as well as all surrounding farmers if planned burns (e.g. to create firebreaks) are planned.
- ◆ Allow fires used for purposes such as cooking (by staff) in designated areas only.

Mitigation:

- ◆ Implement the fire protection and firefighting plan in the event of a fire.
- ◆ Quick response time by trained staff will limit the spread and impact of fire.

Responsible Body:

- ◆ Proponent
- ◆ Contractors

Data Sources and Monitoring:

- ◆ Maintain a register of all incidents on a daily basis. Include measures taken to ensure that such incidents do not repeat themselves.
- ◆ Compile a bi-annual incidents report. The report should also contain dates when fire drills were conducted and when firefighting equipment were tested and training given.

10.1.11 Noise

Noise is generated by various operational and possible construction activities. Machinery like generators, machinery, vehicles and harvesters cause elevated noise levels that may result in hearing impairment after long term exposure. Activities are generally remote from receptors other than the Proponent, his employees and their families residing on the farming unit. The nature of the noise is related mainly to the ongoing operation (for maintenance records) and mechanical maintenance typically on a farm.

| Project Activity/Resource | Nature (Status) | (A1) Importance | (A2) Magnitude | (B1) Permanence | (B2) Reversibility | (B3) Cumulative | Environmental Classification | Class Value | Probability |
|---------------------------|--|-----------------|----------------|-----------------|--------------------|-----------------|------------------------------|-------------|-------------|
| Construction | Excessive noise generated from construction activities – nuisance and hearing loss | 1 | -1 | 2 | 2 | 1 | -10 | -1 | Probable |
| Daily Operations | Noise generated from the operational activities – nuisance and hearing loss | 1 | -1 | 2 | 2 | 1 | -10 | -1 | Definite |

Desired Outcome: To prevent any nuisance and hearing loss due to noise generated.

Actions

Prevention:

- ◆ Follow Health and Safety Regulations of the Labour Act and/or World Health Organization (WHO) guidelines on maximum noise levels (Guidelines for Community Noise, 1999) to prevent hearing impairment.
- ◆ Regularly service all machinery to ensure minimal noise production.

Mitigation:

- ◆ Hearing protectors as standard PPE for workers in situations with elevated noise levels.

Responsible Body:

- ◆ Proponent
- ◆ Contractors

Data Sources and Monitoring:

- ◆ Health and Safety Regulations of the Labour Act and WHO Guidelines.
- ◆ Maintain a complaints register.
- ◆ Bi-annual report on complaints and actions taken to address complaints and prevent future occurrences.

10.1.12 Waste Production

Various waste streams result from the operational and possible construction and maintenance activities. Waste may include hazardous waste associated with hydrocarbon products and chemicals, as well as soil and water contaminated with such products. Construction waste may include building rubble and discarded equipment. Domestic waste will be generated by the residents and employees on the farm. Waste presents a contamination risk and when not removed regularly may become a health and/or fire hazard and attract wild animals and scavengers. Sewage is a form of liquid biological waste that needs disposal. Since no official waste disposal facilities, especially for hazardous waste, are available, all waste that cannot be re-used are burned at dedicated waste sites.

| Project Activity/Resource | Nature (Status) | (A1) Importance | (A2) Magnitude | (B1) Permanence | (B2) Reversibility | (B3) Cumulative | Environmental Classification | Class Value | Probability |
|---------------------------|--|-----------------|----------------|-----------------|--------------------|-----------------|------------------------------|-------------|-------------|
| Construction | Excessive waste production, littering, illegal dumping, contaminated materials | 1 | -2 | 2 | 2 | 2 | -12 | -2 | Definite |
| Daily Operations | Excessive waste production, littering, contaminated materials | 1 | -2 | 2 | 2 | 2 | -12 | -2 | Definite |

Desired Outcome: To reduce the amount of waste produced and prevent pollution and littering.

Actions

Prevention:

- ◆ Implement waste reduction measures. All waste that can be re-used/recycled must be kept separate.
- ◆ Ensure adequate temporary storage facilities for disposed waste are available.
- ◆ Prevent windblown waste from entering the environment.
- ◆ Prevent scavenging (human and non-human) of waste at the storage facilities.
- ◆ Educate employees on the importance of proper waste handling and disposal.

Mitigation:

- ◆ Waste should be disposed of regularly and at appropriately classified disposal facilities, this includes hazardous material (empty chemical containers and contaminated materials, soil and water)
- ◆ Discarded waste should be disposed of and burned regularly at a dedicated site to reduce health and pollution risks.
- ◆ Empty chemical containers that may present a contamination/health risk must be treated as hazardous waste. Workers should not be allowed to collect such containers for purposes of storing water or food. This can be achieved by puncturing or crushing such containers prior to disposal.
- ◆ Liaise with the applicable authorities regarding waste and handling of hazardous waste.
- ◆ Ensure all ablution facilities are connected to properly constructed septic tank systems to prevent groundwater contamination.

Responsible Body:

- ◆ Proponent
- ◆ Contractors

Data Sources and Monitoring:

- ◆ Maintain a register of disposal of hazardous waste. This should include type of waste, volume as well as disposal method/facility.
- ◆ Record any complaints received regarding waste with notes on actions taken.
- ◆ All information to be included in a bi-annual report.

10.1.13 Ecosystem and Biodiversity Impact

Agriculture and related activities are ongoing at the farming unit. Possible expansion is planned on existing cleared areas and no further impacts on vegetation are thus expected from additional land clearing. Pollution of the environment may however impact on the ecosystem and biodiversity. Poaching and illegal collection of plant and animal materials may occur. Irresponsible pesticide use, for example as method of vermin control, may impact on scavengers such as vultures and in the long run on top predators through biomagnification in higher trophic levels. Pesticides by nature are harmful to the environment, planting Bt crops that targets specifically Lepidopterans, reduce the need for spraying insecticides. This result in an increased biodiversity as compared to fields treated with traditional insecticides. Over abstraction of groundwater may detrimentally affect endemic species linked to the groundwater and related caves.

| Project Activity/Resource | Nature (Status) | (A1) Importance | (A2) Magnitude | (B1) Permanence | (B2) Reversibility | (B3) Cumulative | Environmental Classification | Class Value | Probability |
|---------------------------|---|-----------------|----------------|-----------------|--------------------|-----------------|------------------------------|-------------|-------------|
| Construction | Impact on fauna and flora. Loss of biodiversity | 2 | -1 | 3 | 2 | 2 | -14 | -2 | Probable |
| Daily Operations | Impact on fauna and flora. Loss of biodiversity – poaching, poisoning, etc. | 2 | -1 | 2 | 2 | 2 | -12 | -2 | Probable |

Desired Outcome: To avoid pollution of, and impacts on, the ecological environment.

Actions.

Prevention:

- ◆ Strictly adhere to pesticide application instructions and use pesticides only for the purposes for which it is registered and marketed. Importantly, pesticides should not be used to kill vermin unless specifically registered for that purpose, and even then alternative, environmentally friendly methods should be investigated and used.
- ◆ Prevent pesticides from ending up in the hands of potential poachers.
- ◆ Educate all contracted and permanent employees on the value of biodiversity and strict conditions prohibiting harvesting and poaching of fauna and flora must be part of employment contracts. Include prohibitions or regulations on the collection of firewood.
- ◆ Regular inspection of fences, game footpaths and other sites for snares, traps or any other illegal activities.
- ◆ Prevent spray drift by applying pesticides during calm weather conditions.
- ◆ Proper training of operational personnel.
- ◆ Over-abstraction of groundwater may potentially have devastating effects on plant and animal populations reliant on it. It not only include the drying up of springs, dying of trees and migration or dying of animals but also the lowering of cave water levels.
- ◆ Ensure all waste oil handling is conducted on impermeable or bunded areas.

Mitigation:

- ◆ For construction activities, if any, contain construction material to a designated laydown area and prevent unnecessary movement out of areas earmarked for clearing and construction.
- ◆ Report any extraordinary animal sightings to the Ministry of Environment, Forestry and Tourism.
- ◆ Prevent scavenging of waste by fauna.
- ◆ Take disciplinary action against any employees failing to comply with contractual conditions related to poaching and the environment.

Responsible Body:

- ◆ Contractor
- ◆ Proponent

Data Sources and Monitoring:

- ◆ Report on all extraordinary animal or plant sightings or instances of poaching.
- ◆ Keep frequent records of borehole water levels and abstracted water volumes to identify any trends or consistent reduction in water levels.
- ◆ Compile a bi-annual report on all monitoring results.

10.1.14 Groundwater, Surface Water and Soil Contamination

Leakages and spillages of hazardous substances from vehicles, waste oil handling and accidental fuel, oil or hydraulic fluid spills during the operational phase may contaminate the environment. Increase of nutrient levels (from over application of fertilizers or pesticides) in the soil that can leach to the groundwater. Volumes of pesticides used are in fact lower for GM crops, especially for insecticides. Pollution due to sewerage system overflow or leakage may further put the groundwater at risk.

| Project Activity/Resource | Nature (Status) | (A1) Importance | (A2) Magnitude | (B1) Permanence | (B2) Reversibility | (B3) Cumulative | Environmental Classification | Class Value | Probability |
|---------------------------|---|-----------------|----------------|-----------------|--------------------|-----------------|------------------------------|-------------|-------------|
| Daily Operations | Hazardous material, spillages, hydrocarbon leakages from vehicles and machinery. | 2 | -1 | 2 | 2 | 1 | -10 | -2 | Improbable |
| Daily Operations | Over application of fertilizer, pesticides, herbicides, etc. Sewerage system malfunction. | 2 | -1 | 2 | 2 | 1 | -10 | -2 | Improbable |

Desired Outcome: To prevent the contamination of groundwater, surface water and soil.

Actions

Prevention:

- ◆ Appoint reputable contractors.
- ◆ Vehicles may only be serviced on a suitable spill control structure.
- ◆ Regular inspections and maintenance of all vehicles to ensure no leaks are present.
- ◆ All hazardous chemicals and fuel should be stored in a sufficiently bunded area, as per MSDS requirements.
- ◆ Ensure all waste oil handling is conducted on impermeable or bunded areas.
- ◆ Follow prescribed dosage of fertilizers and pesticides / herbicides and to avoid over application.
- ◆ Maintain sewerage systems and conduct regular monitoring.
- ◆ All hazardous waste must be removed from the site and disposed of timeously at a recognised hazardous waste disposal facility, including any polluted soil or water.

Mitigation:

- ◆ All spills must be cleaned up immediately.
- ◆ Consult relevant Material Safety Data Sheet (MSDS) information and a suitably qualified specialist where needed.

Responsible Body:

- ◆ Proponent
- ◆ Contractors

Data Sources and Monitoring:

- ◆ Maintain Material Safety Data Sheets for hazardous chemicals.
- ◆ Soil should be sampled and analysed annually to ensure the correct amounts of fertilizer is applied and soil and groundwater quality is maintained.
- ◆ Groundwater should be sampled and analysed to test for nitrate concentrations from the fertilizer and for traces of chemicals used in pesticides and herbicides.
- ◆ Registers be kept by the Proponent on the type, quantities and frequency of application of fertiliser, pesticides and any other chemicals utilised in crop production.
- ◆ A register of all incidents must be maintained on a daily basis. This should include measures taken to ensure that such incidents do not repeat themselves.
- ◆ All spills or leaks must be reported on and cleaned up immediately.

10.1.15 Groundwater Abstraction

Groundwater abstraction is a very sensitive topic in a dry country where the value of land is drastically reduced if no or unusable groundwater is present on the land. Abstraction of groundwater must be done in a sensible way not to impact on other groundwater users that depend on such groundwater. This includes water abstracted for human and animal use, irrigation, and also ecosystems that depend on groundwater. A typical groundwater balance was compiled to illustrate the potential consequences of over abstraction of groundwater, see Figure 9-1. Recharge to the area is considered to be comparatively high.

In a typical groundwater environment, a water balance would consist of inflow and outflow of the groundwater system. Over time an equilibrium (or steady state) is normally reached with rising water tables following good recharge events and declining water tables when recharge is below average. Inflow into the system would typically be from infiltration following rainfall in the area and in upstream areas. Outflow would be comprised of water leaving the system through springs and as outflow over the lower boundary of the groundwater system as well as evapotranspiration losses. Groundwater abstraction through boreholes is important as this is normally necessary to sustain human and animal demands where such users became essentially dependant on the abstracted groundwater as a reliable and sustainable source.

Typical consequences of over abstraction will include a lowering in the water table. This may further lead to the drying up of boreholes, springs, and shallow wells. Vegetation will also be impacted where such vegetation has access to groundwater.

| Project Activity/Resource | Nature (Status) | (A1) Importance | (A2) Magnitude | (B1) Permanence | (B2) Reversibility | (B3) Cumulative | Environmental Classification | Class Value | Probability |
|---------------------------|--|-----------------|----------------|-----------------|--------------------|-----------------|------------------------------|-------------|-------------|
| Daily Operations | Over-abstraction of the local aquifer, decrease in the local hydraulic head. | 2 | -2 | 2 | 2 | 2 | -24 | -3 | Probable |

Desired Outcome: To utilise the groundwater sustainably.

Actions

Prevention:

- ◆ Spread the water abstraction points over a larger area to diffuse the impact.
- ◆ Monthly water level monitoring.
- ◆ Maintain safe abstraction rates prescribed by test pump evaluations (an abstraction permit with prescribed rates from the MAWLR is a requirement for this project).

Mitigation:

- ◆ Reduce abstraction when the water levels nears 5 m below the average rest water level of each borehole.

Responsible Body:

- ◆ Proponent

Data Sources and Monitoring:

- ◆ Monthly boreholes rest water level monitoring.
- ◆ Baseline values should be reviewed every three years based on all historic water level data.
- ◆ A summary report on all monitoring results must be prepared.
- ◆ The Proponent supply monitoring returns to the MAWLR, as required by the permit.

10.1.16 Visual Impact

This impact relates to the aesthetic appearance of the site during operations. This impact will be minimal due to the area already being disturbed and widely utilised for agricultural activities. The impact will therefore mostly relate to poor housekeeping and waste not disposed of timeously. Operations at the farm are well kept with the highest standard of neatness and cleanliness exhibited throughout all components of the operations, inclusive of employee housing.

| Project Activity/Resource | Nature (Status) | (A1) Importance | (A2) Magnitude | (B1) Permanence | (B2) Reversibility | (B3) Cumulative | Environmental Classification | Class Value | Probability |
|---------------------------|--|-----------------|----------------|-----------------|--------------------|-----------------|------------------------------|-------------|-------------|
| Construction | Aesthetic appearance and integrity of the site | 1 | 1 | 2 | 2 | 2 | 6 | 1 | Probable |
| Daily Operations | Aesthetic appearance and integrity of the site | 1 | 1 | 2 | 2 | 2 | 6 | 1 | Probable |

Desired Outcome: To minimise aesthetic impacts associated with the farm.

Actions

Mitigation:

- ◆ Regular waste disposal, good housekeeping and routine maintenance on infrastructure will ensure that the longevity of structures are maximised and maintain a low visual impact.

Responsible Body:

- ◆ Proponent
- ◆ Contractors

Data Sources and Monitoring:

- ◆ Compile a bi-annual report of all complaints received and actions taken.

10.1.17 Cumulative Impact

Possible negative cumulative impacts (i.e. the build-up of minor impacts to become more significant) associated with the operational phase and any maintenance/construction activities are mainly linked to traffic, reduction in soil and groundwater quality and groundwater availability. The cumulative increase in employees in the area may put more pressure on biodiversity as a result of poaching or harvesting of plant and animal products. The cumulative positive impacts from farming in the Oshikoto Region relates to increased and sustained employment, revenue generation and overall improved living conditions and livelihoods as a result of increased spending power.

| Project Activity/Resource | Nature (Status) | (A1) Importance | (A2) Magnitude | (B1) Permanence | (B2) Reversibility | (B3) Cumulative | Environmental Classification | Class Value | Probability |
|--|---|-----------------|----------------|-----------------|--------------------|-----------------|------------------------------|-------------|-------------|
| Daily Construction and Operations (Negative) | Waste production, pollution, etc. The build-up of minor impacts to become more significant | 2 | -1 | 2 | 2 | 1 | -10 | -2 | Probable |
| Daily Construction and Operations (Positive) | Employment, skills development, revenue generation | 2 | 1 | 2 | 2 | 1 | 10 | 2 | Definite |

Desired Outcome: To minimise cumulative all impacts associated with the farm.

Actions

Mitigation:

- ◆ Addressing each of the individual impacts as discussed and recommended in the EMP would reduce the cumulative impact.
- ◆ Reviewing biannual reports for any new or re-occurring impacts or problems would aid in identifying cumulative impacts. Planning and improvement of the existing mitigation measures can then be implemented.

Responsible Body:

- ◆ Proponent

Data Sources and Monitoring:

- ◆ Reviewing monitoring results based on all other impacts will give an overall assessment of the impacts of the operational phase.

10.2 DECOMMISSIONING AND REHABILITATION

Closure and decommissioning of agricultural and related activities on the farm as a whole is not foreseen during the validity of the environmental clearance certificate or in the near future. However, it is more likely that certain components may be decommissioned. Decommissioning is therefore included for this purpose as well as the fact that construction activities may also include modification and decommissioning of infrastructure. Future land use after decommissioning should be assessed prior to decommissioning and rehabilitation initiated if the land would not be used for future purposes. Should decommissioning occur at any stage, rehabilitation of the area may be required. Decommissioning will entail the complete removal of all infrastructure including buildings and irrigation infrastructure. Any pollution present on the site must be remediated. The impacts associated with this phase include noise and waste production as structures are dismantled. Noise must be kept within WHO standards. Waste should be contained and disposed of at a dedicated waste disposal site and not dumped in the surrounding areas. The EMP for the farm will have to be reviewed at the time of full decommissioning to cater for changes made to the site and to implement guidelines and mitigation measures.

10.3 ENVIRONMENTAL MANAGEMENT SYSTEM

The Proponent could implement an environmental management system (EMS) for their operations. An EMS is an internationally recognized and certified management system that will ensure ongoing incorporation of environmental constraints. At the heart of an EMS is the concept of continual improvement of environmental performance with resulting increases in operational efficiency, financial savings and reduction in environmental, health and safety risks. An effective EMS would need to include the following elements:

- ◆ A stated environmental policy which sets the desired level of environmental performance;
- ◆ An environmental legal register;
- ◆ An institutional structure which sets out the responsibility, authority, lines of communication and resources needed to implement the EMS;
- ◆ Identification of environmental, safety and health training needs;
- ◆ An environmental program(s) stipulating environmental objectives and targets to be met, and work instructions and controls to be applied in order to achieve compliance with the environmental policy; and
- ◆ Periodic (internal and external) audits and reviews of environmental performance and the effectiveness of the EMS.
- ◆ The EMP.

11 CONCLUSION

Agricultural and related activities as performed on the farm Cuxhaven No. 1278 in the Oshikoto Region, contributes positively to the economy of Namibia. Food is produced for national markets and the sale of livestock for meat production to both local and international markets. A number of employment opportunities are sustained and skills development within the local workforce occur. Revenue is generated that contributes to the Namibian economy.

Negative impacts associated with operational and intermittent maintenance and construction activities on the farming unit, as summarised in, section 10 can successfully be mitigated. Implementing a HSE policy will contribute to effective management procedures to prevent and mitigate impacts. All regulations relating to the agricultural and related activities of the Proponent, including health and safety legislation, should be adhered to and implemented where applicable. Groundwater and soil pollution must be prevented at all times and over abstraction of groundwater prevented. Fire prevention should be key, fire response plans in place, and regular firefighting training provided to key employees. The GMO management plan as present in Appendix C must be implemented and strictly adhered to. All staff must be made aware of the importance of biodiversity and the poaching or illegal harvesting of animal and plant products prohibited. This includes the proper handling and correct application of pesticides. Any waste produced must be properly disposed, re-used, or recycled where possible.

The EMP (Section 10) should be used as an on-site reference document for the operations of the farm. Parties responsible for transgression of the EMP should be held responsible for any rehabilitation that may need to be undertaken. The Proponent could use an in-house Health, Safety, Security and Environmental Management System in conjunction with the EMP. All operational personnel must be taught the contents of these documents.

Should the Directorate of Environmental Affairs (DEA) agree with the impacts and related mitigation measures, they may issue an environmental clearance certificate to the Proponent. The environmental clearance certificate will render this document legally binding on the Proponent. The assessment process's aim is not to stop the farming activities, or any of its components, but to rather determine its impact and guide sustainable and responsible development as per the spirit of the EMA.

12 BIBLIORGAPHY

- Atlas of Namibia Project. 2002. Directorate of Environmental Affairs, Ministry of Environment and Tourism (www.met.gov.na) [Accessed from http://www.uni-koeln.de/sfb389/e/e1/download/atlas_namibia/index_e.htm]
- BirdLife International. 2021. *Aquila nipalensis*. The IUCN Red List of Threatened Species 2021: e.T22696038A205452572. <https://dx.doi.org/10.2305/IUCN.UK.2021-3.RLTS.T22696038A205452572.en>. Accessed on 20 June 2023.
- BirdLife International. 2021. *Falco vespertinus*. The IUCN Red List of Threatened Species 2021: e.T22696432A200136196. <https://dx.doi.org/10.2305/IUCN.UK.2021-3.RLTS.T22696432A200136196.en>. Accessed on 20 June 2023.
- BirdLife International. 2021. *Gyps africanus*. The IUCN Red List of Threatened Species 2021: e.T22695189A204461164. <https://dx.doi.org/10.2305/IUCN.UK.2021-3.RLTS.T22695189A204461164.en>. Accessed on 20 June 2023.
- BirdLife International. 2021. *Neophron percnopterus*. The IUCN Red List of Threatened Species 2021: e.T22695180A205187871. <https://dx.doi.org/10.2305/IUCN.UK.2021-3.RLTS.T22695180A205187871.en>. Accessed on 20 June 2023.
- BirdLife International. 2020. *Polemaetus bellicosus*. The IUCN Red List of Threatened Species 2020: e.T22696116A172287822. <https://dx.doi.org/10.2305/IUCN.UK.2020-3.RLTS.T22696116A172287822.en>. Accessed on 20 June 2023.
- BirdLife International. 2020. *Sagittarius serpentarius*. The IUCN Red List of Threatened Species 2020: e.T22696221A173647556. <https://dx.doi.org/10.2305/IUCN.UK.2020-3.RLTS.T22696221A173647556.en>. Accessed on 20 June 2023.
- BirdLife International. 2020. *Terathopius ecaudatus*. The IUCN Red List of Threatened Species 2020: e.T22695289A174413323. <https://dx.doi.org/10.2305/IUCN.UK.2020-3.RLTS.T22695289A174413323.en>. Accessed on 20 June 2023.
- BirdLife International. 2021. *Trigonoceps occipitalis*. The IUCN Red List of Threatened Species 2021: e.T22695250A205380033. <https://dx.doi.org/10.2305/IUCN.UK.2021-3.RLTS.T22695250A205380033.en>. Accessed on 20 June 2023.
- Bosman Q; Botha P. 2016 August; Silverlands Vineyards Agricultural Project Aussenkehr, Namibia Social Impact Assessment Namibia
- Clarias cavernicola* has most recently been assessed for The IUCN Red List of Threatened Species in 2007. *Clarias cavernicola* is listed as Critically Endangered under criteria B1ab(iii)+2ab(iii).
- Climate Engine. (2020). Desert Research Institute and University of Idaho. Accessed on (15/01/2021). <http://climateengine.org>.
- Climate Risk Profile: Namibia (2021): The World Bank Group
- Curtis B. & Mannheimer C. 2005. Tree Atlas of Namibia. National Botanical Research Institute, Windhoek. 674 pages.
- Directorate of Environmental Affairs, 2008. Procedures and Guidelines for Environmental Impact Assessment (EIA) and Environmental Management Plans (EMP), Directorate of Environmental Affairs, Ministry of Environment and Tourism, Windhoek.
- Durant, S.M., Groom, R., Ipavec, A., Mitchell, N. & Khalatbari, L. 2022. *Acinonyx jubatus*. The IUCN Red List of Threatened Species 2022: e.T219A124366642. <https://dx.doi.org/10.2305/IUCN.UK.2022-1.RLTS.T219A124366642.en>. Accessed on 20 June 2023.

- Emslie, R. 2020. *Diceros bicornis*. The IUCN Red List of Threatened Species 2020: e.T6557A152728945. <https://dx.doi.org/10.2305/IUCN.UK.2020-1.RLTS.T6557A152728945.en>. Accessed on 20 June 2023.
- Funk, C., Peterson, P., Landsfeld, M., Pedreros, D., Verdin, J., Shukla, S., Husak, G., Rowland, J., Harrison, L., Hoell, A. and Michaelsen, J., (2015) The climate hazards group infrared precipitation with stations - A new environmental record for monitoring extremes. *Scientific Data*, 2, 150066. Accessed on (15/01/2021). <https://doi.org/10.1038/sdata.2015.66>.
- Geological Survey of Namibia; 1981. Geological Map 1:250,000. Geological Series 2218 Gobabis (S.W.A/Namibia).
- Geological Survey of Namibia; Geological Map 1:1,000,000.
- https://www.meteoblue.com/en/weather/historyclimate/climatemodelled/abenab_namibia_3358822
Accessed May 2023
- IWRM Plan Joint Venture Namibia. (2010). Integrated Water Resources Management Plan. Windhoek: Ministry of Agriculture, Water and Forestry.
- Miller, R.McG. (2008). *The Geology of Namibia: Volume 2. Neoproterozoic to Lower Paleozoic*. Geological Survey of Namibia.
- Namibia Statistics Agency. Namibia 2011 Population and Housing Census Main Report.
- Namibia Statistics Agency. Namibia household Income and Expenditure Survey 2009/2010.
- Pastakia, C.M.R.; 1998; *The Rapid Impact Assessment Matrix (RIAM) – A new tool for Environmental Impact Assessment*.
- Phocaides, A.; 2007; *Handbook on pressurized Irrigation Techniques*; Food And Agriculture Organization of The United Nations.
- Tweddle, D. & Marshall, B.E. 2007. *Oreochromis andersonii*. The IUCN Red List of Threatened Species 2007: e.T60623A12385801. <https://dx.doi.org/10.2305/IUCN.UK.2007.RLTS.T60623A12385801.en>. Accessed on 20 June 2023.

Appendix A: Certificates



MINISTRY OF MINES AND ENERGY

**PETROLEUM PRODUCTS AND ENERGY ACT, 1990
PETROLEUM PRODUCTS REGULATIONS (2000)**

CONSUMER INSTALLATION CERTIFICATE

[Regulation 18 (5)]

| | | | | | | |
|---|--|---|---|------------------------|---|--|
| CONSUMER INSTALLATION CERTIFICATE | | PERMANENT* <input checked="" type="checkbox"/> | PETROL* | Certificate No. | | |
| | | TEMPORARY* | DIESEL* <input checked="" type="checkbox"/> | CI/1770/2002 | | |
| Name of certificate-holder | | Paul S Potgieter | | | | |
| Address of certificate-holder | | Physical address | Postal address | | | |
| | | Farm Cuxhaven 1278 Tsumeb | Box 67 Tsumeb | | | |
| Nature of activity to which certificate relates* | Commercial/Industrial Undertaking | Farming Operation <input checked="" type="checkbox"/> | Mining Operation | | | |
| If storage tank is to be permanently installed, location of site | | N/A | | | | |
| Conditions applicable to Certificate <i>See next page for general and special conditions applicable to licence.</i> | | | | | | |
| Date of issue of certificate | | 11 February 2002 | | | | |
| In the case of a temporary licence, period of validity | | | | | | |
| Issued by the Minister of Mines and Energy in terms of regulation 18(5), on 11 February 2002 at Windhoek | | <table border="1"> <tr> <td>  Minister: Mines and Energy </td> <td> MINISTRY OF MINES AND ENERGY OFFICE OF THE MINISTER Official Stamp (for office use) 1 1 PRIVATE BAG 1297 WINDHOEK OFFICIAL </td> </tr> </table> | | |  Minister: Mines and Energy | MINISTRY OF MINES AND ENERGY OFFICE OF THE MINISTER Official Stamp (for office use) 1 1 PRIVATE BAG 1297 WINDHOEK OFFICIAL |
|  Minister: Mines and Energy | MINISTRY OF MINES AND ENERGY OFFICE OF THE MINISTER Official Stamp (for office use) 1 1 PRIVATE BAG 1297 WINDHOEK OFFICIAL | | | | | |

* Mark the appropriate item

Appendix B: Hydrogeological Specialist Study

**IRRIGATION BASED AGRICULTURAL ACTIVITIES AND
CULTIVATION OF GENETICALLY MODIFIED MAIZE ON FARM
CUXHAVEN OOS - OSHIKOTO REGION
HYDROGEOLOGICAL SPECIALIST STUDY**



Assessed by:



Assessed for:

L M Potgieter

February 2024

| | | |
|-------------------------------|---|---|
| Project: | IRRIGATION BASED AGRICULTURAL ACTIVITIES AND CULTIVATION OF GENETICALLY MODIFIED MAIZE ON FARM CUXHAVEN OOS - OSHIKOTO REGION: HYDROGEOLOGICAL SPECIALIST STUDY | |
| Report Version/Date | V1 February 2024 | |
| Prepared for | L M Potgieter P O Box 761 Tsumeb | |
| Lead Consultant | Geo Pollution Technologies (Pty) Ltd PO Box 11073 Windhoek Namibia | TEL.: (+264-61) 257411 FAX.: (+264) 88626368 |
| Main Project Team | Pierre Botha (Leader) (B.Sc. Geology/Geography); (B.Sc. (Hons) Hydrology/Hydrogeology) Surene Botha (B.Sc. Geology/Geochemistry); (B.Sc. (Hons.) Geology); (M.Sc. Hydrology/Hydrogeology) | |
| Cite this document as: | Botha P & Botha S; February 2024; Irrigation Based Agricultural and Cultivation of Genetically Modified Maize on Farm Cuxhaven Oos, Oshikoto Region: Hydrogeological Specialist Study | |
| Copyright | Copyright on this document is reserved. No part of this document may be utilised without the written permission of Geo Pollution Technologies (Pty) Ltd. | |
| Report Approval | Pierre Botha Managing Director | |

TABLE OF CONTENTS

| | | |
|-----------|---|-----------|
| 1 | INTRODUCTION..... | 1 |
| 2 | SCOPE OF WORK..... | 1 |
| 3 | METHODOLOGY | 1 |
| 4 | ADMINISTRATIVE, LEGAL AND POLICY REQUIREMENTS..... | 1 |
| 5 | DESCRIPTION OF NATURAL ENVIRONMENT | 2 |
| | 5.1 HYDROGEOLOGICAL LOCATION | 2 |
| | 5.2 CLIMATE..... | 3 |
| | 5.3 TOPOGRAPHY & DRAINAGE | 4 |
| | 5.4 SOIL..... | 5 |
| | 5.5 GEOLOGY AND HYDROGEOLOGY | 6 |
| 6 | ASSESSMENT OF WATER LEVEL MONITORING DATA..... | 11 |
| 7 | WATER SUPPLY..... | 15 |
| | 7.1 GROUNDWATER USAGE..... | 15 |
| 8 | ASSESSMENT OF IMPACTS | 17 |
| | 8.1 GROUNDWATER ABSTRACTION | 19 |
| | 8.2 GROUNDWATER, SURFACE WATER AND SOIL CONTAMINATION | 21 |
| 9 | CONCLUSION | 22 |
| 10 | REFERENCES | 22 |

LIST OF FIGURES

| | | |
|------------|--|----|
| FIGURE 1-1 | PROJECT LOCATION..... | 1 |
| FIGURE 5-1 | DAILY AND SEASONAL RAINFALL FROM CHIRPS-2 DATA (FUNK ET AL., 2015) | 4 |
| FIGURE 5-2 | ASPECT SLOPE AND SURFACE DRAINAGE..... | 5 |
| FIGURE 5-3 | DOMINANT SOIL TYPE AND AGRO ECOLOGICAL ZONE | 6 |
| FIGURE 5-4 | HYDROGEOLOGICAL MAP..... | 9 |
| FIGURE 5-5 | GROUNDWATER BASIN WITH RAINFALL AND INFERRED GROUNDWATER FLOW | 10 |
| FIGURE 5-6 | GROUNDWATER QUALITY | 11 |
| FIGURE 6-1 | MONITOR BOREHOLE LOCATIONS, CAVES, SINKHOLES AND SPRINGS | 12 |
| FIGURE 6-2 | REGIONAL WATER LEVEL PROFILE | 13 |
| FIGURE 6-3 | REGIONAL WATER LEVEL CHANGES AND MONTHLY RAINFALL | 14 |
| FIGURE 7-1 | LOCATIONS OF BOREHOLES..... | 16 |
| FIGURE 8-1 | CONCEPTUAL GROUNDWATER BALANCE WITH OVER ABSTRACTION SCENARIO..... | 19 |

LIST OF TABLES

| | | |
|------------|--|----|
| TABLE 4-1. | NAMIBIAN LAW APPLICABLE TO THE PROJECT..... | 2 |
| TABLE 5-1 | RAINFALL STATISTICS BASED ON CHIRPS-2 DATA (FUNK ET AL., 2015)..... | 4 |
| TABLE 5-2 | GROUNDWATER STATISTICS..... | 8 |
| TABLE 7-1 | SUMMARY OF BOREHOLE INFORMATION OBTAINED FROM THE PROPONENT..... | 15 |
| TABLE 8-1 | ASSESSMENT CRITERIA | 17 |
| TABLE 8-2 | ENVIRONMENTAL CLASSIFICATION OF IMPACTS (PASTAKIA 1998)..... | 18 |
| TABLE 8-3 | ASSESSMENT – GROUNDWATER ABSTRACTION | 20 |
| TABLE 8-4 | ASSESSMENT – GROUNDWATER, SURFACE WATER AND SOIL CONTAMINATION | 21 |

1 INTRODUCTION

Geo Pollution Technologies (Pty) Ltd was appointed by L M Potgieter (the Proponent) to undertake a hydrogeological specialist study for irrigation activities and cultivation of genetically modified maize on the farm Cuxhaven Oos (FMB/01278) in the Oshikoto Region. Existing activities on the farm are focussed on irrigated crop cultivation and livestock farming. The Proponent currently utilizes an area of approximately 260 ha for cultivation, of which 76 ha is irrigated by means of centre pivot systems utilising abstracted groundwater, and the remaining 184 ha is used for dry land crop production.




Figure 1-1 Project location

2 SCOPE OF WORK

The aims of the study were to:

1. Conduct a hydrogeological assessment based on data obtained from an in-field hydrocensus survey.
2. Gather historic information and compile a hydrogeological assessment based on the information.

3 METHODOLOGY

Obtain and review all available geological and hydrogeological information/reports for the investigation area. Review and delineation of hydrogeological catchment and sub-catchments within the investigation area. This will be based on historic groundwater level data contained in the Department of Water Affairs (DWA) database and from hydrocensus data gathered on behalf of the Proponent. Prepare a specialist report of the investigation.

4 ADMINISTRATIVE, LEGAL AND POLICY REQUIREMENTS

To protect the environment and achieve sustainable development, all projects, plans, programmes and policies deemed to have adverse impacts on the environment require an environmental impact

assessment (EIA), as per the Namibian legislation. The key legislation provided in Table 4-1 govern the environmental assessment process in Namibia and/or are relevant to the project.

Table 4-1. Namibian Law applicable to the project

| Law | Key Aspects |
|---|--|
| The Namibian Constitution | <ul style="list-style-type: none"> ◆ Incorporate a high level of environmental protection. ◆ Land, water and natural resources below and above the surface of the land and in the continental shelf and within the territorial waters and the exclusive economic zone of Namibia shall belong to the State if they are not otherwise lawfully owned. |
| Environmental Management Act Act No. 7 of 2007, Government Notice No. 232 of 2007 | <ul style="list-style-type: none"> ◆ Defines the environment. ◆ Promote sustainable management of the environment and the use of natural resources. |
| Water Resources Management Act Act No. 11 of 2013 | <ul style="list-style-type: none"> ◆ Provide for management, protection, development, use and conservation of water resources. ◆ Prevention of water pollution and assignment of liability. |
| Soil Conservation Act Act No. 76 of 1969 | <ul style="list-style-type: none"> ◆ Law relating to the combating and prevention of soil erosion, the conservation, improvement and manner of use of the soil and vegetation and the protection of the water sources Namibia. |

Relevant water resource development and related activities listed as activities requiring an environmental clearance certificate are (Government Notice No. 29 of 2012):

- 8.1 The abstraction of ground or surface water for industrial or commercial purposes.
- 8.2 The abstraction of groundwater at a volume exceeding the threshold authorised in terms of a law relating to water resources.
- 8.6 Construction of industrial and domestic wastewater treatment plants and related pipeline systems.
- 8.7 Irrigation schemes for agriculture excluding domestic irrigation.
- 8.8 Construction and other activities in water courses within flood lines.
- 8.9 Construction and other activities within a catchment area.

The relevance of 8.2 is not clear as to under which act such a threshold is defined, if any. The Water Resources Management Act (Act No. 11 of 2013) do not define such a threshold and existing water control areas in which abstraction permits would be required, was not repealed. The repealed Water Act (Act No. 54 of 1956) only requires abstraction permits within water control areas, see Figure 5-5. Abstraction permits are currently issued by the Ministry of Agriculture Water and Land Reform (MAWLR). The project falls inside a control area, thus an abstraction permit is a requirement.

Within the Water Resources Management Act (Act No. 11 of 2013) it is clearly stipulated that the purification and disposal of industrial water and effluents as well as the disposal of effluents by local authorities is subjected to the requirements of the Act. Agricultural activities is not subjected to the requirements of the Act, making the implementation of 8.6 questionable. The return period for flood lines is not provided for, nor a definition of flood lines to make 8.8 applicable. It is however in the Proponent's best interest to ensure that the project area is outside a flood risk area. All land in Namibia is in some form of catchment area, making the practical implementation of 8.9 questionable. It however remains important to consider all activities that would/may impact on the groundwater.

5 DESCRIPTION OF NATURAL ENVIRONMENT

5.1 HYDROGEOLOGICAL LOCATION

The Project is located within the Owambo Groundwater Basin (Figure 5-5). According to the Ministry of Agriculture, Water and Forestry (MAWF, 2006) the farm is located inside the

Tsumeb-Otavi-Grootfontein Subterranean Water Control Area, Government Notice 1969 of 13 November 1970 and Proclamation 278 of 31 December 1976 (Extension). The farm also fall under a sub-division of the water control area (Tsumeb - B2), known as the eastern half of the Tsumeb-Abenab Synclinorium sub-catchment (Bäumle, 2004). Government regulates groundwater usage in this area and all other groundwater related activities like drilling, cleaning or deepening of boreholes and rates of water abstraction. See Figure 5-5 for a map indicating the water control area, groundwater basin and inferred groundwater flow.

Implications and Impacts

Groundwater Basin committees will likely be formed under the Water Resources Management Act, Act No. 11 of 2013. This will likely give more powers to groundwater users in a basin to ensure sustainability of groundwater usage, but also encourage the optimal usage of groundwater. The project area falls inside a declared water control area and permits are required for drilling and rehabilitation of boreholes as well as for groundwater abstraction.

5.2 CLIMATE

The lack of functioning weather stations in Namibia, in especially rural areas, limits the availability of long term, true weather data. In the absence of weather station in the area, the best possible workaround is to use long term climate data obtained from the Atlas of Namibia Project (2002) and the CHIRPS-2 database (Funk et al., 2015), see Table 5-1 and Figure 5-1

The CHIRPS-2 dataset (Climate Hazards Group Infra-Red Precipitation with Station data version 2) consists of long term rainfall data (1981 to near-present) obtained from satellite imagery and in-situ station data. The remote sensing data provides average rainfall over a 25 km² area and such readings will be lower than some localised actual high volume rain events (thunderstorms). The resultant dataset provides a reasonably well represented overview of the climatic conditions and historic weather conditions of a general area. True values for single, site specific meteorological events may however differ to some degree.


According to the Köppen-Geiger Climate Classification system the project is located in a hot semi-arid climate (BSh) (<http://koeppen-geiger.vu-wien.ac.at/present.htm>). This means that the area receives precipitation below potential evapotranspiration, but not as low as a desert climate and has a mean annual temperature of at least 18 °C. Average rainfall received is 450-500 mm/a with a variation of 30-40 %. Monthly rainfall peaks in January. The potential evapotranspiration is 2,400 – 2,500 mm/a. By dividing the mean annual potential evapotranspiration into the mean annual precipitation, an aridity index value for the area was computed as 0.2, which indicates the area to be Semi-Arid. The average annual minimum temperature is 6-8 °C, while the average annual maximum temperature is 32-34 °C, with an average annual temperature range of 26-28 °C. An average diurnal temperature (difference between daily minimum and maximum temperature) for this area is around 16-18 °C. Direct normal solar irradiance for the area is 6.574 kWh/m²/day. (Atlas of Namibia Team, 2022)

The rain season normally starts in October and last until April, peaking in January and February. Heavier rainfall (single day events) occur between February and April, with a single event of 67 mm in Apr (last 40 years data) being the highest (Table 5-1). Most of the single day maximums are less than 50 mm. The average annual rainfall for the last 40 years was calculated as 464 mm/a, with a coefficient of variance of 27 %. Daily and seasonal rainfall data (Funk et al., 2015) is presented in Figure 5-1. Seasonal (July to June) total rainfall, centred on the average line for the last 40 years, is presented, with the daily total rainfall and the seasonal cumulative rainfall. From the figure it is clear that six out of the last ten seasons were much drier than usual.

Table 5-1 Rainfall statistics based on CHIRPS-2 data (Funk et al., 2015)

| Month | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|--------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Minimum (mm) | 12 | 30 | 20 | 5 | 0 | 0 | 0 | 0 | 0 | 0 | 8 | 17 |
| Maximum (mm) | 276 | 211 | 148 | 109 | 5 | 1 | 0 | 0 | 7 | 40 | 100 | 169 |
| Average (mm) | 112 | 109 | 80 | 29 | 1 | 0 | 0 | 0 | 1 | 15 | 41 | 76 |
| Variability (%) | 57 | 44 | 41 | 82 | 245 | 325 | 648 | 0 | 199 | 79 | 52 | 31 |
| Daily maximum (mm) | 44 | 53 | 44 | 67 | 5 | 1 | 0 | 0 | 6 | 19 | 31 | 46 |
| Average rain days | 14 | 12 | 8 | 3 | 0 | 0 | 0 | 0 | 1 | 4 | 7 | 11 |

Season July - June average: 464 mm | Season coefficient of variation: 27 %
Date range: 1981-July-1 to 2023-June-30 | Lat: 19.21037°S; Long: 18.00144°E

**Figure 5-1 Daily and seasonal rainfall from CHIRPS-2 data (Funk et al., 2015)*****Implications and Impacts***

Water is a scarce and valuable resource in Namibia and the extreme variability in seasonal rainfall makes water an extremely vulnerable resource. Rainfall events are typically thunderstorms with heavy rainfall that can occur in short periods of time (cloud bursts). Pollutants that enter the groundwater can pollute this valuable resource. Rainfall is important for groundwater recharge.

5.3 TOPOGRAPHY & DRAINAGE

The project area is located on the border between the Karstveld landscape to the south and the Kalahari Sandveld landscape to the north. Palaeo dunes and pans are characteristic of the Kalahari Sandveld while the Karstveld Landscape is dominated by limestone with little or no surface run-off and a strong development of sinkholes, dolines and caves. Locally the Karstveld Landscape is dominated by the Otavi Mountain Land, with hills rising up to 500 m above the surrounding plains.

Drainage and soils are influenced by topography which, for the project area, can be described as being level to gently undulating with a varying slope of between 0 % and 5 %. Drainage is poorly developed in the area. The site is located within the catchment of the Etosha Pan.




Figure 5-2 Aspect slope and surface drainage

Implications and Impacts

The lack of major surface runoff and drainage may lead to pooling and even flooding of plains during heavy rainfall events. This may negatively impact soil quality and cause localised flooding of infrastructure, if located in flood prone areas, or if such areas are not considered in designs. The risk of erosion is relatively low.

5.4 SOIL

Dominant soil type for this area is Cambic Calcisol which refers to the soil type commonly found in arid or semi-arid regions with dry seasons. They form in calcium and magnesium rich alluvial, colluvial and aeolian deposits and are alternately dampened by rain and dried by evaporation which results in soft masses or hard layers of calcrete. In addition to this, the calcisol of this particular area is known for having subsurface soil layers of pedogenic change without appreciable illuviated material. The composition of soil in this particular area is roughly 70-75 % sand, 10-15 % silt and 30-35 % clay which gives it the characteristics and texture of Loam soil. Bulk density was computed to be 1,450-1,500 mg/cm³ which means that the soil will affect the

root growth of various plants, but not necessarily restrict it. Soils in this area typically reach depths of 170-180 cm, have a pH of 5.5-6 and a cation exchange capacity of 7-10 cmol/kg. Furthermore, this region has a water capacity of 40-60 mm at root depth.

The farm is situated within the Kalk-2 Agro-Ecological Zone (AEZ) with an average growing period of 91 to 120 days. The Kalk-2 AEZ is ranked 2nd in Namibia in terms of agricultural potential and is deemed most suitable for short-maturing crops and large stock grazing. The Kalk-2 area is generally not regarded as suitable for cropping and this is true for some parts of the farm. The areas under irrigation are however located in patches where sufficiently deep, quality soil is present for irrigation of crops.




Figure 5-3 Dominant soil type and agro ecological zone

Implications and Impacts

Soil is considered to be shallow. The clay content is locally high as well as the organic content. The high clay content will reduce the infiltration rate during saturated condition but cracks in the topsoil during unsaturated conditions will form preferred infiltration pathways with high infiltration rates.

Soil seems to be suitable for irrigation.

5.5 GEOLOGY AND HYDROGEOLOGY

The geology underlying the project area formed during the Namibian, Quaternary and Tertiary Age. Locally the geology from the Quaternary and Tertiary Age comprises of the Kalahari Group deposits which consists of sand, calcrete and gravel. The Kalahari Group sediments originate mainly from fluvial deposition with some reworking through aeolian processes. Kalahari sediments at the project location form only a surface cover. The Kalahari Group sediments commonly overlie pre-Kalahari rocks of the Damara Sequence (Namibian Age). At the project location the Damara Sequence consists of dolostones belonging to the Hüttenberg Formation of the Tsumeb Subgroup, which forms part of the Otavi Group (Schneider, 2004). This group belongs to the Northern Platform of the Damara Sequence.

Moderate folding of the strata occurred during the Pan African Orogeny (680-450 Ma) and resulted in the formation of synclines and anticlines, generally trending east - west. To the north of the project area is the Aalborg Syncline, with the Adelaide Syncline to the south. The development of joints and fractures in the rocks are associated with the folding, which have an impact on the hydrogeological characterization of the area.

Various northeast striking magnetic dykes are known to be present in the subsurface, as inferred from aeromagnetic data. The dykes seem to be related to the Paresis intrusion which are situated just south of Otjiwarongo, with dykes radiating from this intrusion. These dykes are locally thought to have shattered the host rocks during its formation. Where dolomite is the host rock, it forms a zone favourable for the development of karst features and groundwater accumulation. At least one of these dykes cross the project area (Figure 5-4).

Figure 5-4 depicts geological structures interpreted from geophysical data for the project area and surroundings. An east-west trending fault is present in the centre of the project area and several north- south trending fault run across the western side of the project area.

Geophysical-interpreted dykes occur in the area and strike towards the northeast. The Remnant dyke intersects the project area. The nature of these dykes tend to be mineralised faults with high hydraulic conductivity values. Both the Tsumeb (further to the west) and Remnant dykes represented a major exploration target for the NamWater exploration water supply programme to Windhoek. The dykes are thought to have shattered the host rocks during its formation (Hoad, 1992). Where dolomite is the host rock, it forms a zone favourable for the development of karst features and groundwater accumulation.

Several known karst features are present in the region. The most well-known of these are the mineralised karst chimney of Abendab Mine ((Bäumle, 2003), as well as several lakes (Otjikoto and Guinas) and caves (e.g. Ghaub).

The Abenab and Abenab West Mines are located approximately 9 km to the southeast of the project area. This hydrothermal deposit represents a highly mineralized zone of which metals like vanadium as well as lead, copper and zinc were mined until 1948 and 1958 respectively when the ore reserves were depleted and the mines were closed (von Bezing, et.al., 2014). The Tsumeb Mine is approximately 29 km to the west; mining ceased in 1994.


A number of springs are present in the Otavi Mountain land and most of these springs are related to the contact zones between relatively impermeable formations of the Grootfontein Metamorphic Complex and more permeable formations of the Damara Sequence. The nearest of these contact zone springs is present approximately 11 km to the southeast of the project farm, see Figure 6-1. No caves or lakes are known of in close proximity (<10 km radius) to the project area.

The project area is situated in the Owambo Groundwater Basin. Localised groundwater flow may take place along preferred flow paths in different directions, but the larger scale groundwater flow is expected to be in a northern direction (Figure 5-5). Local flow patterns may vary due to groundwater abstraction. Groundwater flow is expected to take place through primary porosity in the surface cover, while it is expected to flow along fractures, faults, dykes/mineralised faults or along contact zones (secondary porosity) and other geological structures present within the underlying formations (hard rock formations). Contact zones in the area occur between permeable and impermeable formations and create favourable conditions to promote groundwater flow.

Groundwater quality data is presented in Figure 5-6 as Maucha plots. From the figure it is clear that the groundwater of the project location is mostly of a calcium-magnesium-bicarbonate type water which suggest the water is recently recharged. Groundwater quality from the project area reflect an aquifer that is typical of a dolomitic hard rock formation host where rapid groundwater recharge takes place.

Table 5-2 indicates the groundwater statistics for a radius of 5 km within the study area. The groundwater information was obtained from Department of Water Affairs (DWA) borehole database. The DWA database is generally outdated and more boreholes might be present. Groundwater is widely utilised in the study area, with 11 boreholes within a 5 km radius. The boreholes were drilled to an average depth of 71 m below surface and average yield of 9 m³/h. Groundwater quality falls mainly under Group A category, which indicates that the water is of an excellent quality, based on the provided parameters.

Table 5-2 Groundwater statistics

| Query Centre: Cuxhaven; -19.2104°S; 18.0014°E | | Query Box Radius: 5.0km | | | | | | | | | | |
|---|--|---------------------------|------------|-----------|-------------|---------------------------|-------------------|--------------------|-----------|----------------|---------------|----------------|
|  | | NUMBER OF KNOWN BOREHOLES | LATITUDE | LONGITUDE | DEPTH (mbs) | YIELD (m ³ /h) | WATER LEVEL (mbs) | WATER STRIKE (mbs) | TDS (ppm) | SULPHATE (ppm) | NITRATE (ppm) | FLUORIDE (ppm) |
| Data points | | 11 | | | 8 | 8 | 8 | 4 | 10 | 10 | 10 | 10 |
| Minimum | | | -19.165404 | 17.953751 | 46 | 3 | 21 | 30 | 477 | 6 | 0 | 0 |
| Average | | | | | 71 | 9 | 35 | 42 | 586 | 10 | 6 | 0 |
| Maximum | | | -19.255396 | 18.049049 | 119 | 18 | 46 | 50 | 789 | 28 | 14 | 0 |
| Group A | | | | | 12.50% | 25.00% | 0.00% | 0.00% | 100.00% | 100.00% | 60.00% | 100.00% |
| <i>Limit</i> | | | | | 50 | >10 | 10 | 10 | 1000 | 200 | 10 | 1.5 |
| Group B | | | | | 75.00% | 37.50% | 100.00% | 75.00% | 0.00% | 0.00% | 40.00% | 0.00% |
| <i>Limit</i> | | | | | 100 | >5 | 50 | 50 | 1500 | 600 | 20 | 2.0 |
| Group C | | | | | 12.50% | 37.50% | 0.00% | 25.00% | 0.00% | 0.00% | 0.00% | 0.00% |
| <i>Limit</i> | | | | | 200 | >0.5 | 100 | 100 | 2000 | 1200 | 40 | 3.0 |
| Group D | | | | | 0.00% | 0.00% | 0.00% | 0.00% | 0.00% | 0.00% | 0.00% | 0.00% |
| <i>Limit</i> | | | | | >200 | <0.5 | >100 | >100 | >2000 | >1200 | >40 | >3 |

Statistical grouping of parameters is for ease of interpretation, except for the grouping used for sulphate, nitrate and fluoride, which follow the Namibian guidelines for the evaluation of drinking-water quality for human consumption, with regard to chemical, physical and bacteriological quality. In this case the groupings has the following meaning:

Group A: Water with an excellent quality

Group B: Water with acceptable quality

Group C: Water with low health risk

Group D: Water with a high health risk, or water unsuitable for human consumption




Figure 5-4 Hydrogeological map

The aquifer associated with the project area falls within the Tsumeb-Abenab Synclinorium sub-catchment (Tsumeb - B2), previously part of the Karst Area IV aquifer zoning, which has a high hydraulic conductivity. Bäumlé (Bäumlé, 2003) described the Elandshoek Formation and the Hüttenberg Formation as the most productive aquifers within the investigation area, with the Tschudi Formation considered as an aquitard. Pumping test analysis yields an average transmissivity of 1,725 m²/d for the Hüttenberg and typical storage coefficients in the order of 5 x 10⁻⁴ to 1 x 10⁻³. Although high volume abstraction currently takes place in the Otavi Mountain Land, the only significant cones of depression known to exist were at the Tsumeb Mine (Hoad, 1992) and at the Kombat Mine, which is situated much further to the west and southwest.

During the peak activities of the Tsumeb Mine, the water level was decreased to a depth of about 1,700 m. Groundwater was abstracted on average at 500 m³/h to 600 m³/h and during peak times at 1,000 m³/h. This abstraction lasted for several decades, with a stable cone of depression that developed at a radius of approximately 2 km around the mine shaft (GKW Consult, et. al., 2003).




Figure 5-5 Groundwater basin with rainfall and inferred groundwater flow




Figure 5-6 Groundwater quality

Implications and Impacts

Local groundwater recharge is influenced by a thin veneer of Kalahari deposit, which might influence the sustainability over time. Recharge from the Otavi Mountain Land further to the south is likely the main source of groundwater recharge.

6 ASSESSMENT OF WATER LEVEL MONITORING DATA

Regional water level monitoring data was sourced from the Ministry of Agriculture, Water and Land Reform. A selection of monitoring borehole data was made, roughly following the inferred groundwater flow path, see Figure 6-1. Boreholes WW25922 and WW82463 are located close to the groundwater divide. Boreholes WW25922, WW82463 WW22024, WW27460 and WW200200 are in the more mountainous area, with boreholes WW200203, WW200224 and WW32627 in the flatter plain area. The selected boreholes present a range of boreholes stretching from just south of the water divide, past the project farm and beyond, roughly following the general groundwater flow direction (i.e. from near the water divide and past and beyond the farm). Figure 6-2 presents a cross section of the terrain as well as the minimum, average and maximum water levels of the boreholes. See Figure 6-1 for the profile location and Figure 6-3 for water level information of the relevant boreholes. Figure 6-2 indicates that the boreholes closer to the water divide shows larger groundwater level fluctuations and that the

fluctuations becomes insignificant as one moves away from the water divide. The project area is located between WW200200 and WW200203, suggesting the area would have little water level fluctuations.

In Figure 6-3 it is evident that most of the water level data has a stable water level profile. Rapid increases in water level are mostly associated with monthly rainfall exceeding 150 mm near the project area. These increases are subsequently followed by a steady decrease in water level. The most dramatic increases in water level is observed in data from WW82463 and WW25922, which is located close to the water divide where recharge is considered to be comparatively high. Careful monitoring is however required to properly manage this resource especially with increasing abstraction from the area.




Figure 6-1 Monitor borehole locations, caves, sinkholes and springs




Figure 6-2 Regional water level profile




Figure 6-3 Regional water level changes and monthly rainfall

7 WATER SUPPLY

7.1 GROUNDWATER USAGE

During the recognisance site visit all known boreholes on the farm were documented. Twelve boreholes were visited and data gathered about their status, use and physical description. A number of the boreholes are known DWA boreholes which are sealed. GPS locations of all boreholes were recorded and presented in Table 7-1. Of the boreholes surveyed, two are used for irrigation purposes, two are used for domestic purposes and two are used for stock watering.

The Proponent has a water abstraction permit for 120,000 m³ per year. Renewal of the permit is planned with no immediate increase in the allocation.

Table 7-1 Summary of borehole information obtained from the Proponent

| Map Ref. | Farm Name | Borehole Name(s) | Use | Borehole Depth (m) | Yield (m ³ /h) | Water Level (mbs) |
|----------|----------------------|----------------------------|----------------|--------------------|---------------------------|-------------------|
| EB1 | Cuxhaven Oos No.1278 | WW38010 (Klein pos gat) | Stock watering | 80 | 15 | |
| EB 2 | Cuxhaven Oos No.1278 | WW100065 | DWA BH | 200 | | 31 |
| EB 3 | Cuxhaven Oos No.1278 | | Not used | | 6 | 30 |
| EB 4 | Cuxhaven Oos No.1278 | | Not used | | | 31 |
| EB 5 | Cuxhaven Oos No.1278 | | Stock watering | 80 | 2 | |
| EB 6 | Cuxhaven Oos No.1278 | WW100071 | DWA BH | 200 | 8 | 45 |
| EB 7 | Cuxhaven Oos No.1278 | WW100069 | Irrigation | 200 | 300 | 48 |
| EB 8 | Cuxhaven Oos No.1278 | | Not used | 140 | 60 | 50 |
| EB 9 | Cuxhaven Oos No.1278 | WW 38011 (Huis gat) | Domestic | 100 | 20 | |
| EB 10 | Cuxhaven Oos No.1278 | WW39436 | Not used | 100 | | 42 |
| EB 11 | Cuxhaven Oos No.1278 | WW41033 | Irrigation | 180 | 50 | 73 |
| EB 12 | Cuxhaven Oos No.1278 | | Domestic | 100 | 30 | 49 |




Figure 7-1 Locations of boreholes



Photo 7-1 Borehole pump installation used for irrigation (EB11)



Photo 7-2 Borehole pump installation used for irrigation (WWW100069 / EB7)



Photo 7-3 Stock watering borehole (EB5)



Photo 7-4 DWA Borehole (WW100071 / EB13)

Implications and Impacts

Groundwater is a valuable resource in the farming area and is controlled by a water abstraction licensing system as regulated by the Ministry of Agriculture, Water and Land Reform. Groundwater contamination may negatively impact surrounding boreholes and groundwater is widely utilised for public water supply. No alternative water supply options exist if extensive contamination or deterioration of groundwater occur.

8 ASSESSMENT OF IMPACTS

The purpose of this section is to assess and identify the most pertinent environmental impacts and provides possible mitigation measures that are expected from the project. The Rapid Impact Assessment Method (Pastakia, 1998) will be used during the assessment. Impacts are assessed according to the following categories: Importance of condition (A1); Magnitude of Change (A2); Permanence (B1); Reversibility (B2); and Cumulative Nature (B3) (see Table 8-1).

The Environmental Classification = $A1 \times A2 \times (B1 + B2 + B3)$, see Table 8-2.

The probability ranking refers to the probability that a specific impact will happen following a risk event. These can be improbable (low likelihood); probable (distinct possibility); highly probable (most likely); and definite (impact will occur regardless of prevention measures).

Table 8-1 Assessment criteria

| Criteria | Score |
|---|--------------|
| Importance of condition (A1) – assessed against the spatial boundaries of human interest it will affect | |
| Importance to national/international interest | 4 |
| Important to regional/national interest | 3 |
| Important to areas immediately outside the local condition | 2 |
| Important only to the local condition | 1 |
| No importance | 0 |
| Magnitude of change/effect (A2) – measure of scale in terms of benefit / detriment of an impact or condition | |
| Major positive benefit | 3 |
| Significant improvement in status quo | 2 |
| Improvement in status quo | 1 |
| No change in status quo | 0 |
| Negative change in status quo | -1 |
| Significant negative detriment or change | -2 |
| Major detriment or change | -3 |

| | |
|---|---|
| Permanence (B1) – defines whether the condition is permanent or temporary | |
| No change/Not applicable | 1 |
| Temporary | 2 |
| Permanent | 3 |
| Reversibility (B2) – defines whether the condition can be changed and is a measure of the control over the condition | |
| No change/Not applicable | 1 |
| Reversible | 2 |
| Irreversible | 3 |
| Cumulative (B3) – reflects whether the effect will be a single direct impact or will include cumulative impacts over time, or synergistic effect with other conditions. It is a means of judging the sustainability of the condition – not to be confused with the permanence criterion. | |
| Light or No Cumulative Character/Not applicable | 1 |
| Moderate Cumulative Character | 2 |
| Strong Cumulative Character | 3 |

Table 8-2 Environmental classification of impacts (Pastakia 1998).

| Environmental Classification (ES) | Class Value | Description of Class |
|-----------------------------------|-------------|-------------------------------|
| 72 to 108 | 5 | Extremely positive impact |
| 36 to 71 | 4 | Significantly positive impact |
| 19 to 35 | 3 | Moderately positive impact |
| 10 to 18 | 2 | Less positive impact |
| 1 to 9 | 1 | Reduced positive impact |
| 0 | -0 | No alteration |
| -1 to -9 | -1 | Reduced negative impact |
| -10 to -18 | -2 | Less negative impact |
| -19 to -35 | -3 | Moderately negative impact |
| -36 to -71 | -4 | Significantly negative impact |
| -72 to -108 | -5 | Extremely Negative Impact |

8.1 GROUNDWATER ABSTRACTION

Groundwater abstraction is a very sensitive topic in a dry country where the value of land is drastically reduced if no or poor quality groundwater is present on the land. Abstraction of groundwater must be done in a sensible way not to impact on other groundwater users that depend on such groundwater. This includes water abstracted for human and animal use, irrigation, and also ecosystems that depend on groundwater. A typical groundwater balance was compiled to illustrate the potential consequences of over abstraction of groundwater, see Figure 8-1. Recharge to the area is considered to be high. It is considered that recharge can vary from 0 % to 4 % of rainfall with an average of 2 % of the rainfall. In periods of drought there may be no recharge while in above average rainfall recharge could be above 4 % (Hoad, 1992).

In a typical groundwater environment, a water balance would consist of inflow and outflow of the groundwater system. Over time an equilibrium (or steady state) is normally reached with rising water tables following good recharge events and declining water tables when recharge is below average.

Inflow into the system would typically be from infiltration following rainfall in the area and in upstream areas. The inflow component will further be enhanced by the high secondary porosity nature of the karst aquifer.

Outflow would be comprised of water leaving the system through springs and as outflow over the lower boundary of the groundwater system as well as evapotranspiration losses. Groundwater abstraction from boreholes is important as this is normally necessary to sustain human and animal demands where such users became essentially dependant on the abstracted groundwater as a reliable and sustainable source.

Typical consequences of over abstraction will include a lowering in the water table. This may lead to the collapse of underground cave roofs where the hydrostatic pressure, used to support the roof of a cave, decrease. The increased flow of water may enhance the dissolution of dolomitic rock, leading to an increase in karst structures. Lowering of water tables may further lead to the drying up of boreholes, springs, underground caves and the subsequent loss of organisms that lives in the subsurface and surface water. Vegetation will also be impacted where such vegetation has access to groundwater.

Based on current water level fluctuations in the area, a short term threshold of 5 m below the long term average water level is set from where abstraction rates should be reduced. Note that this level refers to rest water levels and not pump water levels.

All boreholes should be equipped with a dipper pipe to enable safe water level measurements.




Figure 8-1 Conceptual groundwater balance with over abstraction scenario

Table 8-3 Assessment – Groundwater abstraction

| Project Activity/Resource | Nature (Status) | (A1) Importance | (A2) Magnitude | (B1) Permanence | (B2) Reversibility | (B3) Cumulative | Environmental Classification | Class Value | Probability |
|---------------------------|--|-----------------|----------------|-----------------|--------------------|-----------------|------------------------------|-------------|-------------|
| Daily Operations | Over-abstraction of the local aquifer, decrease in the local hydraulic head. | 2 | -2 | 2 | 2 | 2 | -24 | -3 | Probable |

Desired Outcome: To utilise the groundwater sustainably.

Actions

Prevention:

- ◆ Spread the water abstraction points over a larger area to diffuse the impact.
- ◆ Monthly water level monitoring.
- ◆ Maintain safe abstraction rates prescribed by test pump evaluations (an abstraction permit with prescribed rates from the MAWLR is a requirement for this project).

Mitigation:

- ◆ Reduce abstraction when the water levels nears 5 m below the average rest water level of each borehole.

Responsible Body:

- ◆ Proponent

Data Sources and Monitoring:

- ◆ Monthly boreholes rest water level monitoring.
- ◆ Baseline values should be reviewed every three years based on all historic water level data.
- ◆ A summary report on all monitoring results must be prepared.
- ◆ The Proponent supply monitoring returns to the MAWLR, as required by the permit.

8.2 GROUNDWATER, SURFACE WATER AND SOIL CONTAMINATION

Leakages and spillages of hazardous substances from vehicles, waste oil handling and accidental fuel, oil or hydraulic fluid spills during the operational phase may contaminate the environment. Increase of nutrient levels (from over application of fertilizers or pesticides) in the soil that can leach to the groundwater. Pollution due to sewerage system overflow or leakage may further put the groundwater at risk.

Table 8-4 Assessment – Groundwater, surface water and soil contamination

| Project Activity/Resource | Nature (Status) | (A1) Importance | (A2) Magnitude | (B1) Permanence | (B2) Reversibility | (B3) Cumulative | Environmental Classification | Class Value | Probability |
|---------------------------|---|-----------------|----------------|-----------------|--------------------|-----------------|------------------------------|-------------|-------------|
| Daily Operations | Hazardous material, spillages, hydrocarbon leakages from vehicles and machinery. | 2 | -1 | 2 | 2 | 1 | -10 | -2 | Improbable |
| Daily Operations | Over application of fertilizer, pesticides, herbicides, etc. Sewerage system malfunction. | 2 | -1 | 2 | 2 | 1 | -10 | -2 | Improbable |

Desired Outcome: To prevent the contamination of groundwater, surface water and soil.

Actions

Prevention:

- ◆ Appoint reputable contractors.
- ◆ Vehicles may only be serviced on a suitable spill control structure.
- ◆ Regular inspections and maintenance of all vehicles to ensure no leaks are present.
- ◆ All hazardous chemicals and fuel should be stored in a sufficiently bunded area, as per MSDS requirements.
- ◆ Ensure all waste oil handling is conducted on impermeable or bunded areas.
- ◆ Follow prescribed dosage of fertilizers and pesticides / herbicides and to avoid over application.
- ◆ Maintain sewerage systems and conduct regular monitoring.
- ◆ All hazardous waste must be removed from the site and disposed of timeously at a recognised hazardous waste disposal facility, including any polluted soil or water.

Mitigation:

- ◆ All spills must be cleaned up immediately.
- ◆ Consult relevant Material Safety Data Sheet (MSDS) information and a suitably qualified specialist where needed.

Responsible Body:

- ◆ Proponent
- ◆ Contractors

Data Sources and Monitoring:

- ◆ Maintain Material Safety Data Sheets for hazardous chemicals.
- ◆ Soil should be sampled and analysed annually to ensure the correct amounts of fertilizer is applied and soil and groundwater quality is maintained.
- ◆ Groundwater should be sampled and analysed to test for nitrate concentrations from the fertilizer and for traces of chemicals used in pesticides and herbicides.
- ◆ Registers be kept by the Proponent on the type, quantities and frequency of application of fertiliser, pesticides and any other chemicals utilised in crop production.
- ◆ A register of all incidents must be maintained on a daily basis. This should include measures taken to ensure that such incidents do not repeat themselves.
- ◆ All spills or leaks must be reported on and cleaned up immediately.

9 CONCLUSION

Groundwater on the farms is high yielding and of acceptable quality for human consumption. Most boreholes are utilisation for irrigation purposes, although care must be exercised when long term irrigation takes place and nitrate values should be monitored regularly.

Based on current water level fluctuations in the area, as presented in Figure 6-3, a short term threshold of 5 m below the long term average water level is set from where abstraction rates should be reduced. This threshold may require adjustment during drought periods as abstraction from neighbouring farms may also influence the regional water levels. Careful cooperation between neighbouring farms and beyond is required to optimally utilize the groundwater resource without depleting it as depletion will be detrimental to all. This should include self-monitoring and assessment of water levels in the area as data obtained from DWA indicates a lack of sufficient monitoring in the recent years. Proper monitoring data will provide the required information to make informed decisions and will assist to obtain increased abstraction volume permits when needed and if justified.

Groundwater vulnerability to contamination would be the highest around boreholes, around geological structures as well as where shallow groundwater is present. Contaminated surface runoff can create a pathway to the groundwater, putting the groundwater at risk. Potential sources of groundwater pollution include normal runoff from roofs, properties and surfaced areas, e.g. roads. These impacts are normally of a low magnitude and can be managed through proper housekeeping.

Based on current water level and abstraction volumes continuous monitoring is recommended to determine if higher abstraction volumes may be considered. Large scale abstraction from the Abenab Mine for Windhoek may impact on the irrigation potential of farms in the area and thus lead to a reduction in property prices, reduced income and pose a risk to food security in Namibia.

10 REFERENCES

- Atlas of Namibia Team, (2022), Atlas of Namibia: its land, water and life, Namibia Nature Foundation, Windhoek.
- Bäumle R., (2003), Geohydraulic Characterisation of Fractured Rock Flow Regimes -Regional Studies in Granite (Lindau, Black Forest, Germany) and Dolomite (Tsumeb Aquifers, Northern Namibia).- Doctorate Thesis at the University of Karlsruhe, 129 pp.; Karlsruhe.
- Bäumle, R., (2004), Report Compiled by the Geohydrology Division. Revised Criteria to Consider when Allocating Licences for the Abstraction Of of Groundwater for Irrigation Purposes in The Tsumeb-Grootfontein-Otavi Subterranean Water Control Area, MAWF.
- Funk, C., Peterson, P., Landsfeld, M., Pedreros, D., Verdin, J., Shukla, S., Husak, G., Rowland, J., Harrison, L., Hoell, A. and Michaelsen, J., (2015), The climate hazards group infrared precipitation with stations - A new environmental record for monitoring extremes. *Scientific Data*, 2, 150066.
- Hoad, N., (1992), Resource Assessment and Development Potential of Contrasting Aquifers in Central Northern Namibia. Submitted as partial fulfilment of the M.Sc. Hydrogeology Course University College London.
- Laukamp, C., (2006), Dissertation: Structural and Fluid System Evolution in the Otavi Mountainland (Namibia) and its Significance for the Genesis of Sphide and Nnsulphide Mineralisation. Heidelberg: Ruprecht-Karls-Universität.
- Miller, R. M., (2008), The Geology of Namibia Volume 2 Neoproterozoic to Lower Palaeozoic. Windhoek: Geological Survey.
- Schneider, G., (2004), The Roadside Geology of Namibia. Stuttgart: Gebrüder Borntraeger.
- von Bezing, L., Bode, R., & Jahn, S., (2014), Namibia Minerals and Localities Volume 1. Bode Verlag GmbH.

Appendix C: GM Maize and Cotton in Namibia Specialist Report

ENVIRONMENTAL RELEASE OF GENETICALLY MODIFIED MAIZE AND COTTON FOR AGRICULTURAL PURPOSES IN NAMIBIA




Assessed by:

Assessed for:



Agricultural Industry

October 2023

| | | |
|-------------------------------|--|---|
| Project: | ENVIRONMENTAL RELEASE OF GENETICALLY MODIFIED MAIZE AND COTTON FOR AGRICULTURAL PURPOSES IN NAMIBIA: SPECIALIST ASSESSMENT | |
| Report Version/Date: | Final October 2023 | |
| Lead Consultant | Geo Pollution Technologies (Pty) Ltd PO Box 11073 Windhoek Namibia | TEL.: (+264-61) 257411 FAX.: (+264) 88626368 |
| Main Project Team: | André Faul (B.Sc. Zoology/Biochemistry); (B.Sc. (Hons) Zoology); (M.Sc. Conservation Ecology); (Ph.D. Medical Bioscience) | |
| Cite this document as: | Faul A. 2023. Environmental Release of Genetically Modified Maize and Cotton for Agricultural Purposes in Namibia: Strategic Environmental Assessment Report | |
| Copyright | Copyright on this document is reserved. No part of this document may be utilised without the written permission of Geo Pollution Technologies (Pty) Ltd. | |
| Report Approval |  André Faul Environmental Scientist | |

EXECUTIVE SUMMARY

Introduction

Stakeholders in the agricultural sector intend to apply for the registration of genetically modified (GM) maize for environmental release in Namibia. Under the Biosafety Act and Environmental Management Act of Namibia, the environmental release of a GM organism requires an environmental risk assessment to be conducted. This document reports on an assessment conducted by Geo Pollution Technologies (Pty) Ltd for the environmental release of GM maize and cotton. The specific GM events for maize are MON 810, MON 89034, NK 603 and stacks (combinations) of these events, and for cotton MON 88913 and the stacked event MON 88913 × MON 15985.

These maize and cotton events have primarily been designed to provide insect and / or herbicide resistance. Insect resistance is targeted at the fall armyworm and African maize stalk borer in maize, and the African boll worm in cotton. These pests can cause significant crop losses within days of infestation in traditional non-GM maize and cotton fields. In insect resistant events, moth larvae are controlled by specific proteins that were introduced into the maize and cotton through genetic engineering. Herbicide resistant maize and cotton are resistant to the systemic, non-selective herbicide glyphosate. This enables farmers to manage all weeds in maize and cotton fields by applying glyphosate without harming the maize and cotton itself.

Scope and Methodology

A specialist assessment report was prepared by conducting an extensive literature review and interviewing experts in the field of agricultural economics, specifically in the Namibian environment. The report addresses both GM maize and GM cotton and can thus be used as literature source in environmental impact assessments for farmers' who wish to cultivate GM maize and/or GM Cotton.

Literature Review and Aspects of GM Maize and Cotton Cultivation

A vast amount of scientific and unscientific (popular) publications are available. To separate fact from myth requires in-depth consideration of various publications. A number of expert scientific reviews on the topic of genetically modified organisms (GMOs) are available. The most recent of these covering two decades worth of literature and data.

The main concerns related to the cultivation of GMOs in general are the potential health effects they may have on the consumers as a result of their changed genetic composition, and the potential impact on biodiversity as a result of their environmental release.

Based on the review of existing scientific literature, no concrete evidence could be found that substantiate the various claims of negative impacts caused by GMOs. What became evident is that many anti-GMO lobbyists portray GMOs in a negative light without critical examination of the existing scientific data. Some of these campaigns have been so successful that amidst a severe shortage of food, Zambia's government refused a consignment of food aid consisting of GM maize. Thus far, the only real argument that has some scientific credibility pertaining to negative impacts of GM crops, is that insect and weed resistance can develop in light of the designed GM traits. However, this is not more so than resistance development in conventional non-GM maize (and other crop) cultivation activities.

Conclusion

Economically, the cultivation of GM maize and cotton have been shown, more often than not, to be more profitable and higher yielding (especially for insect resistant crops), than its non-GM counterpart. This is evident in the complete adoption of GM cotton in South Africa with no traditional cotton being planted anymore. The profitability and yields also increase significantly during years of significant pest infestations. In a country like Namibia, with mostly marginal agronomic potential, and likely to be significantly affected by climate change, it makes sense to diversify agronomic practices by introduction GM crops into the system. This assessment report will guide the implementation process and provide a framework within which adopters of GM maize and cotton for cultivation must operate. It remains the responsibility of each farmer to perform the necessary calculations to establish feasibility of GM maize and cotton cultivation for his / her specific circumstances.

TABLE OF CONTENTS

| | | |
|----------|--|-----------|
| 1 | BACKGROUND AND INTRODUCTION | 1 |
| 2 | OBJECTIVES | 1 |
| 3 | NEED AND DESIRABILITY | 1 |
| 4 | LITERATURE REVIEW | 2 |
| 4.1 | GENETICALLY MODIFIED ORGANISMS | 2 |
| 4.1.1 | <i>Selective Breeding</i> | 2 |
| 4.1.2 | <i>Brief History of Genetically Modified Organisms</i> | 3 |
| 4.1.3 | <i>Genetic Engineering Methodology</i> | 4 |
| 4.1.4 | <i>Global Status of Genetically Modified Crop Production</i> | 4 |
| 4.2 | GENETICALLY MODIFIED MAIZE FOR AGRICULTURE IN NAMIBIA | 7 |
| 4.2.1 | <i>Event MON 810</i> | 7 |
| 4.2.2 | <i>Event MON 89034</i> | 8 |
| 4.2.3 | <i>Event NK 603</i> | 8 |
| 4.2.4 | <i>Gene Stacked Events</i> | 8 |
| 4.3 | GENETICALLY MODIFIED COTTON FOR AGRICULTURE IN NAMIBIA | 8 |
| 4.3.1 | <i>Event MON 88913</i> | 9 |
| 4.3.2 | <i>Event MON 88913 x MON 15985</i> | 9 |
| 4.4 | ASPECTS OF CULTIVATING GM MAIZE AND COTTON | 9 |
| 4.4.1 | <i>Genetically Modified Crop Yield</i> | 9 |
| 4.4.2 | <i>Pesticide Use</i> | 10 |
| 4.4.3 | <i>Comparison of Costs and Benefits</i> | 10 |
| 4.4.4 | <i>Trade and Marketing Issues</i> | 16 |
| 4.4.5 | <i>Biodiversity</i> | 17 |
| 4.4.6 | <i>Bt Toxin Resistance</i> | 18 |
| 4.4.7 | <i>Herbicide Resistance</i> | 19 |
| 4.4.8 | <i>Gene Flow</i> | 20 |
| 4.4.9 | <i>GMOs as Food and Livestock Feed</i> | 21 |
| 4.5 | GENETICALLY MODIFIED MAIZE AND COTTON IN SOUTH AFRICA | 22 |
| 5 | ALTERNATIVES | 23 |
| 5.1 | <i>No Go ALTERNATIVE</i> | 24 |
| 6 | ADMINISTRATIVE, LEGAL AND POLICY REQUIREMENTS | 24 |
| 7 | IDENTIFIED IMPACTS | 30 |
| 7.1 | ECONOMIC | 30 |
| 7.1.1 | <i>Employment</i> | 30 |
| 7.1.2 | <i>Economic Resilience</i> | 31 |
| 7.1.3 | <i>Yield and Revenue</i> | 32 |
| 7.1.4 | <i>Meat Exports</i> | 33 |
| 7.2 | PHYSICAL / CHEMICAL | 34 |
| 7.2.1 | <i>Pesticides in Soil and Groundwater</i> | 34 |
| 7.2.2 | <i>Soil Erosion</i> | 35 |
| 7.3 | BIOLOGICAL | 36 |
| 7.3.1 | <i>Pesticides Resistance</i> | 36 |
| 7.3.2 | <i>Biodiversity / Non-Target Species</i> | 37 |
| 7.3.3 | <i>GM Crops Becoming Invasive</i> | 38 |
| 7.3.4 | <i>Horizontal Gene Transfer</i> | 39 |
| 7.4 | SOCIAL | 40 |
| 7.4.2 | <i>Feelings and Aspiration for the Future</i> | 41 |
| 7.4.3 | <i>Social Cohesion</i> | 42 |
| 7.4.4 | <i>Community Health and Welfare</i> | 43 |
| 7.4.5 | <i>Cultural Aspects</i> | 45 |
| 8 | PERMIT APPLICATION | 46 |
| 9 | CONCLUSION | 46 |

| | |
|----------------------------|-----------|
| 10 REFERENCES | 47 |
|----------------------------|-----------|

LIST OF APPENDICES

| | | |
|-------------|--|----|
| APPENDIX A: | NEWSPAPER CLIPPINGS | 52 |
| APPENDIX B: | COST GUIDE FIGURES FOR BT MAIZE AND NON-GMO MAIZE | 56 |
| APPENDIX C: | COST GUIDE FIGURES FOR GM COTTON: DRY-LAND VS. IRRIGATED | 65 |
| APPENDIX D: | CONSULTANT’S CURRICULUM VITAE..... | 70 |

LIST OF FIGURES

| | | |
|------------|--|----|
| FIGURE 4-1 | MAIZE PRODUCTION TREND IN SOUTH AFRICA OVER THE LAST 50 YEARS (SOURCE: ESTERHUIZEN & CLADWELL, 2021) | 6 |
| FIGURE 4-2 | SOUTH AFRICAN AVERAGE MAIZE YIELD/HA INCREASE (SOURCE: ESTERHUIZEN & CLADWELL, 2021)..... | 6 |
| FIGURE 4-3 | SCHEMATIC REPRESENTATION OF PESTICIDE RESISTANCE DEVELOPMENT (SOURCE: IRAC 2011)..... | 19 |
| FIGURE 4-4 | GLOBAL WEED RESISTANCE (HTTP://WWW.WEEDSCIENCE.ORG/GRAPHS/GEOCHART.ASPX) | 20 |

LIST OF TABLES

| | | |
|-----------|--|----|
| TABLE 4-1 | SUMMARY TABLE: COMPARATIVE PRODUCTION COSTS AND INCOMES FOR GM AND NON-GM MAIZE CULTIVARS UNDER KNOWN SOUTH AFRICAN CONDITIONS (2019 FIGURES USED) | 13 |
| TABLE 4-2 | SUMMARY TABLE: PRODUCTION COSTS AND INCOMES FOR GM COTTON CULTIVARS UNDER KNOWN SOUTH AFRICAN CONDITIONS (2022 AND 2023 DATA USED)..... | 14 |
| TABLE 5-1 | ALTERNATIVE MAIZE AND COTTON TYPES FOR CULTIVATION | 23 |
| TABLE 6-1 | NAMIBIAN LAW APPLICABLE TO GMOS | 25 |
| TABLE 6-2 | RELEVANT MULTILATERAL ENVIRONMENTAL AGREEMENTS..... | 28 |
| TABLE 6-3 | STANDARDS OR CODES OF PRACTISE | 29 |

LIST OF PHOTOS

| | | |
|-----------|---|---|
| PHOTO 4-1 | TEOSINTE EAR..... | 3 |
| PHOTO 4-2 | MAIZE EAR..... | 3 |
| PHOTO 4-3 | COMPARISON OF THREE SPECIES OF WILD AND DOMESTICATED COTTON (FROM HTTPS://FACULTY.SITES.IASTATE.EDU)..... | 3 |

LIST OF ABBREVIATIONS

| | |
|---------------|--|
| ATF | Namibian Agricultural Trade Forum |
| Bt | <i>Bacillus thuringiensis</i> |
| Btk | <i>Bacillus thuringiensis krustaki</i> |
| DDT | Dichlorodiphenyltrichloroethane |
| DNA | Deoxyribonucleic acid |
| EIA | Environmental Impact Assessment |
| EMA | Environmental Management Act No 7 of 2007 |
| EPSPS | Enolpyruvylshikimate-3-phosphate synthase |
| EU | European Union |
| FAO | Food and Agriculture Organization |
| GE | Genetically Engineered |
| GM | Genetically Modified |
| GMO | Genetically Modified Organism |
| HGT | Horizontal Gene Transfer |
| HIV | Human Immunodeficiency Virus |
| HT | Herbicide Tolerant |
| ISAAA | The International Service for the Acquisition of Agri-biotech Applications |
| ISPM | International Standards for Phytosanitary Measures |
| IUCN | International Union for the Conservation of Nature |
| LMO | Living Modified Organism |
| NASEM | National Academies of Sciences, Engineering, and Medicine |
| NAU | Namibia Agricultural Union |
| NBA | National Biosafety Authority |
| NCRST | National Commission on Research Science and Technology |
| NDP5 | Fifth National Development Plan |
| SADC | Southern African Development Community |
| SPS | Sanitary and Phytosanitary |
| Subsp. | Subspecies |
| UNFCCC | United Nations Framework Convention on Climate Change |
| US | United States |
| USA | United States of America |
| WHO | World Health Organization |
| WTO | World Trade Organization |

GLOSSARY OF TERMS

Actual Yield – The real tonnage/ha harvested, which typically are less than potential yield because of reducing factors, limiting factors and less than perfect conditions.

Assessment - The process of collecting, organising, analysing, interpreting and communicating information relevant to decision making.

Competent Authority - means a body or person empowered under the local authorities act or Environmental Management Act to enforce the rule of law.

Cumulative Impacts - in relation to an activity, means the impact of an activity that in itself may not be significant but may become significant when added to the existing and potential impacts eventuating from similar or diverse activities or undertakings in the area.

Environment - As defined in the Environmental Assessment Policy and Environmental Management Act - “land, water and air; all organic and inorganic matter and living organisms as well as biological diversity; the interacting natural systems that include components referred to in sub-paragraphs, the human environment insofar as it represents archaeological, aesthetic, cultural, historic, economic, palaeontological or social values”.

Environmental Release – For purposes of this document this means the release of genetically modified crops for controlled agricultural purposes.

Genetic Modification / Genetic Engineering – the process of altering the genetic material of an organism to produce a genetically modified organism.

Genetically Modified Organism - organisms whose genetic material (genome) has been artificially altered, through genetic engineering, to express favourable physiological traits or produce desired biological products.

Herbicide Resistance – The ability of a plant, typically referring to weeds, to withstand the effects of a herbicide.

Horizontal Gene Transfer – The transfer of genetic material between single cell and / or multicellular organisms where offspring is not produced.

Insect Resistance – The ability of a plant to resist insect damage either through natural means or as a result of genetic modification.

Mitigate - The implementation of practical measures to reduce adverse impacts.

Potential Yield - The maximum tonnage/ha that a crop can produce given no reducing factors (weeds, pests, diseases, etc.), an abundance of water and nutrients, and optimum carbon dioxide levels, radiation, temperature, etc.

Significant Effect/Impact - means an impact that by its magnitude, duration, intensity or probability of occurrence may have a notable effect on one or more aspects of the environment.

Sustainable Development - “Development that meets the needs of the current generation without compromising the ability of future generations to meet their own needs and aspirations” – the definition of the World Commission on Environment and Development (1987). “Improving the quality of human life while living within the carrying capacity of supporting ecosystems” – the definition given in a publication called “Caring for the Earth: A Strategy for Sustainable Living” by the International Union for Conservation of Nature (IUCN), the United Nations Environment Programme and the World Wide Fund for Nature (1991).

Tolerance – The ability of a plant or animal to tolerate a pesticide. Often used interchangeably with resistance.

1 BACKGROUND AND INTRODUCTION

Worldwide, vast expanses of land has been cleared to make way for crop farming to produce food and other agricultural products. With the human population exceeding eight billion, traditional methods of practising agriculture are struggling to meet the demand for food. This is further exacerbated by climate change impacts on rainfall and desertification. Therefore, the agricultural sector continuously investigates and apply increasingly intensive farming methods, to maximise yield and profitability per farming unit. Modern biotechnology has the potential to revolutionize the agricultural industry by developing genetically modified organisms (GMOs) that, due to specific engineered traits, can increase yields and profits while simultaneously simplifying crop cultivation.

Agriculture is one of the key economic sectors in Namibia and one of the major contributors to employment. To meet the growing demand for maize for food and feed production as well as cotton, it is the intention of some farmers to register genetically modified (GM) maize and cotton for environmental release in Namibia. To achieve this, such farmers must apply for permission from the Biosafety Council, of the National Commission on Research, Science and Technology (NCRST), to cultivate GM maize in Namibia. To allow for the registration of GMOs in Namibia, an environmental impact assessment, and an associated management plan, is required as per the Environmental Management Act (EMA) of Namibia (Act No. 7 of 2007). The GM maize earmarked for registration expresses three different genetically engineered (GE) traits, being 1) insect resistance (Mon 810 and Mon 89034), 2) glyphosate resistance (NK 603), and 3) both insect and glyphosate resistance (Mon 810 × NK 603 and Mon 89034 × NK 603). The GM cotton earmarked for registration is 1) glyphosate resistance (MON 8913) and 2) glyphosate and insect resistance (MON 88913 × MON 15985).

2 OBJECTIVES

The main objective of this study is to provide sufficient information to feed into environmental impact assessments for individual farmers who wish to cultivate GM maize and cotton. This will be achieved by:

1. Providing a brief explanation of what constitutes a GMO.
2. Presenting a literature review on GM maize and cotton, the potential benefits, impacts and main concerns related to GM maize and cotton and GMOs in general.
3. Providing a summary of the legal and regulatory framework related to GMOs in Namibia.
4. Evaluating the potential environmental impacts that may result from the cultivation of the selected GM maize and cotton strains in Namibia.
5. Identifying a range of management actions to mitigate the potential adverse impacts to acceptable levels.

3 NEED AND DESIRABILITY

The Fifth National Development Plan of Namibia (NDP5) recognises the importance of the agricultural sector in Namibia. Currently agriculture supports approximately 70% of Namibians and provide employment to roughly a third of the workforce. The NDP5's desired outcome is to see a reduction in food insecurity through an increase in food production [agriculture]. A reduction in agricultural potential (yield) is however expected in light of climate change and desertification. In addition, the occurrence of periodic drought cycles drastically reduce agricultural productivity in Namibia. Therefore, technological advancements are required should Namibia wish to increase food production by means of agriculture.

Maize is one of the staple foods in Namibia and a key ingredient in many animal feed products. Due to the lack of rainfall, the commercial cultivation of maize is only feasible in selected areas, and on relatively small scale. As a result, Namibia is a net importer of maize. Local maize production volumes are dependent on rainfall (dryland cropping), sufficient volumes of stored water (groundwater and dams) and suitable soils. Cotton is an ideal small-scale cash crop in drier climates, due to its resilience under lower rainfall conditions. Yields of both maize and cotton are affected by the outbreak of pests like the fall armyworm and boll worm that can rapidly damage vast stands of maize and cotton

respectively. (See Appendix A for examples of newspaper articles making headlines on pests affecting the agriculture sector.).

Genetically modified crops have the ability to resist or withstand some of the obstacles in crop cultivation. This may result in various direct and indirect benefits and ultimately contribute to food and feedstuff security. Benefits of cultivating pest and herbicide resistant GM crops include:

- ◆ Increased actual yields leading to enhanced food and feedstuff security for local and international markets.
- ◆ Resilience in the agricultural sector.
- ◆ Increased income and thus spending power.
- ◆ Increased revenue paid to government.
- ◆ Decreased insecticide use.
- ◆ More convenient and potentially safer pest control.
- ◆ More time for additional income generating activities which can in turn lead to more employment.

4 LITERATURE REVIEW

In the first part of the literature review, a short explanation of the basics of GMOs is provided.

4.1 GENETICALLY MODIFIED ORGANISMS

The World Health Organisation (WHO) defines and explains GMOs as follows:

“Organisms (i.e. plants, animals or microorganisms) in which the genetic material (DNA) has been altered in a way that does not occur naturally by mating and/or natural recombination. The technology is often called “modern biotechnology” or “gene technology”, sometimes also “recombinant DNA [deoxyribonucleic acid] technology” or “genetic engineering”. It allows selected individual genes to be transferred from one organism into another, also between nonrelated species. Foods produced from or using GM organisms are often referred to as GM foods” (WHO 2014).

Genetically modified organisms are thus organisms whose genetic material (genome) has been artificially altered, through genetic engineering, to express favourable physiological traits or produce desired biological products. Genetic modification is not a new concept, the method however has changed significantly in the last four to five decades.

4.1.1 Selective Breeding

As far back as 30,000 years ago, people selectively bred wolves that shared similar favourable phenotypic traits. The result of this selective breeding is that the offspring is more likely to have the genes responsible for that specific trait. In turn, by selecting the offspring with the trait, and again breeding with them, increases the chances of the offspring containing those genes. As this process is repeated, a wolf with a different genotype and phenotype is eventually produced. This is exactly how the numerous dog breeds in existence today, originated (e.g. doberman, labrador, beagle, etc.). Their original ancestors were wolves, but their genotype, and thus phenotype, are now completely different. So much so that dogs are regarded as an entirely new species.

Maize and cotton are no exceptions when it comes to selective breeding. Originally, maize was a wild grass, teosinte, with tiny ears and very few kernels (Photo 4-1). Through selective breeding, dating back as far as 9,800 years, maize now produce large ears with many kernels (Photo 4-2). Cotton was also bred to have more and longer fibres than their wild relatives (Photo 4-3) (<https://faculty.sites.iastate.edu>).

Selective breeding is thus a slow process of changing the genome of an organism, in order to develop traits favourable to man. Other examples include the numerous colours in budgies, canaries and some parrots, seedless watermelons, larger fruits and vegetables, cattle better suited for specific environments, cows producing more milk, etc.



Photo 4-1 Teosinte ear




Photo 4-2 Maize ear




Photo 4-3 Comparison of three species of wild and domesticated cotton (from <https://faculty.sites.iastate.edu>)

4.1.2 Brief History of Genetically Modified Organisms

Although selective breeding also results in organisms that are genetically different (modified), “GMO” typically refers to the modern techniques of genetic engineering.

In 1973, the first GM bacteria was engineered (Cohen et. al. 1973) when scientists succeeded in “cutting” a gene from one strain of bacteria and “pasting” it into the genome of another bacterium. By 1974, the first GM mammal, a mouse, was engineered (Jaenisch and Mintz 1974). Eight years later, in 1982, the first medication produced by a GMO was approved for human use (Ladisch and Kohlmann 1992). In the latter case, bacteria was engineered to

synthesize insulin in large enough amounts to allow its purification and subsequent administration to patients. The insulin had the exact same structure as that produced in humans. By the early 1990's, the first commercially available GM tomato was placed on the market as food (Bruening and Lyons 2000). In 2017 GM salmon, the first GM animal approved as food, were placed on the market in Canada. GM animals for food production are however still controversial and generally not well received by the general public.

4.1.3 Genetic Engineering Methodology

The genetic modification of an organism is known as an “event”. An event can be a single modification or multiple modifications. Where multiple modifications are present it is referred to as a “gene stacked event”.

A variety of genetic engineering (GE) techniques exists. A lengthy and complete description / explanation of each of the technologies falls outside of the scope of this report. Instead, brief, non-technical descriptions of some of the techniques are provided as background information. The descriptions of the techniques were obtained from National Research Council (US) Committee on Identifying and Assessing Unintended Effects of Genetically Engineered Foods on Human Health (2004). What is important to know is that genetic code is a “universal language”, meaning one organism has the ability to read and encode the genes of almost all other organisms. It is as a result of this ability that GE is possible.

Microbial vectors – The plant disease causing *Agrobacterium tumefaciens* naturally occur in soil and has the special trait of being able to transfer a portion of its own DNA into a host plant cell. By doing so, it causes gall disease in susceptible plants. In the 1980s, *Agrobacterium* lacking the disease causing genes were developed, while still maintaining its ability to insert DNA into the host. Substituting the disease causing *Agrobacterium* DNA, with DNA from another species that expresses desirable traits, allows *Agrobacterium* to insert the “new” DNA into a host plant. The DNA is subsequently integrated into the host's cells. By growing a fertile plant from the modified plant cell, produces a plant that may express the desired trait. Since it is a universal language, the host plant will express the traits of the inserted gene by producing the proteins it codes.

Microprojectile Bombardment – With this method, DNA is attached to microscopic pellets, which are “shot” at plant cells. This way, DNA is inserted into the plant cell, and subsequently expressed.

Retroviral Vectors – Retroviruses are viruses able to transport their own genes into the cells they infect. The genes are then integrated with the host cells' genome. With retroviral vectors, certain genes of the virus are removed and replaced by the gene to be introduced into the host. When the virus delivers the new gene together with some enzymes to the host cell, the gene is integrated into the host, which can then express the desired trait. The virus therefore acts like a “Trojan horse”.

4.1.4 Global Status of Genetically Modified Crop Production

In general terms, the economic benefits of cultivating GMO crops are well-researched and well-known globally. Empirical evidence of the economic benefits has been available for decades. Countries that adopted GMO technology during the early years have proceeded to steadily increase the area under GM crop cultivation, as well as the number GM varieties grown in their territories. This trend still continues. At the same time, more and more countries are joining this trend by either lifting or relaxing previously introduced bans and restrictions on the importation of GMO food and feedstuffs and/or allowing the cultivation of GMO crops. The main crop producing and exporting countries have almost all adopted GMO technology and, as a result, have continued to expand their agricultural production base, as well as their overall agricultural output and exports.

Cotton was one of the first crops to be bio-engineered and adopted at a global level. It was much easier to accept the introduction of bio-engineered cotton (as a non-food crop) in

contrast to bio-engineered crops cultivated as foodstuffs (both for human and for animal consumption) such as maize, wheat and to a lesser extent, soybeans. Due to less controversy and sensitivity around crops that were not meant to be used as human or animal food, genetically modified cotton became the first crop that was accepted for introduction in farming systems on the African continent and have been cultivated in Africa as far back as the 1990s (Hofs, J.L. & Kirsten, J., Working Paper, 2001-17).

In a regional context, cotton was also the first genetically modified crop to be approved for commercial cultivation in South Africa and, at present, 100% of cultivated cotton in South Africa is from GM seeds. The economic and management benefits obtainable from the use of GM cotton has resulted in a situation where there is no longer any conventional cotton being planted in South Africa.

In 1996, 2.8 million hectares of GM crops were cultivated in the United States of America, China, Canada, Argentina, Australia and Mexico (ISAAA, 1997). By 2018, this figure has grown to 191.7 million hectares in 26 countries by approximately 17 million farmers (ISAAA, 2018). An additional 44 countries imported GMOs for food and feed purposes, which brings the total number of countries adopting GM crops to 70 (ISAAA, 2018). As of 2017, the top five countries growing GMOs in terms of crop area are the United States, Brazil, Argentina, Canada and India. It is also interesting to note that these countries are amongst the biggest organic agricultural producers in the world, along with China, Australia and the EU, which is in itself interesting as it shows the complementarities that do exist between the two fields of agricultural production. They are not mutually exclusive and can indeed co-exist and flourish in the same country. In 2019, the number of countries in Africa that have approved GM crops for food, field trials and/or environmental release doubled from three to six (ISAAA 2019) and by 2023, nine African countries are listed on the ISAAA website as having approvals for GM crops (<https://www.isaaa.org/>). South Africa, being the largest producer of GM crops with a total of 72 events approved for canola, cotton, maize, rice and soybeans (<https://www.isaaa.org/>). During the 2020/21 marketing year, 3.3 million hectares of land were cultivated with maize, cotton and soybeans of which approximately 2.8 million hectares were planted with GM variants (Esterhuizen & Cladwell, 2021). All cotton produced in South Africa in this period were GM variants while 85% of maize were GM variants.

The commercialisation of GM crops has occurred at a rapid rate since the mid-1990s, with important changes in both the overall level of adoption and impact occurring in 2016. Positive gains have been divided 48% to farmers in developed countries and 52% to farmers in developing countries. There continues to be very significant net economic benefits at the farm level amounting to US\$18.2 billion in 2016 and US\$186.1 billion for the period 1996–2016 (in nominal terms). PG Economics (2018) estimates that farmers in developing countries received US\$5 for each dollar invested in genetically engineered crop seeds in 2017. About 65% of the gains have derived from yield and production gains with the remaining 35% coming from cost savings.

Genetic engineering technology has also made important contributions to increasing global production levels of the four main crops, having, for example, added 213 million tonnes and 405 million tonnes respectively, to the global production of soybeans and maize since the introduction of the technology in the mid-1990s. Cultivating GMO crops has provided significant benefits to farmers globally, including increased yield and lower production costs. Importantly, GMOs also help to alleviate poverty for the millions of resource-poor farmers and farm families around the world. As countries look to expand their domestic GM product pipelines and crop production, even more farmers will have access to improved seeds and the benefits they provide (PG Economics, 2018).

South Africa and Sudan have had great successes with GM crops (Abdallah 2014; Pellegrino et al. 2018). South Africa is the ninth largest GM crop producing country in the world (Esterhuizen & Cladwell, 2021). South Africa's production of maize (non-GM and GM maize) increased over the last four decades while the area planted, decreased (Figure 4-1)

(Esterhuizen & Cladwell, 2021). Average maize yields per hectare increased from 2.2 tons per hectare to 4.5 tons per hectare since the adoption of GM maize (Figure 4-2) (Esterhuizen & Cladwell, 2021).




Figure 4-1 Maize production trend in South Africa over the last 50 years (source: Esterhuizen & Cladwell, 2021)




Figure 4-2 South African average maize yield/ha increase (source: Esterhuizen & Cladwell, 2021)

As mentioned, seventy-two GM events have approval for feed, food or environmental release in South Africa. Since the first GM crops were adopted in South Africa a shift in the perception of the public on GM crops and food has occurred. A public perception survey indicated that the understanding and awareness of biotechnology increased significantly between 2004 and 2015

(Gastrow et al. 2017). Forty-nine percent of the population believe GM food is safe to eat while 53% believe it is good for the economy. The increase in positive perception is largely attributable to increased education and knowledge on biotechnology.

4.2 GENETICALLY MODIFIED MAIZE FOR AGRICULTURE IN NAMIBIA

Namibia is a net maize (*Zea Mays*) importer, mostly from South Africa. White maize is one of Namibia's staple food grains while yellow maize are mostly used for animal feed. White maize is planted mainly as a dry-land crop, but also under irrigation where surface water (Hardap Dam, Kavango River) or groundwater is abundant. The main white maize cultivation areas are the maize triangle (Otavi – Grootfontein – Tsumeb), along the Kavango River (Green Schemes), Hardap Scheme and eastern Namibia around Hochfeld and Summerdown. Limited production of white maize also occur in the Zambezi and Omusati Regions. On communal farms maize production is mainly for own use.

The main pests encountered in the cultivation of maize in Namibia is the Lepidopterans (moths and butterflies) *Busseola fusca* (African maize stalk borer), *Spodoptera frugiperda* (fall armyworm) and to a lesser degree *Spodoptera exempta* (African army worm). All three are the larval stage of species of moths. The African maize stalk borer is native to sub-Saharan Africa while the fall armyworm is an alien invasive from the Americas, first encountered in Africa in 2016. The larval stages of these moths can cause massive destruction in maize fields if detected too late or if not actively controlled through pesticide application.

Weeds typically compete with a crop's resources and must in most case be actively managed. This can be achieved either by mechanical removal (tillage and manual labour like hoeing) or herbicide application. Herbicides can be non-selective or selective in nature. Non-selective herbicides will kill all plants it comes in contact with. Selective herbicides will selectively kill certain plants while not damaging others. Selectivity can be based on a plant's age or growing stage, morphology, absorption potential, etc. A feature often used in selective weed control is the difference between monocotyledonous (grasses) plants and dicotyledonous (broadleaf) plants. Certain herbicides will kill only broadleaf weeds while others target only grasses. Since maize is a monocotyledonous plant, herbicides controlling broadleaf plants can be sprayed onto post-emergent maize, but not herbicides for controlling grasses.

Existing GM maize events for agricultural purposes are insect resistance, glyphosate herbicide resistance, as well as both insect and glyphosate resistance. Namibian farmers wish to be granted permission to cultivate GM maize in order to reduce losses in maize production from pests and weeds, as well as reduce costs in cultivation of maize. The following sections discuss the specific events for which permission is required.

4.2.1 Event MON 810

Event MON 810 developed by Monsanto (now incorporated into Bayer) is marketed under the trade name YieldGard[®]. It is an insect resistant strain, specifically targeting the order Lepidoptera, which comprises of moths and butterflies (and their larvae). It is engineered to express insecticidal toxins from the bacterium *Bacillus thuringiensis* subsp. *kurstaki*, commonly referred to as Btk. *B. thuringiensis* *krustaki* is a gram-positive, rod-shaped bacterium widely distributed in soil. In nature, Btk produces a delta-endotoxin with insecticidal properties against the orders Lepidoptera, Coleoptera (beetles), Hymenoptera (ants, wasps, bees and sawflies) and Diptera (true flies) as well as the phylum Nematoda (round worms). The endotoxin is in the form of parasporal crystals comprised of one or more proteins – Cry and Cyt proteins. When Btk bacteria is ingested by these organisms, these proteins adversely affects their digestive systems, leading to their death. Due to this ability, Btk is used as biological pest control agent against lepidopterans.

In MON 810, the gene coding for the Cry1Ab protein in Btk was isolated and inserted into the genome of maize. This event allows for the maize, known as Bt maize, to produce the same Cry1Ab protein with insecticidal properties. When larvae of the typical maize pests,

African maize stalk borer and fall armyworm, eat the leaves of Bt maize, they suffer the same fate as ingesting the bacterium itself, thus protecting the maize against these pests.

4.2.2 Event MON 89034

Monsanto also developed event MON 89034 marketed under the tradename YieldGard® VT PRO. It is based on the same Btk bacterium, but expresses the proteins Cry1A.105 and Cry2Ab2, and has improved insecticidal properties over MON 810.

4.2.3 Event NK 603

Event NK 603, marketed as Roundup Ready® maize, is also a Monsanto product. It is resistant to glyphosate, a non-selective post-emergent systemic herbicide. Glyphosate (N-phosphonomethyl-glycine) is absorbed by plants and binds to the plant enzyme enolpyruvylshikimate-3-phosphate synthase (EPSPS). By binding to EPSPS, glyphosate blocks the enzyme's function in the shikimic pathway, preventing the production of aromatic amino acids and metabolites. This ultimately results in plant death by "starvation". Glyphosate is the active ingredient in the herbicide Roundup.

The bacterium, *Agrobacterium* sp. strain CP4, is a common soil bacterium that expresses a glyphosate-tolerant EPSPS enzyme. Glyphosate resistant maize is produced by inserting the CP4 EPSPS gene into maize. Glyphosate resistance is thus brought on by the EPSPS enzyme, now produced by the maize, which continues to function in the shikimic pathway. This enables the continued production of aromatic amino acids and metabolites for growth, despite the presence of glyphosate.

4.2.4 Gene Stacked Events

In addition to the single events proposed to be planted in Namibia as discussed above, combinations of these events, or gene stacked events, are also under consideration. Event MON 89034 × NK 603 for example express both insect and glyphosate resistance and was developed by inserting the genes CP4 EPSPS, Cry1A.105 and Cry2Ab2 into maize. Similarly, NK 603 × Mon 810 contains the genes for CP4 EPSPS and Cry1Ab, also providing for insect and glyphosate resistance.

4.3 GENETICALLY MODIFIED COTTON FOR AGRICULTURE IN NAMIBIA

Namibia is a net cotton (*Zea Mays*) exporter as there are no cotton ginneries in Namibia. Cotton is planted mainly as a dry-land crop, but also under irrigation where surface water (Hardap Dam) or groundwater is abundant. The main cotton cultivation areas are the maize triangle (Otavi – Grootfontein – Tsumeb), along the Kavango River (Green Schemes) and the Hardap Scheme.

The main insect pest encountered in the cultivation of cotton in Namibia is the Lepidopteran, *Helicoverpa armigera* subsp. (African bollworm). The larval stage of this moth, the caterpillar, feeds on, not only cotton, but a variety of other crops' leaves, flowers buds, pods, fruits and seeds. In cotton they bore into the seed pod (the cotton boll) where they are relatively well protected against typical pesticides. The African bollworm can result in significantly decreased cotton yields where infestations occur and increases cotton production costs as a result of increased requirements for the use of pesticides.

Weeds also compete with cotton's resources and must, similarly to maize (section 4.2), be actively managed through mechanical removal (tillage and manual labour like hoeing) or herbicide application. In contrast to maize, cotton is a dicotyledonous plant, and herbicides controlling broadleaf plants cannot be sprayed onto post-emergent cotton. Only herbicides selective for monocotyledonous plants (i.e. grasses) can be sprayed on cotton.

Existing GM cotton events for agricultural purposes are insect resistance, glyphosate herbicide resistance, as well as both insect and glyphosate resistance. Namibian farmers wish to be granted permission to cultivate GM cotton in order to reduce losses in cotton production from pests and weeds, as well as reduce costs in cultivation of cotton. The following sections discuss the specific

events for which permission is required (the traits, and biotechnology behind them, are similar to that of the maize events, and such similarities will not be repeated here).

4.3.1 Event MON 88913

Event MON 88913, marketed as Roundup Ready™ Flex™ Cotton, is also a Monsanto product. It is resistant to glyphosate and the traits and biotechnology used are similar to, for example, NK 603 maize.

4.3.2 Event MON 88913 x MON 15985

In addition to the single event MON 88913 proposed to be planted in Namibia, a combination of events, or gene stacked event, MON 88913 × MON 15985 developed by Monsanto, is also considered. It is marketed under the trade name Roundup Ready™ Flex™ Bollgard II™ Cotton. Event MON 88913 × MON 15985 expresses both insect and glyphosate resistance and was developed by inserting the genes CP4 EPSPS, Cry2Ab2 and cry1Ac into cotton. While, in terms of its insect resistance trait it is similar to MON 810 and MON 89034 maize by producing Bt proteins, it expresses cry1Ac proteins which is not present in the maize events.

4.4 ASPECTS OF CULTIVATING GM MAIZE AND COTTON

The production and environmental release of GMOs for food and feed purposes is a controversial topic. Opinions are divided on GMOs and arguments for and against it are centred on, among others, health concerns, biodiversity impacts, food security and ethics. In this section, a summary is provided on various aspects of cultivating GM maize and cotton. A major source used is a very extensive and objective review, of hundreds of studies on GM crops, summarised in the book *Genetically Engineered Crops: Experiences and Prospects*. The book was compiled by The National Academies of Sciences, Engineering, and Medicine, of the United States of America (USA), hereafter referred to as NASEM, who is tasked, among others, to provide independent, objective analysis and advice to the nation of the USA (National Academies of Sciences, Engineering, and Medicine (NASEM), 2016). Where other literature is cited, the source is referenced.

4.4.1 Genetically Modified Crop Yield

The significant increase in the global human population is increasing pressure on food security. Since the early 1800's the world population has increased from one billion to over 8 billion in 2023. By 2050 it is expected to reach 9.7 billion (United Nations, 2019). In order to ensure food security, food and feedstuff production have to become more intensive / productive in order to get better yields without increasing the amount of land cleared for agriculture. Approximately three quarters of global maize production is used as animal feed. It is a high-energy feed for livestock and is fed either unprocessed or processed as an ingredient of feed.

In terms of crop yield, one should distinguish between potential yield and actual yield. Potential yield is the maximum tonnage/ha that a crop can produce given no reducing factors (weeds, pests, diseases, etc.), no limiting factors (i.e. an abundance of water and nutrients) and optimum carbon dioxide levels, radiation, temperature, etc. The actual yield is the real tonnage/ha harvested, which typically are less than potential yield because of reducing factors, limiting factors and less than perfect conditions.

NASEM (2016) concluded that genetic engineering of crops to increase potential yield, does not seem to be more effective than selectively breeding crops for the same purpose. However, GM crops outperforms non-GM crops in terms of actual yield (Brookes 2019; Esterhuizen 2019; Pellegrino 2018). Based on 21 years of data on cultivation of insect resistant GM maize in Spain and Portugal, an increase in yield of 11.5% and more was observed. This, together with reduced expenditure on pesticides (see section 4.4.2), resulted in an average increase in farm income of €173/ha/year (N\$2,819 at current exchange rate) (Brookes 2019). In South Africa, the estimated economic gain from using biotech crops in the period 1998 to 2016 is

US\$2.3 billion while for 2016 alone it is US\$330 million (N\$33.8 billion and N\$4.8 billion respectively) (Brookes and Barfoot, 2018; ISAAA, 2017). In a meta-analysis of 21 years' worth of field data, Pellegrino et al. (2018) confirmed a 10.1% average actual yield increase in maize. Similarly, a meta-analysis by Klümper and Qaim (2014) indicated a 22% yield increase for Bt maize and cotton (as an aggregate) when compared to traditional non-GM variants. Khuda (2017) modelled the average effects of Bt cotton on short-run profits, yields and farm inputs in Pakistan in 2008/9. In his study he found that Bt cotton yields increased by 9% in comparison with traditional cotton cultivars.

4.4.2 Pesticide Use

Intensive commercial farming methods include the use of insecticides and herbicides to control unwanted (pest) species. In maize, the African maize stalk borer, fall armyworm and African bollworm can account for massive crop losses, if not controlled. These pests were initially controlled with organochlorines and later with organophosphates. Although organophosphates are considered less toxic than the organochlorines, both are still considered to be highly detrimental to the environment. Newer insecticides contain active ingredients such as pyrethroids, carbamates, neonicotinoids and ryanoids.

Reviewing various case studies, NASEM (2019) concluded that reduced volumes of insecticides are applied on Bt crops when compared to non-Bt crops. This is supported by Brookes and Barfoot (2017), Khuda (2017), Pellegrino (2018) and Brookes (2019). The latter noting that 678,000 kg less insecticide active ingredient was used in Spain alone for the period 1998 to 2018. Where Bt and non-Bt fields are near to each other, it has been shown that even non-Bt crops required less insecticides. This is due to the nearby Bt crops reducing pest population sizes. There seems to be some instances where reduction in herbicide use is noted when herbicide resistant crops are planted. However, there is not enough sound scientific evidence to support decreased (or increased) use of herbicides (NASEM 2019). Herbicide resistant crops do however make weed control easier and more effective. Some instances of increased actual yields are also associated with herbicide tolerant crops (Brookes and Barfoot 2018).

4.4.3 Comparison of Costs and Benefits

The decision to allow the cultivation of GM crops in Namibia can be influenced or informed by various aspects and criteria. One of these aspects is the economic costs and benefits of introducing GM crops. It has already been proven at a global stage that GM crops hold substantial financial benefit over conventional crops, especially when faced with extreme climatic conditions and natural disasters such as increased pests' activity. Overall, there continues to be a considerable and growing body of evidence, in peer reviewed literature, that quantifies the positive impacts of crop biotechnology, including its economic benefits. Research over the last two decades has provided overwhelming positive results in favour of GM crops when it comes to the benefits of introduction of GM crops. Graham Brookes and Peter Barfoot have tracked farm income and production impacts since 1996 when the first GM crops were introduced, and their analysis has demonstrated over time that GM crops have a financial benefit over conventional crops. Their analysis concentrated on gross farm income effects because these are a primary driver of adoption amongst farmers (both large commercial and small-scale subsistence). They also quantified the (nett) production impact of the technology, and recognised that broader economic impacts exist, such as on labour usage, household incomes, local communities and economies.

Their research has concluded that in the last 21 years, crop biotechnology has helped farmers grow more food using fewer resources by reducing the damage caused by pests and better controlling weeds. The highest yield increases have occurred in developing countries and this has contributed to a more reliable and secure food supply base in these countries. In South America, herbicide tolerant technology has helped farmers reduce tillage, shortening the time between planting and harvesting, allowing them the opportunity to grow an additional soybean crop after wheat in the same growing season.

With higher yields and less time and money spent managing pests and weeds, farmers have earned higher incomes (also because they have more time at hand to spend on other income-generating activities). This has proved to be especially valuable for farmers in developing countries where, in 2016, an average of \$5 was received for each extra dollar invested in biotech crop seeds.

The widespread use of GM crop technology is also changing agriculture's land footprint by allowing farmers to grow more without needing to use additional land. To maintain global production levels at 2016 levels, without biotech crops, would have required farmers to plant an additional 10.8 million hectares (ha) of soybeans, 8.2 million ha of maize, 2.9 million ha of cotton and 0.5 million ha of canola, an area equivalent to the combined land area of Bangladesh and Sri Lanka. To put this in perspective, this is approximately 27% of Namibia's total land area.

Because Namibia has not formally introduced GM crops into its production systems as yet, historical exact data is unavailable and one cannot calculate the exact financial costs/benefit compared to conventional crops at this stage (Namibian Agricultural Trade Forum (ATF), 2023).

While maize cultivation in Namibia has been ongoing uninterruptedly in Namibia for decades, the same is not true for cotton cultivation. Qualitatively, Namibia's agronomic crops and fodder production areas, where traditionally maize, wheat, sunflowers, groundnuts, millet/mahangu, oats and lucerne have been grown, are all suitable for cotton production. Historically, cotton was successfully grown prior to independence and for some time thereafter on the Hardap irrigation scheme as well as in the dryland production areas of the 'maize triangle', the area around Grootfontein and in the Kavango Region. In those years, dryland yields varied from 300 kg to 1.6 tons per hectare (rainfall dependent), while irrigation farmers' averages were around 5 tons per hectare, with some farmers harvesting up to 7 tons/hectare and sometimes 9 tons/hectare on very good soils. Dryland yields per hectare on an annual basis were in most cases not profitable, and a practice whereby cotton stood over to be harvested in year 2 and sometimes year 3 as well, was the only way to ensure overall profitability could be realised (Francois Wahl, Personal Communication, 2023).

In the early 2000's a fundamental shift occurred in the agronomic industry in Namibia and cotton production declined drastically as a result. Two main reasons for this decline include i) the prices of wheat and maize that increased more than twofold; and ii) synthetic fibre prices declined, which competed head-on with cotton fibre, thereby making it no longer that lucrative to produce cotton. In addition, globally, GM cotton was introduced more and more at the time, making other countries', including South Africa's cotton production, more competitive vis-à-vis conventional cotton production that was still being practiced in Namibia, and, as a result, prices in South Africa was also driven down. Namibian cotton production almost came to a complete halt as a result.

Currently in Namibia, there are more and more farmers from traditional cattle farming areas in the north, north east and east of Namibia with access to land and water for irrigation, that are diversifying into agronomic, oilseed and horticulture production – thereby expanding the areas in Namibia where land can be cultivated successfully. Fibre production, such as cotton, will also be suitable in these new environments.

As mentioned previously, cotton has been proven as an ideal small-scale cash crop in drier climates, due to its resilience under lower rainfall conditions. It can therefore be deemed as a suitable alternative cash crop in Namibia as well for small-scale and dryland farmers, based on successes achieved elsewhere - globally and in Africa. The main stumbling blocks in convincing small-scale farmers into cotton farming has traditionally been their reluctance to plant non-edible cash crops instead of food crops like mahangu and maize, the lack of a nearby markets and local ginneries, lack of economies of scale/critical mass, long transport distances, transport costs and bulkiness of the product, the labour intensive production system for hand-

picked cotton, and pest/weed control challenges as a result of insect infestation and the need to make use of manual weeding, inter alia (Namibia Agricultural Trade Forum, 2023).

From a quantitative or economic perspective, in order to provide some indication of what the financial costs/benefits could be, a commodity budget can be drawn up to show the estimated costs and incomes to be derived from specific crops. Commodity budgets have been calculated historically by several entities in South Africa for example and for different climatic and farming conditions, many of these similar to the farming conditions and practices that are used in Namibia. An excellent commodity budget tool (© 2023 - Profarmer) has been developed by the Griqualand West Cooperative in South Africa. Many farmers in South Africa and Namibia have historically been using tools such as the Profarmer© Tool to calculate the costs and benefits of farming with specific crops/cultivars in their respective farming areas. Maize and cotton production are also covered by the Profarmer© Tool and updated figures are prepared on an annual basis. An annual subscription allows users access to the Tool and users can include own data and yield/cost/income figures to allow for specific circumstances.

Maize

Maize production and input costs as well as yields and price information have been obtained from the Profarmer© Tool. The results thereof are included in Appendix B. The information has been summarised in Table 4-1 and contains cost calculations for both non-GM (current conventional maize being grown in Namibia) and GM maize (BT maize) and for both dryland and irrigation conditions. Information for yellow maize varieties is also included.

Whereas the exact figures will vary for Namibian conditions and from farm to farm, the important aspect that we are trying to highlight here is the comparison between GM maize and non-GM maize. It is clear that there are some notable differences between conventional maize and GM maize production systems. In general terms, the GM maize is expected to realize higher yields per hectare compared to the conventional maize (due to less damage from insects for example). Notably, there will also be a differentiation when it comes to the production costs. The GM cultivars provide for a lower total production cost/ha compared to conventional maize. This is mainly due to lower costs as a result of reduced pesticide/insecticide/herbicide applications and less tillage. GM maize seeds are however priced at a premium compared to conventional seeds and input costs will be higher as a result, especially under irrigation conditions where a huge investment will be made if yield expectations are to be maximised and 80,000 - 90,000 seed kernels are planted per hectare. The cost of seed will therefore be quite high; however this will be offset by the estimated higher yields, which overall would provide for a positive benefit.

Even though the figures in Table 4-1 are for South African farming systems, the net results should be more or less the same from a Namibian point of view, especially for the production cost side. Namibian production costs are overall around 20-30% higher than South Africa and these costs must be substituted into the budget tool by individual Namibian farmers with their real figures in order to get the exact comparisons. Assumptions need to be made at farm level regarding a couple of variables, such as the price of maize, expected yield, costs of inputs such as fuel, labour, fertiliser, interest rates, etc. At the moment, Namibian maize farmers are receiving higher prices for their maize than farmers in South Africa (ATF, 2019). Depending on the actual price of maize, the break-even yield/ha could be substantially influenced, which could make maize production either more, or less profitable, compared to South African conditions. All that needs to be done is to substitute the Namibian prices for inputs and for the maize harvest for those that currently apply to South African farmers. With a higher maize price, the breakeven yield for Namibian farmers would be much lower and profit margins could materialise at much lower yields. This could influence the decision on how much GM maize seeds are to be planted, which would lower input/production costs even further.

Table 4-1 Summary table: comparative production costs and incomes for GM and non-GM maize cultivars under known South African conditions (2019 figures used)

| | Dryland non-GM White Maize | Dryland Bt White Maize | Dryland non-GM Yellow Maize | Dryland Bt Yellow Maize | Irrigated non-GM White Maize | Irrigated Bt White Maize | Irrigated Non-GM Yellow Maize | Irrigated Bt Yellow Maize |
|---|----------------------------|------------------------|-----------------------------|-------------------------|------------------------------|-----------------------------|-------------------------------|---------------------------|
| Expected Yield (ton/ha)¹ | 5.5 | 5.5 | 5.5 | 5.5 | 13.0 | 13.0 | 13.0 | 13.5 ² |
| Expected Price (R/ton) | R3,420 | R3,420 | R3,450 | R3,450 | R3,420 | R3,420 | R3,450 | R3,450 |
| Gross Value (R/ha) | R18,810 | R18,810 | R18,975 | R18,975 | R44,460 | R44,460 | R44,850 | R46,575 |
| Production costs (R/ha incl. interest) | R15,594 | R15,516 ³ | R15,597 | R15,519 | R40,692 ⁴ | R41,979 | R40,700 | R42,113 |
| Breakeven yield (ton/ha) | 4.56 | 4.54 | 4.52 | 4.5 | 11.9 | 12.27 | 11.8 | 12.21 |
| Margin (R/ha) | R3,216 | R3,294 | R3,378 | R3,456 | R3,768 | R2,481 | R4,150 | R4,462 |
| Nett benefit/cost (R/ha)⁵ | (R78) | R78⁶ | (R78) | R78⁶ | R1,287 | (R1,287)⁷ | (R312) | R312⁸ |

¹ Expected yield/ha depends on a number of issues, including the specific cultivar that has been developed and released for a particular production year and peculiar on-farm conditions, both controllable (for example seeds planted/ha) and non-controllable (such as rainfall).

² Higher yield/ha according to cultivar developer specifications, compared to non-GMO varieties

³ Lowest production cost/ha for dryland conditions

⁴ Lowest production cost/ha for irrigation conditions

⁵ Compared to its GM or non-GM counterpart

⁶ Highest nett benefit for dryland conditions

⁷ Nett cost for irrigation conditions

⁸ Highest nett benefit for irrigation conditions

Cotton

In order to provide an economic comparison for purposes of this report, the production and input costs as well as yields and price information of the 2022/23 summer planting season and 2023 winter planting season have been obtained from the Profarmer© Tool and has been included in this report (Appendix C). The information has been summarised in Table 4-2 and contains cost calculations for GM Cotton for both dry-land and irrigation conditions. As mentioned earlier, there are for a couple of decades already no longer any conventional cotton grown in South Africa (the country is 100% GM in so far as cotton production is concerned); hence it is not possible and in any event pointless to obtain production figures for non-GM cotton varieties. Secondly, since Namibia is not producing any cotton currently, and historic production figures for Namibia are also not available any longer, a proper cost/benefit comparison is not possible between Namibian cotton production vis-à-vis GM cotton production. At most, an individual farmer will have to use its own production figures and variables over time in order to build a record of costs/benefits for comparison purposes.

Whereas the exact figures will vary for Namibian conditions and from farm to farm, the important aspects that we are trying to highlight here are the profitability variables for dryland and irrigated GM Cotton. Price of seed cotton per tonne as well as yields per hectare are the most critical and will determine whether cotton in general and GM cotton in particular can be grown profitably under Namibian production conditions or not. The South African averages that have been collected over many years have pointed to a scenario where cotton production is profitable with breakeven yields as per above table (in relation to a specific price obtained in the market for the product).

Should a Namibian farmer be able to obtain a higher (or lower) yield per hectare or a higher (or lower) price for his/her cotton, then obviously the profitability outcome and breakeven yield will be influenced (either positively, or negatively). Namibian farmers will also have to take into account additional transport costs as there are currently no ginneries in Namibia and most probably all cotton will need to be sold in South Africa, hence an additional input cost that needs to be factored in. There are also variances in relation to hand-picked cotton (labour component) vis-à-vis machine picked cotton (capital cost and machinery cost including fuel). All this is farmer unit/system specific and therefore has to be calculated on a case-by-case basis for each farmer.

However, despite the absence of conventional cotton production data for comparative purposes, the results obtained under South African conditions indicates that both dryland and irrigated GM cotton is profitable, with breakeven yields in 2023 and 2022 of 4.58 and 4.66 tons/ha (for irrigated cotton) and 1.24 and 1.41 tons/ha (for dryland cotton) respectively. This was achieved against a 2022 winter cotton price of R11,950/ton and a 2023 summer cotton price of R11,870/ton. A sensitivity analysis, factoring in various price and yield scenarios, is therefore important for each farmer.

Table 4-2 Summary table: production costs and incomes for GM cotton cultivars under known South African conditions (2022 and 2023 data used)

| | Dryland GM Cotton (2023 Winter) | Dryland GM Cotton (2022/23 Summer) | Irrigated GM Cotton (2023 Winter) | Irrigated GM Cotton (2022/23 Summer) |
|--|--|---|--|---|
| Expected Yield (ton/ha)⁹ | 1.5 | 1.5 | 5.5 | 5.5 |
| Expected Price (R/ton) | R11,950 | R11,870 | R11,950 | R11,870 |
| Gross Value (R/ha) | R17,925 | R17,805 | R65,725 | R65,285 |

⁹ Expected yield/ha depends on a number of issues, including the specific cultivar that has been developed and released for a particular production year and peculiar on-farm conditions, both controllable (for example seeds planted/ha) and non-controllable (such as rainfall).

| | | | | |
|---|---------------|---------------|----------------|---------------|
| Production costs (R/ha incl. interest) | R14,820 | R16,765 | R54,683 | R55,295 |
| Breakeven yield (ton/ha) | 1.24 | 1.41 | 4.58 | 4.66 |
| Breakeven price (R/ton) | R9,880 | R11,177 | R9,942 | R10,054 |
| Margin (R/ha) | R3,105 | R1,040 | R11,042 | R9,990 |

Source: Profarmer©. 2023

In general terms, as with other GM crops such as maize, wheat and soybeans, the GM cotton cultivars are expected to realize higher yields per hectare compared to conventional cotton (due to less damage from insects for example). Notably, there will also be a differentiation when it comes to the production costs. The GM cultivars provide for a lower total production cost/ha compared to conventional crops. This is mainly due to lower costs as a result of reduced pesticide/insecticide/herbicide applications, less mechanical weed control and tillage, and reduced fuel and machinery costs. GM seeds are however often priced at a premium compared to conventional seeds and input costs will be higher as a result, especially under irrigation conditions where a huge investment will be made if yield expectations are to be maximised and many seed kernels are to be planted per hectare. The cost of seed will therefore be quite high; however, this will be offset by the estimated higher yields, which overall would provide for a positive benefit.

Assumptions also need to be made at farm level regarding a couple of other variables, such as the price of cotton, expected yield, costs of inputs such as fuel, labour, machinery cost, packing material, transport, fertiliser, interest rates, etc. Since Namibia does not have a cotton gin, the assumptions regarding where the cotton is to be sold is very important. Likewise, transport differentials will need to be taken into account if the buyers are from outside Namibia. In the past, Namibian cotton was hand-picked and sold to South African Ginners in the Mpumalanga and Limpopo Provinces in South Africa. Towards the latter stages of cotton production in Namibia, a ginningery in Modder River in South Africa provided a ginning service to Namibian farmers at a fee. A contract price will therefore need to be negotiated with buyers prior to planting to ensure that accurate profitability calculations can be made in the budgeting process.

Depending on the actual price of cotton, the break-even yield/ha could be substantially influenced, which could make cotton production either more, or less profitable, compared to South African conditions. Of course, the input costs for Namibian conditions will also differ from South African conditions, hence the breakeven yield under Namibian conditions could be higher. Historically, Namibian production costs are overall more expensive than that of South Africa and these costs must be substituted into the budget tool by individual Namibian farmers with their real figures in order to get the exact comparisons. All that needs to be done is to substitute in the budget tool the prices of all inputs and expected yields with Namibian estimates/actual figures, instead of using the provided figures, which currently apply to South African farmers. Also, with a lower cotton price, the breakeven yield for Namibian farmers would be much higher and profit margins could be under pressure, requiring higher yields, and vice versa.

As mentioned elsewhere in this report, yield losses and crop devastation and related financial losses as a result of pests such as the African maize stalk borer, fall armyworm, Africa army worm and cotton boll worm have amplified the need for alternatives that could safeguard crops and yields against these devastating natural phenomena. The negative financial impact that a reduction in yield result in, coupled with the additional costs of spraying of pesticides (direct cost of pesticides as well as additional costs of manpower, fuel and mechanisation costs), all amplify the benefit that the introduction of BT maize could bring for both the small-scale and largescale farmer in Namibia. Army worm breakouts can devastate household food security in a matter of days, while the reduction in yields and additional costs of pesticide application could render largescale commercial irrigated maize non-profitable.

Other alternative GM crops that could show great potential in the traditional Namibian dryland farming areas, as well as the areas under irrigation could also be introduced. These include GM Maize, GM Wheat and GM Soybeans, with these crops having the potential to serve as excellent food and cash crops for farmers in addition to the traditional crops that are currently being produced, both for household food security and commercially. There is currently an upward demand for cotton - worldwide and in the region - and this could be the catalyst to introduce BT cotton as an alternative cash crop to farmers in Namibia. Cotton have already proven to be successfully grown in Namibia in the 1980's and 1990's and a collapse in world prices compared to alternatives such as maize and wheat, as well as synthetic fibres, was some of the reasons why farmers stopped producing cotton (Namibia Agricultural Trade Forum, 2023).

The SADC (Southern African Development Community) Industrialisation Strategy and Roadmap 2015–2063, and in particular the SADC Industrial Development Policy Framework, aims to promote industrialisation, enhance competitiveness, and deepen regional integration through structural transformation, leading to increased manufactured goods and exports. The SADC Region has prioritised the clothing and textile sector as one of nine key sectors to be supported in its industrial development ambitions, and the production of cotton in Namibia could be a catalyst for Namibia to enter into and participate in the highly-valued textiles and garment manufacturing cross-border value chain (<https://www.tralac.org/documents/resources/sadc/1281-sadc-industrial-development-policy-framework-2014/file.html>).

4.4.4 Trade and Marketing Issues

Given the rapid increase in the production of GMOs and the ever-expanding capabilities of biotechnology applied to food production, it is surprising that in sub-Saharan Africa—the poorest region in the world with the lowest agricultural productivity—very few countries cultivate GM crops. In fact, many countries have instituted outright bans on imported food containing GM products. One of the most high-profile examples was Zambia's ban on GM food imports, including famine relief shipments in the face of millions suffering from starvation, in 2002.

It seems that the main “stumbling block” that prevents the introduction of GM products into consumer markets or the cultivation of GM crops remains the “perception” that GM products are frowned upon by consumers. Their preferences may very well dictate what products will sell best at the corner shop; however, it is not based on a legal requirement or the results from scientific research. Consumers' perceived preference to consume non-GM products remain a voluntary preference and as a result it has been for decades wrongfully perceived that certain countries have “banned” food and feedstuffs containing GM products/ingredients. Countries across Africa and Asia that have been hesitant to introduce GMO crops, have cited the risk of future export losses as a rationale for rejecting GM technology. The reasoning behind this is because they believed that supermarket chains in major markets like the EU and Japan have instituted private standards to avoid GM ingredients in the products they sell (Gruère and Sengupta, 2009).

Over the years however, the perception that the EU has regulations/import bans in place against the importation of foodstuffs from outside that contains GMOs, has proofed to be a myth. Not only do some countries in the EU actively produce GM feed and foodstuffs; but they all allow the importation of GM feed and foodstuffs (even into those countries that may not have actively adopted GM technology in their agricultural production systems). In Germany for example, GM crops are not allowed to be planted, however they do allow feed and foodstuffs containing GMOs to be imported, which is then either consumed directly by the German consumer or finds its way into the agricultural value chains. In the EU, 60% of animal feed is imported (European Commission, 2015). The protein-rich soya in that feed comes overwhelmingly from countries that plant GM soybeans - Brazil, Argentina and the

US. The imported GM soybeans end up in the dairy, beef, pork, chicken industries, amongst others.

In the African context, in recent years, even countries like Zambia, who had a zero tolerance for anything “GMO-like”, have started to allow the importation of foodstuffs obtained from GMO products (such as breakfast cereals and prepared foodstuffs containing GMO ingredients) into its domestic market. In July 2019, the National Biosafety Authority (NBA) of Zambia has granted four companies new permits to import products that may contain GMOs (Zambia Reports, 2019). The permits were granted to Gatbro Distributors, Pick n Pay, Southern National Import and Export Limited and Choppies Super Stores. Permits were issued following a recommendation from the Scientific Advisory Committee of the NBA, to the Board, to issue the permits after risk assessment was conducted on the products that may contain GMOs and were found to be safe for human consumption.

According to the Namibian Agricultural Trade Forum (ATF), the importation of GM feed and foodstuffs have never been disallowed in Namibia. Almost all processed agricultural products and foodstuffs are imported from outside Namibia (mainly the EU and South Africa) and the majority of these contain GM ingredients. Namibia is also a net importer of cereals and other agronomic crops such as wheat, maize (both white and yellow), rice, soybeans, potatoes, etc. Argentina, Canada and South Africa are main suppliers to Namibia and they are mostly cultivating GM crops. In addition to foodstuffs, almost all of Namibia’s animal feeds are produced using mostly imported ingredients that contain GMOs (soybeans, cotton seed, oil cake and yellow maize for example). These animal feeds are used by our livestock industries (beef, small stock, chicken, game, dairy, pork), including those livestock sectors that have traditionally been exporting to overseas markets such as Norway and the European Union, as well as regionally to South Africa. These markets historically accepted meat and meat products from countries that either utilise GM products as animal feed or actively grow GM crops themselves. In addition, these markets also allow the use of GM-based animal feed (either imported or locally-produced) in their own meat production value chains. Any sudden or new restrictions or bans on the export of meat from Namibia to these markets (EU, Norway, South Africa for example) - should Namibia start to allow GM crops to be cultivated locally - would therefore be far-fetched and irrational, given that these countries currently allows and historically allowed meat and meat products into their own domestic markets that already historically contained and currently contains GM ingredients (either directly or indirectly in the value chain/manufacturing). The GM crops/events that Namibia intends to cultivate, are also not new, but have been on the market for many years and are well-known, so no new or additional risks are to be introduced into the meat value chains that does not already exist (if any).

The ATF also indicated that the Meat Board of Namibia has confirmed that the export status to the European Union are not negatively influenced by the fact that Namibian animal feed already contains GM ingredients. No legal basis therefore exist that could restrict Namibian meat exports to the EU as a result of GMOs in animal feed. At most, it could be a marketing issue, linked to consumer preferences in specific markets. The latter is however only a voluntary standard, which every consumer is entitled to, and similar to the issue of consumer preference for fair trade or organic-produced products for example.

4.4.5 Biodiversity

It is argued that non-target and beneficial species are also affected in Bt crop fields, resulting in overall reduced biodiversity. Various investigations indicate that Bt crop fields have either no impact on non-target species (Pellegrino 2018) or even result in higher biodiversity than non-Bt fields sprayed with insecticides (NASEM 2019; Carpenter 2011). The literature review by Pellegrino (2018) found only Hymenoptera to be affected and specifically a parasitic wasp, *Macrocentrus cingulum*. However, since the main hosts for this wasp are stalk borers, a decrease in its presence is expected if there is a decrease in stalk borers as a result of the Bt maize.

Typical insecticides sprayed on non-Bt maize are not selective and orders other than Lepidoptera may also be harmed, as well as other classes of animals. Since Bt crops target mainly Lepidoptera, increased biodiversity can realistically be expected in Bt crop fields. For example, all existing studies indicate that honey bees are not affected by Bt crops (Duan et al. 2008, Ricroch et al. 2018).

Weed diversity in glyphosate resistant crops, after spraying with glyphosate, seems to largely depend on the type of crop (NASEM 2019). In some instances weed diversity is lower and thus arthropod diversity is also lower. In general, in the United States of America, glyphosate resistant crops sprayed with glyphosate, had similar or increased weed diversity than non-GM fields.

In terms of biodiversity within different varieties of the same crop (crop diversity), limited studies are available. However, those that have been conducted do not indicate decreased genetic variety since GM crops were introduced (Carpenter 2011).

4.4.6 Bt Toxin Resistance

Organisms continuously evolve because of random mutations at genetic level and selection pressure. For example, trees, that because of a random mutation had thorny protrusions, were not preferred by herbivores for browsing (i.e. selection pressure). Since these trees had a higher chance of survival, they had a higher chance of cross-pollinating, and thus an increased chance of containing and expressing the genetics for thorny projections. In this way, the random mutation coupled with the selection pressure, resulted in the evolution of thorns for protection against herbivores. This is a very similar process to selective breeding applied by humans to produce certain traits in organisms.

Random mutations can also lead to resistance in insects against the active ingredients of insecticides (see Figure 4-3 for a schematic representation of the process). The best-known example is the resistance that developed in *Anopheles* mosquitoes to dichlorodiphenyltrichloroethane (DDT), during the fight against malaria (Fossog et al. 2013). In addition to DDT resistance, *Anopheles* mosquitoes have also developed resistance against pyrethroids and to some degree against carbamates (Wanjala et al. 2015). Insects that reproduce quickly, with large numbers of offspring, are more prone to developing resistance.

Similar to the insecticide resistance mentioned, insects can also become resistant to Bt toxins in Bt crops. When Bt crops were first approved for agriculture, the prediction by some scientists were, that insects will rapidly become resistant to Bt proteins. The reality was that although incidents of resistance in insect populations against Bt toxins have been described (van den Berg et al. 2013; van Rensburg 2007), it took much longer than initially predicted (Kunert et al. 2011).

Different strategies, which are mostly applicable to both GM crops and normal insecticide use, can delay evolution of resistance in insects. The first is by ensuring a high enough dose of the Bt toxin and / or more than one toxin is produced by the GM crop. In a population of insects, there will be individuals more susceptible to an insecticide, as well as those less susceptible. Spraying low dosages of an insecticide will only kill those more susceptible while the resistant individuals survive. A high dosage of an insecticide is more likely to kill less susceptible (resistant) individuals, thus delaying the evolution of resistance. Using multiple insecticides will also delay resistance, as it is more unlikely for an organism to be resistant to more than one insecticide. The same principle is true with GM crops. Those expressing more than one toxin and / or toxins of a higher dosage will delay evolution of resistance.

The second method used to delay resistance is to plant refuges of similar non-GM crops close to GM crop fields. A refuge of non-Bt maize will, for example, allow for the pests in question to feed and reproduce in the absence of a toxin and thus in the absence of a selection pressure. The population of insects sustained in the refuge will have a lower incidence of resistance. When these individuals mate with Bt toxin resistant individuals, it decreases the number of resistant offspring and delays the evolution of resistance.

Evidence suggest that the high dose / multiple toxins / refuge strategy can successfully delay development of resistance (NASEM 2019). The success will depend on the expression of toxins in the crop as well as appropriately sized refuges. Another factor to consider is that because Bt crops can reduce pest populations significantly, it may become feasible to plant only non-Bt crops in some years, thus further delaying the evolution of resistance.

It should be noted that resistance is possible to both traditional insecticides and Bt toxins. It is a matter of proper management and correct agricultural practices to delay the evolution of resistance. For example, planting of Bt maize and cotton should not completely negate the use of insecticides, but the two should be used together.




Figure 4-3 Schematic representation of pesticide resistance development (source: IRAC 2011)

4.4.7 Herbicide Resistance

All plants or weeds have the ability to become herbicide resistant / tolerant (Brookes and Barfoot 2018). Hundreds of weeds are herbicide resistant without the involvement of GM crops. These are listed on the International Survey of Herbicide Resistant Weeds website (<http://www.weedscience.org>) (Figure 4-4). Weeds have also evolved glyphosate resistance before the first herbicide tolerant GM crops were released. However, glyphosate resistance was also encountered where environmental release of glyphosate resistant crops occurred (NASEM 2019; Brookes and Barfoot 2018). Evolution of resistance is mostly similar to that of animals and so are the methods to delay resistance. Integrated weed management practices such as a combination of herbicides, manual hoeing or ploughing will delay evolution of resistance.




Figure 4-4 Global weed resistance (<http://www.weedscience.org/Graphs/GeoChart.aspx>)

4.4.8 Gene Flow

Concerns about the potential transfer of the modified gene sequences from a GM crop to closely related species or weeds through horizontal gene transfer (HGT) exist. Horizontal gene transfer is the transfer of genetic material, between single cell and / or multicellular organisms that did not originate from a parental donor. This is in contrast to vertical gene transfer, which is the transfer of genetic material from parent to offspring during reproduction. Horizontal gene transfer is a natural process and forms an important part of evolution. For example, hundreds of genes in humans appears to have originated from bacteria and through HGT they ended up in vertebrates, and ultimately in humans, at some point during vertebrate evolution (Heilig et al. 2001). HGT is common in prokaryotes while HGT between eukaryotes are considered scarce due to numerous obstacles that have to be overcome to achieve successful HGT (Philips et al., 2022).

The concern with gene flow involving GMOs is that the genetic material inserted into a GM organism may be transferred to other organisms and have detrimental effects. Examples include the HGT of antibiotic resistance genes to pathogens (Bennett et al. 2004, Keese, 2008) and virus to virus gene transfer resulting in new diseases (Falk and Bruening 1994; Keese, 2008).

Horizontal gene transfer from a plant to other organisms is a very rare occurrence and is expected to be less frequent than normal background rates (Keese, 2008, WHO 2014, Philips et al., 2022). Furthermore, maize is categorised as low risk in terms of its probability for gene flow to occur (Viljoen and Chetty, 2011; Tsatsakis et al., 2017). Viljoen and Chetty (2011) calculated cross-pollination success over distance. They found that at 45 m the chance for cross-pollination to occur is between 1.0% and 0.1%, at 145 m between 0.1% and 0.01% and at 473 m between 0.01% to 0.001%. Cross-pollination success over distance for cotton also shows a rapid decline in cross-pollination success over distance (Llewellyn et al. 2007). The percentage of seeds testing positive for Cry1A and Cry2A in conventional cotton segregated

from the Bt field by 1 m was 7.9%, at 12.6 m it was 1%, at 25.2 m it was 0.88% and at 48.6 m, 0.79%. Based on the results, Llewellyn et al. (2007) recommend a buffer between GM and conventional cotton of 20 m. Similar results were obtained by Sen et al. (2004) who indicated that as little as 8 to 9 m can provide good isolation. However, for both maize and cotton the success rate for cross-pollination is significantly influenced by external factors such as wind, topography, etc. Also for cotton, the presence of very high numbers of honey bees also increase cross-pollination at greater distances (Llewellyn et al., 2007).

Gene flow is considered to have negligible risks to humans and the environment (Keese, 2008; WHO, 2014) and no cases of adverse environmental effects as a result of HGT between GM crops and wild, related plants have been observed (NASEM, 2019) nor have any reports been made by 2022 of adverse impacts on human health or environmental safety due to HGT from GM plants (Philips et al., 2022).

4.4.9 GMOs as Food and Livestock Feed

Multiple arguments on the safety of GM food and feedstuffs and the risks they pose to humans and animals exist. A very long and detailed discussion falls outside of the scope of this assessment. However, a brief summary of various studies and literature reviews are presented below.

A general health concern is that the modified genes of a GM crop can be transferred to, and incorporated into the genome of, a consumer of a GM crop or its products. Potential adverse health effects may then result from this new genetic material. For example, the Cry1Ab fragments of Bt genes have been detected in animal organs (Mazza et al. 2005). The Bt gene as a whole was however not detected. It should be noted that with all food that is eaten, the fragments of genes can find its way into organs. It is not restricted to GM food only. Thus, should harmful effects realise because of gene fragments entering organs, it can occur with any of the food we eat. A second concern is that the specific protein that is expressed by the inserted gene(s), will be harmful when consumed and that allergens can be produced.

NASEM (2019), Vince et al. (2018) and de Vos et al. (2017) all reviewed existing literature on the health effects of GM feed on livestock. The conclusion reached by all three papers is that there is a lack of published evidence of adverse effects in livestock fed with GM feed. NASEM (2019) concluded: *“On the basis of detailed examination of comparisons of currently commercialized GE and non-GE foods in compositional analysis, acute and chronic animal-toxicity tests, long-term data on health of livestock fed GE foods, and human epidemiological data, the committee found no differences that implicate a higher risk to human health from GE foods than from their non-GE counterparts.”*

The conclusion makes sense since proteins, natural and GM, undergo the same process of denaturation into peptides (segments of amino acids) during the digestion process. Once denatured into amino acids, the characteristics of the original protein are no longer present.

Health impacts of glyphosate sprayed maize are also questioned. A significant contributor to people being sceptic about the health effects of eating glyphosate resistant maize stems from a 2012 study (Séralini et al. 2012). It presented data indicating that the long-term toxicity of glyphosate (specifically in Roundup®) and maize event NK603 (Roundup Ready®) on rats have severe health impacts. This resulted in large public outcry. However, the study was in the meantime retracted due to a lack of scientific accuracy, after the validity of the data was questioned and re-examined. Steinberg et. al., (2019) repeated a similar study and found that after two years of feeding rats NK603 maize, both treated with Roundup and untreated, no adverse health effects could be discerned.

Whereas no evidence of adverse health effects could be found, instances of health benefits are documented. Pellegrino et al. (2018) analysed long-term data on GM maize and stated that lower concentrations of mycotoxins (-28.8%), fumonisin (-30.6%) and thricotecens (-36.5%) are present in maize. NASEM (2019) concluded their review as follows: *“There is*

some evidence that GE insect-resistant crops have had benefits to human health by reducing insecticide poisonings and decreasing exposure to fumonisins.”

4.5 GENETICALLY MODIFIED MAIZE AND COTTON IN SOUTH AFRICA

South Africa's GM maize and cotton cultivation is an example to consider in deciding the future of GM maize production in Namibia. It provides some information on the potential advantages, disadvantages and impacts. The following list summarises some of the findings in no specific order of importance:

Positive

- ◆ Twenty one years of GM maize cultivation and related studies show that actual yield of GM maize is 5.6% to 24.5% higher than their non GM counterparts (Pellegrino et. al. 2018). Yields for cotton are also higher (Morse et al., 2006).
- ◆ Higher yields and reduced pesticide use results in increased profit margins (Morse et al., 2006).
- ◆ Since the introduction and widespread cultivation of Bt maize in 1998 in South Africa, the volume of chemical insecticides used has reduced significantly (Kunert, 2011; Mwamahonje and Mrosso, 2016). The same was found for insecticide use on Bt cotton, with significantly less insecticides applied than on conventional cotton for the period 1997 to 2001 (Morse et al., 2006).
- ◆ GM maize kernels have 28.8% lower concentrations of toxic compounds naturally produced by fungi which can cause various adverse health effects in humans and livestock. Collectively these toxins are called mycotoxins, and of the mycotoxins, fumonisin is 30.6% less and thricotecens 36.5% less (Pellegrino et. al., 2018).
- ◆ Evidence point towards Bt toxins not affecting non-target organisms (Pellegrino et. al. 2018).
- ◆ The adoption of GM maize for cultivation in South Africa has led to the stabilisation in the growth rate of the wholesale maize price, thus reducing price risk (Abidoeye and Mabaya, 2014).
- ◆ Smallholder farmers value the labour-saving benefit (mostly women and children) and increased yields (mostly men) of GM maize and GM cotton (Morse et al., 2008; Gouse, 2012; Gouse et al., 2016). Greater yields provide more income which in turn is spend on education of children, more investment in agriculture, and payment of debt (Morse et al., 2008).

Negative

- ◆ Some Bt resistance was detected in the African stalk borer in the Vaalharts irrigation scheme (van Rensburg, 2007). It seems that the lack or wrong implementation of refuges as well as the planting regime (late planting of maize as well as variance in time of planting) may have contributed to the evolution of resistance (van Rensburg, 2007; Kruger et. al., 2009).
- ◆ Lack of GM seed availability and cost to smallholder farmers may hamper the adoption of GM cropping in communal areas (Gouse et al., 2016).
- ◆ Cross pollination between GM and non-GM maize can occur where fields are near to each other (see Section 4.4.8) (Viljoen and Chetty, 2011).

Whereas most cotton plantations globally are of GM nature, there is a high, albeit small, demand for organic cotton in some niche markets. To exploit this possible opportunity, organic cotton research had been tested in South Africa in the past, but yields were not profitable, and as farmers are not subsidised as in other countries to farm organically, this venture never took off. Organic cotton production requires the use of non-GM (conventional) cottonseed, and since organic production is not commercially viable in South Africa, there is also no conventional cottonseed available. Thus, no organic cotton or conventional cotton are produced in South Africa (<https://cottonsa.org.za/cotton-facts/>).

Cotton GM varieties are as a result also all deregulated in South Africa, while all cottonseed sold in South Africa contains the Bt-gene. By-products, such as cottonseed oil and cottonseed oilcake that is used for animal feeding, are therefore also effectively genetically modified products.

5 ALTERNATIVES

Table 5-1 highlights the major advantages and disadvantages of traditional non-GM maize and cotton and various strains of GM maize and cotton.

Table 5-1 Alternative maize and cotton types for cultivation

| Alternative | Advantages | Disadvantages | Preferred Option |
|---|--|--|--|
| Maize type | | | |
| Traditional non-GM maize and cotton | <ul style="list-style-type: none"> ◆ Long established crops of which the positive and negative properties are well known ◆ Cheaper seeds ◆ Seeds easily available ◆ Can keep some harvested maize seed for next planting season | <ul style="list-style-type: none"> ◆ Highly susceptible to crop damage by insects ◆ Reduced crop yields when significant pest outbreaks occur ◆ Maize is only broad leaf herbicide tolerant ◆ Cotton is only grass herbicide tolerant ◆ More labour intensive ◆ More spraying result in more fuel use and thus greenhouse gasses ◆ Increased water use due to need for dilution of insecticides | <ul style="list-style-type: none"> ◆ Cultivation of GM maize and cotton with traditional maize and cotton as refuges. Planting a combination of GM maize and cotton events, or varying GM maize and cotton events between planting seasons, will contribute to delaying the onset of insect resistance. |
| MON 810 | <ul style="list-style-type: none"> ◆ Resistant to main pests like fall armyworm and African stalk borer ◆ Increased actual yields ◆ Reduced insecticide use ◆ Less labour intensive ◆ Less greenhouse gas emissions due to reduced fuel use for spraying ◆ Reduced water use due to less need for dilution of insecticides | <ul style="list-style-type: none"> ◆ Only one BT toxin can potentially lead to more rapid insect resistance to Bt ◆ Seed is more expensive ◆ Seed is less easily obtainable ◆ Requires special knowledge and proper management to prevent potential negative impacts | |
| MON 89034 (Maize) MON 15985 (Cotton) | <ul style="list-style-type: none"> ◆ Resistant to main pests like fall armyworm and African stalk borer ◆ Two Bt toxins has high efficiency and delay insect resistance ◆ Increased actual yields ◆ Reduced insecticide use ◆ Less labour intensive ◆ Less greenhouse gas emissions due to reduced fuel use for spraying | <ul style="list-style-type: none"> ◆ Seed is more expensive ◆ Seed is less easily obtainable ◆ Requires special knowledge and proper management to prevent potential negative impacts | |

| Alternative | Advantages | Disadvantages | Preferred Option |
|--|--|--|------------------|
| | <ul style="list-style-type: none"> ◆ Reduced water use due to less need for dilution of insecticides | | |
| NK 603 (Maize) MON 88913 (Cotton) | <ul style="list-style-type: none"> ◆ Easier weed control ◆ Increased actual yields | <ul style="list-style-type: none"> ◆ Weeds can become resistant to glyphosate ◆ Requires special knowledge and proper management to prevent potential negative impacts | |
| Stacked events | <ul style="list-style-type: none"> ◆ Both insect resistance and easier weed control ◆ Increased actual yields ◆ Reduced insecticide use ◆ Less labour intensive ◆ Less greenhouse gas emissions due to reduced fuel use for spraying ◆ Reduced water use due to less need for dilution of insecticides | <ul style="list-style-type: none"> ◆ Pests and weeds can become resistant to Bt proteins and glyphosate ◆ Requires special knowledge and proper management to prevent potential negative impacts | |

5.1 NO GO ALTERNATIVE

Maize and cotton production volumes on the existing cleared land for crop production will remain the same, or may even reduce in light of climate change, if the environmental release of GM maize and cotton are not allowed. Namibia will continue to rely heavily on maize imports (which also is GM maize) for most of the country's maize consumption. This results in a net cash outflow from the country. More land will need to be cleared to increase local maize and cotton production. Maize and cotton producers will remain vulnerable to pest outbreaks.

6 ADMINISTRATIVE, LEGAL AND POLICY REQUIREMENTS

The legislation and standards provided in Table 6-1 to Table 6-3 are relevant to the proposed environmental release of GM maize and cotton in Namibia.

Table 6-1 Namibian law applicable to GMOs

| Law | Key Aspects | Applicability to Environmental Release of GM Maize |
|---|---|---|
| The Namibian Constitution | <ul style="list-style-type: none"> ● Promote the welfare of people ● Incorporates a high level of environmental protection ● Incorporates international agreements as part of Namibian law | <ul style="list-style-type: none"> ● Genetically modified maize can potentially provide increased food security, thus promoting the welfare of people ● Introduction of GMOs may have environmental impacts |
| Environmental Management Act Act No. 7 of 2007, Government Notice No. 232 of 2007 | <ul style="list-style-type: none"> ● Defines the environment ● Promotes sustainable management of the environment and the use of natural resources ● Provides a process of assessment and control of activities with possible significant effects on the environment | <ul style="list-style-type: none"> ● Introduction of GMOs is a listed activity requiring environmental assessment |
| Environmental Management Regulations Government Notice No. 28-30 of 2012 | <ul style="list-style-type: none"> ● Commencement of the Environmental Management Act ● List activities that requires an environmental clearance certificate ● Provide Environmental Impact Assessment Regulations | <ul style="list-style-type: none"> ● Regulates the environmental assessment process |
| Biosafety Act Act No. 7 of 2006, Government Notice No. 223 of 2006 | <ul style="list-style-type: none"> ● Regulate activities involving the research, development, production, marketing, transport, application and other uses of genetically modified organisms and specified products derived from genetically modified organisms ● Prohibits planting of GMOs without registration ● Provides for formation of the Biosafety Council ● Government Notice No. 259 of 2018 declares certain products in Namibia as GMOs or GMO containing products. The schedule includes the maize events MON810, MON89034 and NK603 as well as stacked events of these | <ul style="list-style-type: none"> ● Main legislation dealing with the environmental release of GM maize |

| Law | Key Aspects | Applicability to Environmental Release of GM Maize |
|--|--|--|
| Biosafety Act Regulations Government Notice No. 210 | <ul style="list-style-type: none"> ● Provide regulations for obtaining permits to place GMO Food and feedstuff on the market, for contained use of GMOs, and for field trials and environmental release ● Provides for risk assessment reports and risk management plans for GMO related activities ● Provides for regulations pertaining to handling, labelling, transport, accidental release, etc. of GMOs | <ul style="list-style-type: none"> ● Regulates the environmental release of GM maize and incidental matters ● Regulates the environmental assessment process |
| Research, Science and Technology Act Act No. 23 of 2004, Government Notice No. 283 of 2004 | <ul style="list-style-type: none"> ● Provide for the promotion, co-ordination and development of research, science and technology in Namibia ● Establish the National Commission on Research, Science and Technology | <ul style="list-style-type: none"> ● Establishes the National Commission on Research, Science and Technology |
| Agronomic Industry Act Act No. 20 of 1992, Government Notice No. 107 of 1992 | <ul style="list-style-type: none"> ● Governs the prohibition, restriction and permitting on the sale, import and export of controlled products | <ul style="list-style-type: none"> ● Legislation pertaining to the agronomic industry who will cultivate GM maize |
| Fertilizers, Farm Feeds, Agricultural Remedies and Stock Remedies Act Act No. 36 of 1947, Government Notice No. 1239 of 1947 | <ul style="list-style-type: none"> ● Governs the registration, importation, sale and use of fertilizers, farm feeds, agricultural remedies and stock remedies ● Various amendments and regulations | <ul style="list-style-type: none"> ● Genetically modified maize will ultimately form part of farm feeds |
| Seed and Seed Varieties Act Act No. 23 of 2018, Government Notice No. 368 of 2018 | <ul style="list-style-type: none"> ● Provides for restrictions on the importation, production and sale of seed ● Not in force yet | <ul style="list-style-type: none"> ● Expected to control GM seed once enforced |
| Import and Export Control Act Act No. 30 of 1994, Government Gazette Notice No. 224 of 1994 | <ul style="list-style-type: none"> ● Controls imports into and exports from Namibia ● Provides for issuing of permits with respect to imports and exports | <ul style="list-style-type: none"> ● Genetically modified seed imports and potential GM maize or GM maize containing food and feed exports |
| Soil Conservation Act Act No. 76 of 1969 | <ul style="list-style-type: none"> ● Law relating to the combating and prevention of soil erosion, the conservation, improvement and manner of use of the soil and vegetation and the protection of the water sources in Namibia | <ul style="list-style-type: none"> ● Genetically modified crops allow for easier implementation of conservation tillage (reduced erosion) and less pesticide use |

| Law | Key Aspects | Applicability to Environmental Release of GM Maize |
|---|--|--|
| Public Health Act Act No. 36 of 1919 | <ul style="list-style-type: none"> ◆ Provides for the protection of health of all people | <ul style="list-style-type: none"> ◆ Potential health effects of consuming GMOs |
| Labour Act Act No 11 of 2007, Government Notice No. 236 of 2007 | <ul style="list-style-type: none"> ◆ Provides for Labour Law and the protection and safety of employees ◆ Labour Act, 1992: Regulations relating to the health and safety of employees at work (Government Notice No. 156 of 1997) | <ul style="list-style-type: none"> ◆ Application of herbicides on herbicide tolerant GM maize pose potential health impacts, but not more so than pesticide application on non-GM maize |
| National Agricultural Policy, 1995 | <ul style="list-style-type: none"> ◆ Aims to realize the national objectives of reviving and sustaining economic growth, creating employment opportunities, alleviating poverty and reducing inequalities in income ◆ Aims to maintain or increase levels of agricultural productivity | <ul style="list-style-type: none"> ◆ Genetically modified maize can potentially contribute to reaching the aims of the policy by providing increased yields |
| Namibia Food Safety Policy, 2014 | <ul style="list-style-type: none"> ◆ Aims to ensure food safety for all consumers in Namibia, and provide sufficient food safety guarantees on all food products traded nationally, or exported to other countries ◆ This policy ensures that control standards are established and adhered to as regards food production safety, food product hygiene, animal health and welfare, plant health and preventing the risk of contamination from external substances ◆ It lays down conditions for regulations on appropriate labelling for these foodstuffs and food products | <ul style="list-style-type: none"> ◆ Health concerns related to consumption of GMOs |

Table 6-2 Relevant multilateral environmental agreements

| Agreement | Key Aspects | Applicability to Environmental Release of GM Maize |
|---|--|--|
| Stockholm Declaration on the Human Environment, Stockholm 1972. | <ul style="list-style-type: none"> ◆ Recognizes the need for a common outlook and common principles to inspire and guide the people of the world in the preservation and enhancement of the human environment | <ul style="list-style-type: none"> ◆ In agreement with the Namibian Constitution regarding enhancement of the human environment ◆ Genetically modified maize can potentially provide increased food security, thus promoting the welfare of people |
| United Nations Framework Convention on Climate Change (UNFCCC) | <ul style="list-style-type: none"> ◆ The Convention recognises that developing countries should be accorded appropriate assistance to enable them to fulfil the terms of the Convention | <ul style="list-style-type: none"> ◆ Genetically modified crop production can reduce greenhouse gas emissions through the reduced need for spraying pesticides and thus less exhaust gasses from farm implements |
| Convention on Biological Diversity, Rio de Janeiro, 1992 | <ul style="list-style-type: none"> ◆ Under article 14 of The Convention, EIAs must be conducted for projects that may negatively affect biological diversity | <ul style="list-style-type: none"> ◆ Cultivation of GM crops can affect biodiversity through the reduction of pest species and onset of resistance in pests |
| Cartagena Protocol on Biosafety, 2000 | <ul style="list-style-type: none"> ◆ Adopted by the Convention on Biological Diversity in 2000 and came in force in 2003 ◆ Objective is to protect biological diversity from the potential risks posed by safe transfer, handling and use of Living Modified Organisms (LMOs)[GMOs] resulting from modern biotechnology ◆ Considers risks to human health | <ul style="list-style-type: none"> ◆ Address GMOs directly |
| International Treaty on Plant Genetic Resources for Food and Agriculture, 2001 | <ul style="list-style-type: none"> ◆ Promote conservation, exploration, collection, characterization, evaluation and documentation of plant genetic resources for food and agriculture ◆ Promote the sustainable use of plant genetic resources for food and agriculture | <ul style="list-style-type: none"> ◆ Cultivation of GM crops can potentially affect plant genetic resources |
| International Plant Protection Convention, Rome, 1951 | <ul style="list-style-type: none"> ◆ Promote controlling pests and diseases of plants and plant products and preventing their introduction and spread across national boundaries | <ul style="list-style-type: none"> ◆ Although not directly dealing with GMOs it established International Standards for Phytosanitary Measures (ISPMs) with applicability to GMOs (Table 6-3) |
| World Trade Organization (WTO) | <ul style="list-style-type: none"> ◆ Global international organization dealing with the rules of trade between nations ◆ The primary purpose of the WTO is to open trade for the benefit of all | <ul style="list-style-type: none"> ◆ The use of GMOs in the production of food and feedstuff may influence international trade |

| Agreement | Key Aspects | Applicability to Environmental Release of GM Maize |
|--|--|---|
| WTO Agreement on the Application of Sanitary and Phytosanitary Measures (SPS Agreement) | <ul style="list-style-type: none"> ◆ Applies to all sanitary and phytosanitary measures which may, directly or indirectly, affect international trade | <ul style="list-style-type: none"> ◆ The use of GMOs in the production of food and feedstuff may influence international trade |
| Table 6-3 Standards or codes of practise | | |
| Standard or Code | Key Aspects | Applicability to Environmental Release of GM Maize |
| International Standards for Phytosanitary Measures (ISPMs) | <ul style="list-style-type: none"> ◆ Compiled under the International Plant Protection Convention of 1951 ◆ Various standards related to GMOs that deals with plant pests | <ul style="list-style-type: none"> ◆ Some GM plants may present a phytosanitary risk ◆ The potential of GM maize becoming a pest / invasive |
| Food and Agriculture Organization (FAO) / World Health Organization (WHO): Codex Alimentarius | <ul style="list-style-type: none"> ◆ Provides international standards for all the principle foods, whether processed, semi-processed or raw, for distribution to the consumer ◆ Includes provisions in respect of food hygiene, food additives, residues of pesticides and veterinary drugs, contaminants, labelling and presentation, methods of analysis and sampling, and import and export inspection and certification. | <ul style="list-style-type: none"> ◆ Provides: <ul style="list-style-type: none"> ○ standards for maize (CXS 153-1985) and whole maize meal (CXS-154-1985) ○ principles for the risk analysis of foods derived from modern biotechnology (CXG 44-2003) ○ guidelines on performance criteria and validation of methods for detection, identification and quantification of specific DNA sequences and specific proteins in foods (CXG 74-2010) ○ compilation of Codex texts relevant to the labelling of foods derived from modern biotechnology (CXG 76-2011) ○ guideline for the conduct of food safety assessment of foods derived from recombinant-DNA plants (CXG 45-2003) |

7 IDENTIFIED IMPACTS

The following section provides a brief description of potential impacts (positive and negative) of cultivating GM maize and cotton and highlights the objective for each. The impacts are categorised according to economic, physical / chemical, biological and social impacts.

7.1 ECONOMIC

The financial feasibility of planting GM versus traditional crop cultivars will have to be considered for each farming unit. Ultimately, the goal of introducing GM crops is, among others, to increase profitability by increasing actual yields at times of pest outbreaks or by being able to plant crops in short planting seasons (i.e. late onset of rain in case of dry land cropping). Factors that can decrease profitability include administrative costs related to permitting, more expensive seeds, lower tonnage price for GM vs non-GM crops, additional expenses incurred to ensure GM crops remains contained and segregated from non-GM variants, and possible insurance costs to cover GM crop related events such as product spills during transport, costs for coexistence with neighbours planting non-GM crops, and resistance management. In case of incidents pertaining to GM crops (e.g. non-GM and GM crop contamination), there may be additional costs incurred, for example for decontamination, product withdrawals, compensation or legal costs.

7.1.1 Employment

Objective: To promote sustainable employment.

Planting of certain GM crops, such as Round-Up Ready maize, can lead to reduced labour requirements to perform certain tasks (e.g. manual hoeing of weeds). A lesser component of mostly seasonal and/or temporary workforce may result in the cultivation of such a variant. However, the introduction of GM cotton for cultivation in Namibia may entice more farmers to start planting cotton. To harvest cotton, many farmers will rely on seasonal and/or temporary employees to handpick the cotton, thereby increasing the seasonal and/or temporary workforce component of operations. Furthermore, diversification of farming activities by cultivating GM crops, may increase the overall sustainability of the farm and allow for the time and resources to pursue additional revenue streams. This may offset possible job losses resulting from the planting of GM crops. Many of the farming units in Namibia, have diverse agricultural production units which include agronomy, livestock farming, charcoal production and tourism.

Actions

Enhancement:

- ◆ Opportunities for additional income generating activities to be investigated in order to sustain employment.
- ◆ Employment of local and Namibians first. Where feasible, employment of the same seasonal and/or temporary workforce year on year.
- ◆ Adhere to all the requirements of the Labour Act.

Responsible Body:

- ◆ Proponent

Data Sources and Monitoring:

- ◆ Keep in good standing with Social Security Commission.
- ◆ Updated employment records and contracts on file.

7.1.2 Economic Resilience

Objective: Contribution to local and national treasury as well sustaining a stable earning potential for employees and industry.

The impact is based on the assumption that the net economic benefit of GMO cultivation (on a specific farm and in general), will exceed the net benefit of non-GMO cultivation. The assumption is required as the net economic benefit may in some instances not realise (e.g. when no significant pests are present). Should the assumption be correct, the benefit will be experienced greatly by the Proponent, where after multiplier effects will result in increased economic resilience in the regional and national agricultural sectors. Planting of the GMO crops will require less input in terms of pesticide application (including fuel and water) and labour, depending on the GM events planted. Therefore, producers will make time available for additional revenue generating activities to be considered. More successful harvests translates into a more sustainable flow of revenue per agricultural unit, resulting in an increase in the stability of revenue flow.

Cultivation of especially GM maize will reduce the risk to harvest failure and or losses. An indirect impact of the increased economic resilience will see increased planning ability for socio-economic aspects such as health and education.

Actions

Prevention:

- ◆ Prior to embarking on the cultivation of GM maize and cotton, each farmer must do feasibility calculations taking specific local conditions into consideration.
- ◆ Where feasible and possible, economic gains should be invested into the local agricultural sector and related communities.

Responsible Body:

- ◆ Proponent

Data Sources and Monitoring:

- ◆ Feasibility reports on file

7.1.3 Yield and Revenue

Objective: To increase maize and cotton yields and thus revenue generation at all levels i.e. employee, employer, supplier, processor and national treasury.

At present, there is no difference in the potential yield between conventional maize and cotton and GM variants. However, actual yields for GM variants may be higher due to decreased insect damage, especially during a heavy infestation or plague, and competition with weeds. Coupled to this is the potential for increased profit margins if reduced volumes of pesticides are used, which also mean less fuel and water consumption. GM seed are typically more expensive and crop producers will likely consider the financial benefits of GM maize and cotton vs. conventional maize and cotton in deciding which to plant. Refer to Appendix B and Appendix C for examples of cost guide figures.

Actions

Mitigation:

- ◆ Prior to embarking on the cultivation of GM maize or cotton, each farmer must do feasibility calculations taking specific local conditions into consideration.

Responsible Body:

- ◆ Proponent

Data Sources and Monitoring:

- ◆ Feasibility reports on file.

7.1.4 Meat Exports

Objective: No impact on producers of meat for export purposes.

Concerns were raised that international markets may be closed if Namibian livestock consumes GMO containing feed. The reality is that feed produced in, or imported to, Namibia have for a long time contained GM ingredients. The Meat Board of Namibia also confirmed that meat exports to the EU are not negatively influenced because of livestock consuming GMO containing feed (ATF 2019).

Actions

Prevention:

- ◆ Continue to adhere to the regulations and legislation pertaining to the agricultural industry which may impose certain restrictions on crops that may be cultivated or how crops are utilized.

Responsible Body:

- ◆ Proponent

Data Sources and Monitoring:

- ◆ Legal register

7.2 PHYSICAL / CHEMICAL

7.2.1 Pesticides in Soil and Groundwater

Objective: No or minimal impact on soil and groundwater as a result of pesticide use.

Pesticides can enter soil, and where porosity is high with shallow groundwater, can reach the water table. Unless organic farming is practiced, pesticide use will persist in both non-GM and GM cropping. As discussed earlier, evidence shows that the volumes of pesticides used are in fact lower for GM crops, especially for insecticides. In terms of herbicides, the concern is that where glyphosate resistant maize or cotton are planted, excessive volumes of glyphosate will be applied to combat weeds. Apart from the additional costs involved with excessive herbicide spraying, the regulations for herbicide use are the same, regardless of the choice of crop (GM vs non-GM). It will therefore be in the best interest of the farmer to maintain a pest management program that is sensible, with reduced potential impacts.

Actions

Prevention:

- ◆ Limit herbicide application as far as is practically possible.
- ◆ Application of glyphosate herbicide as per the prescribed concentration and application procedures.
- ◆ Prevent spray drift by applying herbicides during calm weather conditions.
- ◆ Proper training of operational personnel.

Responsible Body:

- ◆ Proponent; HSE Officer.

Data Sources and Monitoring:

- ◆ Keep record of all instances of herbicide application.

7.2.2 Soil Erosion

Objective: No or minimal soil erosion.

Globally, millions of tons of soil is lost through erosion each year. A significant portion of this is because of poor farming practices and tillage. Tilling is often employed to uproot weeds prior to planting of fields. This ensures all broad leaf and grassy weeds are removed. By planting glyphosate resistant maize and cotton the need for tillage is made redundant and conservation agriculture can be practiced since post emergent weeds among crops can be controlled. By practicing conservation tillage, there is less likelihood of soil loss due to water runoff and wind.

Actions

Prevention:

- ◆ Implement conservation tillage practises.

Responsible Body:

- ◆ Proponent

Data Sources and Monitoring:

- ◆ None

7.3 BIOLOGICAL

7.3.1 Pesticides Resistance

Objective: To delay, or ideally prevent, the onset of pesticide resistance in insects and weeds.

In GM crop fields, pesticide resistance has been reported in insects (against Bt proteins) and weeds (against glyphosate). This is however no different from pesticide resistance reported in non-GM crop fields. Over reliance on the use of glyphosate and the lack of crop and herbicide rotation by farmers, in some regions, contribute to the development of weed resistance. In order to address this problem and maintain good levels of weed control, farmers have increasingly adopted more integrated weed management strategies incorporating a mix of herbicides, other herbicide tolerant crops and cultural weed control measures. These include, using other herbicides with glyphosate rather than solely relying on glyphosate; using herbicide tolerant crops that are tolerant to other herbicides, such as glufosinate; and using cultural practices such as mulching. These add cost to the GM herbicide tolerant production systems compared to about 10–15 years ago, although relative to the current conventional alternative, the GM herbicide tolerant technology continues to offer important economic benefits.

Actions

Prevention:

- ◆ Develop and implement an insect and weed resistance management plan in collaboration with the seed supplier.
- ◆ The plan should among others include:
 - all farmers must adhere to the refuge strategy as stipulated by the GM seed supplier.
 - as part of the insect resistance management plan, intermittently apply insecticides to kill any pest insects that may have developed Bt resistant traits.
 - application of glyphosate herbicide as per the prescribed concentration (i.e. not lower or higher concentrations as this may be ineffective) and application procedures.
 - weed control prior to planting which should include herbicides of alternative active ingredients to allow killing of weeds that may have developed resistance to glyphosate.
 - weed control prior to its production of viable seeds.
 - cleaning of farm implements to prevent distribution of potential resistant weeds.
 - crop rotation.

Responsible Body:

- ◆ Proponent; HSE Officer; seed supplier

Data Sources and Monitoring:

- ◆ Insect and weed resistance management plan.
- ◆ Regular inspection of all fields to ensure early detection of extraordinary damage to crops that would indicate Bt resistance.
- ◆ If Bt resistance is expected, implement the insect resistance management plan and notify the NCRST and seed supplier.
- ◆ Inspection of all fields after application of glyphosate to ensure early detection of surviving weeds that may indicate resistance.
- ◆ If glyphosate resistance is expected, implement the weed resistance management plan and notify the NCRST and seed supplier.
- ◆ Keep record all instances of suspected insect or weed resistance. Note at least the species, date, extent and measures taken.
- ◆ Keep record of all instances of insecticide and herbicide application as a measure to combat weeds or to prevent / delay resistance in insects and weeds. Note at least the date, insecticide and/or herbicide used, concentration of active ingredients as applied, and the reason for application.

7.3.2 Biodiversity / Non-Target Species

Objective: To prevent or minimize impacts on biodiversity and non-target species.

Pesticides by nature are harmful to the environment. Since typical insecticides are not species specific, they affect many non-target species. Planting Bt crops that targets specifically Lepidopterans, reduce the need for spraying insecticides. Using less insecticides are overall more beneficial for the environment and results in increased biodiversity as compared to fields treated with traditional insecticides.

The aim with weed control is to rid the crop fields of all weeds. Therefore, whether it is achieved by spraying a broad-spectrum herbicide like glyphosate, or by using a combination of manual and chemical control, the result is the same. The only instance where non-target species will be affected by herbicide application, is where spray drift occurs. Spray drift can be prevented by applying pesticides during calm conditions.

Actions

Prevention:

- ◆ Limit pesticide application as far as is practically possible.
- ◆ Application of pesticides as per the prescribed concentration and application procedures.
- ◆ Prevent spray drift by applying pesticides during calm weather conditions.
- ◆ Proper training of operational personnel.

Responsible Body:

- ◆ Proponent

Data Sources and Monitoring:

- ◆ Keep record of all instances of insecticide and herbicide application. Note at least the date, insecticide and/or herbicide used, concentration of active ingredients as applied, and the reason for application.

7.3.3 GM Crops Becoming Invasive

Objective: No GM maize establishing outside of farmland.

Concerns have been raised regarding the possibility of GM crops establishing themselves outside of farmland with the potential of becoming invasive. After decades of planting traditional maize and cotton, no instances of this have been recorded and it is highly unlikely that the GM cultivars will be any different. Neither maize nor cotton has any closely related species occurring naturally within Namibia, thus further decreasing the possibility of them establishing and becoming invasive.

Actions

Prevention:

- ◆ Contain GM seeds and prevent spillages during transport.
- ◆ Spill clean-up plan where accidental spills occur during transport.
- ◆ Prevent theft of GM crop seeds.

Responsible Body:

- ◆ Proponent

Data Sources and Monitoring:

- ◆ Spill management plan.
- ◆ Record all spills and include maize strain, date, location and spill clean-up measures with photo records.
- ◆ Submit the spill report to the NCRST.

7.3.4 Horizontal Gene Transfer

Objective: No health impacts as a result of horizontal gene transfer and no conflict with organic or non-GMO farmers.

As discussed in this report, HGT is considered to have negligible risks to humans and the environment and no cases of adverse environmental effects as a result of HGT between GM crops and wild, related plants have been observed.

Actions

Prevention:

- ◆ Communicate the intention to plant GM variants to neighbours indicating buffer zones to neighbours who do not plant GM variants,
- ◆ Maintain a buffer zone of 800 m (or a distance as directed by the seed supplier) between GM and non-GM fields.

Responsible Body:

- ◆ Proponent

Data Sources and Monitoring:

- ◆ Seed supplier guidelines and contractual obligations of farmer.
- ◆ Keep record of any potential cross-contamination events and report to NCRST.

7.4 SOCIAL

Evaluating social aspects associated with the cultivation of GM produce, are complex and sensitive at times. Impacts range from feelings about the subject (and related future aspirations) to demographic change processes such as seasonal migration of a workforce. Community structures and belief systems are different on a local, regional and even sometimes, national scale. Therefore, the local context of every producer will have to be considered during individual assessments on a microeconomic scale. The assessment should consider the specific crop or trait, or the combination, which may be important as to determine which indicators to measure / evaluate. Aspects to be covered should include the following during their assessment: benefits to society, economically linked prosperity, health and welfare, freedom of choice, food supply, cultural heritage, safety, biodiversity and environmental services. The first two aspects also form part of the economic considerations of the his report while the latter two are included in the biophysical considerations. Of importance is to note that there is very little information or research done considering the social impact of GMOs in Namibia.

For this report the following main aspects, are broadly covered:

- ◆ Feelings and aspirations for the future,
- ◆ Social cohesion,
- ◆ Community health, and
- ◆ Cultural aspects.

If more sustainable employment realises in the agricultural sector, migration of workers to rural farming areas (limited to geographical areas which support maize and cotton production) may occur as workers search for employment. Increased migration to farming units may increase the integration of various cultural groups. Integration of culture and increased migration of labourers may increase the spread of HIV/AIDS. It is expected that possible migration to rural areas will not significantly affect the current migration trend in Namibia which has seen increased rates of urbanization.

7.4.2 Feelings and Aspiration for the Future

Objective: To achieve optimal consensus regarding the cultivation of GM crops and promote the positive aspects in terms of agricultural markets.

Concerns about the use and cultivation of GM variants, mainly maize, permeates certain communities and action groups throughout the world. In Namibia, concerns have also been raised on a national scale and include perceived threats to the Namibian beef export markets as well as community health concerns in consuming related food. Aspiration for the future are bleak and negatively perceived as it is anticipated that GM cultivation will affect the meat trade and the overall health economy of Namibia negatively. Camped in with these concerns, are those questions related to the possible economic harm of non-GMO farmers. Cross pollination organic non-GM crops and GM crops, as well as the risk of pesticide spray drift are issues which have been raised. For the latter, there is no difference in the risk between fields of non-GM and GM crops near organic fields. The potential for cross-pollination in maize and cotton decrease relatively quickly with increased distances between fields. Trials by Viljoen and Chetty (2011) on maize indicated a maximum distance of 650 m at which cross-pollination occurred under South African conditions. The use of buffers between non-GM and GM maize can prevent cross-pollination. This may however not be feasible where farms are small and near each other. It will be the responsibility of the GM maize farmer to establish the buffer zones as contractually agreed with the seed supplier.

An opposition view, concerning the cultivation of GM variants, reflects positive aspirations for the Namibian agricultural sector with increased local food production for human and animal use. Successful cultivation of GM maize and cotton is considered to increase the stability of markets through more reliable yield and harvest expectations. The aspiration focusses on increased food security in Namibia with secondary spin-offs such as improved soil conservation and reduced greenhouse gas emissions, etc. Improved security of supply is considered to affect the markets favorably. Both views towards GM cultivation are applicable on a National, regional and local scale.

Farmers will continue to have a choice between farming systems and choice of crop. Cultivation of GM maize in Namibia will remain optional, as is organic or non-GM cropping / farming. With increasing GM crop production, a positive spinoff for organic producers is the creation of a niche market, targeting a sector of the community who are willing to pay more for food perceived as healthier (organic).

Actions

Prevention / Enhancement:

- ◆ Education and dissemination of accurate, scientific, information pertaining to the cultivation of GMOs.
- ◆ Maintain a buffer zone of 800 m (or a distance as directed by the seed supplier) between GM and non-GM fields.

Responsible Body:

- ◆ Proponent
- ◆ Consultants

Data Sources and Monitoring:

- ◆ Seed supplier guidelines and contractual obligations of farmer.
- ◆ Keep record of any potential cross-contamination events and report to NCRST.

7.4.3 Social Cohesion

Objective: To achieve optimal coexistence between GMO and non-GMO cultivating farmers and consumers.

Social change processes which could affect community cohesion, mostly on a local scale, include changes in social structure of a community, conflicts and community adaptability. Criteria for measurement of the aspect are too complex for a national scale, however, well achievable for local evaluations. For example, an increased potential for conflicts between neighbouring farmers, (which cannot easily be separated from the overall effects of conventional agriculture). An increased potential conflict risk may result between neighbouring farmers, should coexistence measures not be applied properly (by either), or if fear of contamination increases. These conflicts could lead to serious community rifts, especially in small rural communities where people depend, to some extent, on each other (e. g. neighbourly help, shared machinery). Such conflicts could be amplified by a change in social structure due to negative economic effects. For example, if a non GMO farmer's fields are contaminated by GMO crops, the non GMO farmer may sustain economic losses which could affect their role in the community and related structure.

In contrast to the above, farming communities who share the same position towards the cultivation of GMO's, could be unified and have increased levels of community cohesion, corporation and collaboration. For the purposes of this report, both possibilities and related mitigation and or enhancement measures have been included.

Actions

Prevention/Enhancement:

- ◆ Education and dissemination of accurate, scientific, information pertaining to the cultivation of GMOs during community meetings.
- ◆ Communication of plans and intentions to cultivate GMO crops.
- ◆ Agreements on the specific GMO management measures such as the setting and adherence to buffer zones, contamination contingency plans (inclusive of remuneration for losses / insurance etc.).
- ◆ Agreement, prior cultivation of GMOs, on conflict remediation measures to be taken.
- ◆ Sharing, where feasible, information and challenges with local neighbours in addressing concerns prior to them becoming unresolvable.

Responsible Body:

- ◆ Proponent

Data Sources and Monitoring:

- ◆ Communication record kept on file.
- ◆ Any neighbour agreements kept on file.

7.4.4 Community Health and Welfare

Objective: To reduce environmental contamination, increase food security and livelihoods.

Although there are points of view that the consumption of GMO produce have detrimental health effects, there are many independent research articles which refutes this. The matter will not be discussed within the scope of this report as it ties in with the feelings towards GMO cultivation. Rather, community health and welfare will be considered in a broader sense, looking at aspects such as food security, labour (income) and environmental degradation.

GMO cultivation has the potential to safeguard crops against pests, thereby increasing the overall yield. Cumulatively, this could (considering GMO maize), increase the amount of food available locally, if and when such crops are plagued by pests. The overall gain would be an increase in food security which could be beneficial for the largest segment of the Namibian population. An increase in food security, affects the overall community health, especially for those living in poverty. An increase in production of GMOs might however also see a reduced availability of non-GMO produce, thereby reducing the food choices available to those who are against its cultivation and / or consumption.

Changed labour conditions may result in the cultivation of GM variants. Labour and remuneration directly affect households and related communities. Seasonal labour is considered as one of the groups which may be affected the most. Increased employment opportunities in for example the cultivation of cotton, may increase earning potential of the seasonal workforce, which are also employed during harvesting of many other vegetables such as onions, potatoes, pumpkins, table grapes, etc. Increased labour requirements could also result in a change in regional migration patterns. The opposite is true for those instances where reduced employment opportunities realise (such in the case of Round-Up ready maize). In such case, the probability of poverty/vulnerability increases. Therefore, community health could be negatively impacted.

An overall cultivation plan includes the aim to reduce the use of pesticides on crops (Bt crops) while also enabling less reliance on tillage. Both of these fundamental approaches in agriculture, contribute to overall global conservation efforts. Reducing reliance on chemical pesticides, reduces the risks of contamination though over application of pesticides, while safeguarding non-target species such as bees. Finally, the reduced use of pesticides, especially for BT maize and cotton, will reduce human contact with chemicals. There would thus be a decrease in potential medically important consequences of exposure to pesticides and chemicals.

The greatest risks related to environmental health, however still include the misuse or over application of herbicides such as Round-Up; and the build up of chemical resistance in target species. The former is not directly related to the GMO product, but rather to the individual using the product. Chemical mismanagement is not only linked to GMO producers, but can also occur on non-GM crop producing farms. Unlike non GMO producers though, GMO farmers have a strict reporting regime in efforts to kibosh chemical mismanagement and related affects. Should resistance in insects develop, for example with BT maize and cotton, an application of an alternative pesticide will be required to eliminate such resistance. It should be noted that resistance may also develop where GM crops are not involved, such as the well documented case of resistance in mosquitos to insecticides (Riveron et al. 2016).

In Namibia, conservation agriculture was identified as one of vices to combat soil degradation. Eliminating or even just reducing tillage, reduces Namibia's greenhouse gas emission rate which is linked to the reduced rate of tractor use. Since planting of glyphosate tolerant GM crops makes it easier to practise conservation tillage, it could, if done responsibly, contribute positively to Namibia's overall soil conservation and climate change strategies.

Actions

Prevention/Enhancement:

- ◆ Keep to cultivation plan of GM variants and report any resistance development according to reporting requirements.
- ◆ Identify technically and financially feasible pollution prevention and control techniques to avoid or minimize adverse impacts on human health and the environment.
- ◆ Where applicable (located close to communities), avoid or minimize the potential for community exposure to hazardous materials (chemicals) and substances that may be released through cultivation.
- ◆ Ensure that appropriate mitigation and management measures are taken to address risks and potential impacts on community health and safety arising from an influx of project-related workers (for example, ensure adequate water and sanitation is available to all seasonal employees).
- ◆ Promote the preservation of water quality, along with integrated pest management and integrated soil fertility management to minimize the use of agrochemicals and ensure that wastewater is properly treated before it is discarded.
- ◆ A pest management plan must be developed when the use of a significant volume of pesticides is foreseen.
- ◆ When required to be used to reduce probability of insect or weed resistance, hazards of pesticide must be carefully considered, and the least toxic pesticides must be selected that are: (i) known to be effective; (ii) have minimal effects on non-target species and the environment; and (iii) minimize risks and impacts associated with the development of resistance in pests.
- ◆ Measures must be taken to avoid or minimize adverse impacts on ecosystem services from project activities. Any risks or potential adverse impacts on ecosystem services that may be exacerbated by climate change, should be identified and an mitigation plan provided, (for example over abstraction of groundwater for crop cultivation).
- ◆ Provide safety and health training, including on the proper use and maintenance of machinery and personal protective equipment.
- ◆ Employ local and Namibians first.
- ◆ Where implementable, use of technologies, practices and models that generate more and better employment opportunities (both directly and indirectly) for men and women equally, including the youth.
- ◆ Adhere to all requirements of the Labour Act and the Environmental Health Act.

Responsible Body:

- ◆ Proponent

Data Sources and Monitoring:

- ◆ Pesticide use register.
- ◆ Keep all records if any resistance reporting was conducted.
- ◆ Keep records of employment.
- ◆ Keep records of health and safety training.
- ◆ Keep records of soil and water (quality sampling).

7.4.5 Cultural Aspects

Objective: Conserve and coexist with cultural tradition related to conventional and traditional crop cultivation.

In considering the preservation of cultural heritage in terms of agriculture, reference is made to the traditional practises as well as the autonomy of local populations. The former refers to traditional production techniques or the use of specific crop variants, whereas the latter refers to the freedom of the population to decide on GMO-free production or GMO-free areas. Additional heritage or archaeological resources will be subject to standard chance-find-procedures.

Maize and cotton are not crops which are traditionally grown in the rural areas of Namibia. Nonetheless, the Namibian Government has set up various projects in assisting farmers in producing crops for commercial and own use. Some of these are irrigation based projects which aim at increasing the contribution of agriculture to the country's gross domestic product and to simultaneously achieve the social development and upliftment of communities, located within areas suitable for crop farming. The bulk of maize production in Namibia is however achieved through commercial farming techniques, driven by the availability and traditional use of existing implements and seed available. The bulk of producers use conventional tillage and planting techniques of non-GMO maize, some planted as dryland crops and some under irrigation (mainly pivot irrigation systems). Recent years have seen an increase in irrigation based production, which in itself, signifies a change in the traditional methods of cultivation. Cultivation of GMO maize will both impact conventional crop production techniques (of those who plant it) as well as the traditional label of Namibia, being a GMO maize producing country.

Introduction of GMO maize and related cultivation methods, have the potential to overshadow GMO-free / organic production leading to reduced sustainability of such cultivation.

Actions

Prevention:

- ◆ Education and dissemination of accurate, scientific, information pertaining to the cultivation of GMOs.
- ◆ Should any aspect of the cultivation, utilize cultural heritage, including knowledge, innovations or practices of local communities (specifically) to benefit the project or for commercial purposes, communities should be informed of: (i) their rights under national law; (ii) the scope and nature of the proposed use; and (iii) the potential consequences.
- ◆ The public consultation process should include groups affected by the project, main users, custodians, local communities, relevant government authorities and interested NGOs.
- ◆ For archaeological resources, about the chance find procedures for the preservation of such resources.

Responsible Body:

- ◆ Proponent

Data Sources and Monitoring:

- ◆ Keep consultation record

8 PERMIT APPLICATION

The approval of EIAs for the cultivation of GMOs, and subsequent issuing of a clearances, does not automatically allow farmers to import seeds to cultivate GM maize. An application for environmental release has to be submitted to the Biosafety Council, NCRST, by each farmer. This application must among others be accompanied by the SEA and its SEMP and an emergency response plan. Standard procedures for importation of seeds continue to apply, except that more stringent regulations are in place for its transport and handling.

Typically, environmental release of a GMO for agricultural purposes is preceded by field trials. For the proposed GM maize and cotton events and their stacks, sufficient evidence is available in the form of scientific literature spanning two decades and more of GM maize and cotton cultivation in South Africa, as well as various other countries worldwide. During this period some lessons were learned, specifically for example the importance of pest management plans to prevent development of resistance. This information is now freely available. The need for field trials are therefore considered to be redundant in the Namibian context.

9 CONCLUSION

Members of the APA intends to apply for the registration of GM maize (MON 810, MON 89034, NK 603 and stacks thereof) and GM cotton (MON88913 and MON88913 × MON15985) for purposes of environmental release in Namibia. These events provide for crops with insect resistance, glyphosate resistance as well as a combination of insect and glyphosate resistance. In general terms, GMOs are ideally placed to support the Namibian economy and the Namibian Government in its endeavours to ensure food security and food self-sufficiency. With less and less resources available due to climate change, more frequent droughts and outbreaks of pests and diseases, the negative effect of chemicals and pesticides on the Namibian fauna and flora, it is more than opportune to introduce GM crops for cultivation into Namibia. Such a step could turn otherwise marginal agronomic areas into profitable production areas and assist in the alleviation of hunger and poverty for those small-scale farmers that produce for household food security.

A large part of the population objects to the idea of genetic engineering and the consumption of GM foods. While some of the objections are based on moral and ethical beliefs, other objections stem from being misinformed or being selective in the sourcing of literature to support anti-GMO campaigns. Some objections, however, do warrant caution as is the concern about development of resistance in pests. Resistance in pests is however not restricted to GM crops, but results from poor pest management practises in both non-GM and GM crop cultivation.

In a country like Namibia, with mostly marginal agronomic potential, and likely to be significantly affected by climate change, it makes sense to diversify agronomic practices by introduction GM crops into the system. Based on extensive literature reviews as touched on in this report, there is no concrete evidence that GM maize and cotton's negative impacts are such that it should not be allowed for environmental release. That being said, it remains important for farmers to be obligated to follow the regulations and recommendations prescribed for each specific GM event. This includes the management plan prepared as part of the environmental impact assessment. Furthermore, GMOs pose very little threat to organic initiatives, as these can co-exist in the same country, as already proven in many other countries that have adopted both organic and GM production systems. The US for example has the biggest organic market in the world and it is growing at an impressive rate, despite the US also being one of the biggest producers and exporters of GM crops (FiBL & IFOAM – Organics International, 2018).

10 REFERENCES

- Abdallah N. 2014. The story behind Bt cotton: Where does Sudan stand? *GM Crops & Food* 5 (4): 241-243.
- Abidoye B, Mabaya E. 2014. Adoption of genetically modified crops in South Africa: Effects on wholesale maize prices, *Agrekon: Agricultural Economics Research, Policy and Practice in Southern Africa*. 53(1): 104-123.
- Bennett P, Livesey C, Nathwani D, Reeves D, Saunders J and Wise R. 2004. An assessment of the risks associated with the use of antibiotic resistance genes in genetically modified plants: report of the Working Party of the British Society for Antimicrobial Chemotherapy. *Journal of Antimicrobial Chemotherapy* 53:418–431.
- Brookes G and Barfoot P. 2014. Economic Impact of GM Crops. *GM Crops & Food* 5(1): 65-75.
- Brookes G and Barfoot P. 2017. Environmental impacts of genetically modified (GM) crop use 1996–2015: Impacts on pesticide use and carbon emissions, *GM Crops & Food*, 8:2, 117-147.
- Brookes G and Barfoot P. 2018. Farm income and production impacts of using GM crop technology 1996–2016. *GM Crops & Food* 9(2): 59-89.
- Bruening G and Lyons J. 2000. The case of the *FLAVR SAVR* tomato. *California Agriculture* 54(4): 6-7.
- Carpenter J. 2011. Review: Impacts of GE Crops on Biodiversity. *GM Crops* 2: 1-17.
- Cohen S, Chang A, Boyer H and Helling R. 1973. Construction of Biologically Functional Bacterial Plasmids *In Vitro*. *Proceedings of the National Academy of Science of the United States of America* 70 (11): 3240-3244.
- CottonSA. 2023. <https://cottonsa.org.za/cotton-facts/>
- De Vos C and Swanenburg M. 2017. Health effects of feeding genetically modified (GM) crops to livestock animals: A review. *Food and Chemical Toxicology* 117: 3-12.
- Duan J, Marvier M, Huesing J, Dively G and Huang Z. 2008. A meta-analysis of effects of Bt crops on honey bees (Hymenoptera: Apidae). *PLoS One*. 3 (1): e1415.
- Esterhuizen D, Cladwell A. 2021. Agricultural Biotechnology Annual: Republic of South Africa (Biotechnology and Other New Production Technologies). United States Department of Agriculture: Foreign Agricultural Service. Report No. SF2021-0060 https://apps.fas.usda.gov/newgainapi/api/Report/DownloadReportByFileName?fileName=Agricultural%20Biotechnology%20Annual_Pretoria_South%20Africa%20-%20Republic%20of_10-20-2021.pdf
- Esterhuizen D. 2019. Agricultural Biotechnology Annual: Biotechnology in South Africa. Gain Report. USDA Foreign Agricultural Service. https://apps.fas.usda.gov/newgainapi/api/report/downloadreportbyfilename?filename=Agricultural%20Biotechnology%20Annual_Pretoria_South%20Africa%20-%20Republic%20of_2-5-2019.pdf accessed 21 November 2019.
- European Commission Communication. 2015. <https://www.bbc.com/news/world-europe-32450268>

- Falk B, Bruening G. 1994. Will transgenic crops generate new viruses and new diseases? *Science* 263: 1395–1396.
- FiBL & IFOAM – Organics International .2018. The World of Organic Agriculture. Frick & Bonn. <https://orgprints.org/34674/1/willer-etal-2018-world-of-organic-summary.pdf>
- Fossog Tene B, Poupardin R, Costantini C, Awono-Ambene P, Wondji CS,.... 2013. Resistance to DDT in an Urban Setting: Common Mechanisms Implicated in Both M and S Forms of *Anopheles gambiae* in the City of Yaounde' Cameroon. *PLoS ONE* 8 (4): e61408.
- Wahl F. Horticultural/Agronomic Consultant & Farmer, Personal Communication
- Gastrow M, Roberts B, Reddy V and Ismail S. 2017. Public perceptions of biotechnology in South Africa. *South African Journal of Science*. 114 (1/2).
- Gouse M, Sengupta D, Zambrano P, and Zepeda J. 2016. Genetically Modified Maize: Less Drudgery for Her, More Maize for Him? Evidence from Smallholder Maize Farmers in South Africa. *World Development* 83: 27-38.
- Gouse M. 2012. GM maize as subsistence crop: the South African smallholder. *AgBioForum* 15 (2): 163-174.
- Brookes G, Barfoot P. 2014. Economic Impact of GM Crops, *GM Crops & Food*, 5:1, 65-75, DOI: 10.4161/gmcr.28098. <https://doi.org/10.4161/gmcr.28098>
- Gruère G, Debdatta S. 2009. “GM-free private standards and their effects on biosafety decision-making in developing countries.” *Food Policy* 34 (2009) 399–406. <https://www.ictsd.org/bridges-news/biores/news/gmo-trade-in-a-world-of-fragmented-consumer-preferences-and-needs>
- Heilig R, Eckenberg R, Petit J, Fonknechten N, Da Silva C, Cattolico L, ... 2004. International Human Genome Sequencing Consortium. *Nature* 431, 931-945. *Nature*. 431. 931-945. 10.1038/nature03001.
- Hofs JL, Kirsten J. Working Paper, 2001-17 (2001): Genetically Modified Cotton in South Africa: The Solution for Rural Development?, Department of Agricultural Economics, University of Pretoria, South Africa
- <https://allianceforscience.cornell.edu/blog/2019/02/egypt-poised-lead-africa-ag-biotech-innovation/> accessed 21 November 2019
- <https://faculty.sites.iastate.edu/jfw/project/cotton-fiber-and-domestication-process> accessed 25 August 2023
- <https://www.isaaa.org/gmaprovaldatabase/countrylist/default.asp> accessed 25 August 2025
- Ihsan S, Mustafa O, Yuksel B, Bekir C, Nefi K, Selahattin A. 2004. Assessing the Out-crossing Ratio, Isolation Distance and Pollinator Insects in Cotton (*Gossypium hirsutum* L.). *Asian Journal of Plant Sciences*, 3: 724-727.
- ISAAA. 2019. Brief 55: Global Status of Commercialized Biotech/GM Crops: 2019 <https://www.isaaa.org/resources/publications/briefs/55/default.asp>

- ISAAA. 1997. Global Status of Transgenic Crops in 1997. Brief 5. <https://www.isaaa.org/resources/publications/briefs/05/default.html> accessed 21 November 2019.
- ISAAA. 2018. Executive Summary: Global Status of Commercialized biotech/Gm Crops in 2018: Biotech Crops Continue to help meet the Challenges of Increased Population and Climate Change. Brief 54. <http://isaaa.org/resources/publications/briefs/54/executivesummary/pdf/B54-ExecSum-English.pdf> accessed 21 November 2019.
- ISAAA. 2019. Global Status of Commercialized Biotech/GM Crops in 2017: Biotech Crop Adoption Surges as Economic Benefits Accumulate in 22 Years. Brief 53 <http://www.isaaa.org/resources/publications/briefs/53/download/isaaa-brief-53-2017.pdf> accessed 21 November 2019.
- Jaenisch R, Mintz B. 1974. Simian Virus 40 DNA Sequences in DNA of Healthy Adult Mice Derived from Preimplantation Blastocysts Injected with Viral DNA. *Proceedings of the National Academy of Science of the United States of America* 71(4): 1250-1254.
- Keese P. 2008. Review Article: Risks from GMOs Due to Horizontal Gene Transfer. *Environmental Biosafety Research* 7(3): 129-149.
- Khuda B. 2017. Impacts of Bt cotton on profitability, productivity and farm inputs in Pakistan: use of panel models. *Environment and Development Economics*. 22. 1-19. 10.1017/S1355770X17000080.
- Kruger M, van Rensburg J, van den Berg J. 2009. Perspective on the development of stem borer resistance to Bt maize and refuge compliance at the Vaalharts irrigation scheme in South Africa. *Crop Protection* 28: 684-689.
- Kunert K. 2011. How effective and safe is Bt-maize in South Africa? *South African Journal of Science* 107 (9/10).
- Ladisch M, Kohlmann K. 1992. Recombinant Human Insulin: Review. *Biotechnology Progress* 68(6): 471 – 478.
- Llewellyn D, Tyson C, Constable G, Duggan B, Beale S, Steel P. 2007. Containment of regulated genetically modified cotton in the field. *Agriculture, Ecosystems & Environment*. 121. 419-429. 10.1016/j.agee.2006.11.019.
- Mazza R, Soave M, Morlacchini M, Piva G and Marocco A. 2005. Assessing the transfer of genetically modified DNA from feed to animal tissues. *Transgenic Research* 14: 775–784.
- Meat Board of Namibia: Dr Anja Boshoff-de Witt, Manager Meat Standards, PO Box 38, Ausspannplatz, Windhoek, Namibia: Personal communication
- Meat Board of Namibia: Mr Paul Strydom, General Manager, PO Box 38, Ausspannplatz, Windhoek, Namibia: Personal communication
- Meat Board of Namibia: Mr Paul Strydom, General Manager, PO Box 38, Ausspannplatz, Windhoek, Namibia: Personal communication
- Morse S and Bennett R. 2008. Impact of Bt cotton on farmer livelihoods in South Africa. *International Journal of Biotechnology*. 10. 10.1504/IJBT.2008.018355.

- Morse S, Bennett R, Ismael Y. 2006. Environmental Impact of Genetically Modified Cotton in South Africa. *Agriculture, Ecosystems & Environment*. 117: 277-289. 10.1016/j.agee.2006.04.009.
- Mwamahonje A, Mrosso L. 2016. Review: Prospects of Genetic Modified maize crop in Africa. *African Journal of Biotechnology* 15 (15): 571 – 579.
- Namibia Agricultural Trade Forum: Anton Faul, Senior Trade Advisor, Personal Communication
- Namibian Agronomic Board: Antoinette Venter, P.O. Box 5096, Ausspannplatz, Windhoek, Namibia: Personal communication
- Namibian Agronomic Board: Mr Gilbert Mulonda, P.O. Box 5096, Ausspannplatz, Windhoek, Namibia: Personal communication
- National Academies of Sciences, Engineering, and Medicine. 2016. *Genetically Engineered Crops: Experiences and Prospects*. Washington, DC: The National Academies Press. doi: 10.17226/23395.
- National Research Council (US) Committee on Identifying and Assessing Unintended Effects of Genetically Engineered Foods on Human Health. *Safety of Genetically Engineered Foods: Approaches to Assessing Unintended Health Effects*. Washington (DC): National Academies Press (US); 2004. 2, *Methods and Mechanisms for Genetic Manipulation of Plants, Animals, and Microorganisms*. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK215771/>
- Network, Canadian Biotechnology Action, Ann Slater, and Cathy Holtslander (2015). "Where in the world are GM crops and foods?"
- Pellegrino E, Bedini S, Nuti M and Ercoli L. 2018. Impact of genetically engineered maize on agronomic, environmental and toxicological traits: a meta-analysis of 21 years of field data. *Scientific Reports* 8: 3113.
- PG Economics, Graham Brookes & Peter Barfoot (2018): Farm income and production impacts of using GM crop technology 1996–2016, *GM Crops & Food*, 9:2, 59-89. <https://doi.org/10.1080/21645698.2018.1464866>
- Philips JG, Martin-Avila E, Robold AV. 2022. Horizontal gene transfer from genetically modified plants - Regulatory considerations. *Frontiers in Bioengineering and Biotechnology*. 10: 971402
- Profarmer© cost guide, 2019, Griqualand West Cooperative, De Villiers Street, Douglas 8730, Northern Cape, e-mail: info@profarmer.co.za telephone: 053 298 8200; <https://cguide.profarmer.co.za/>
- Profarmer© cost guide, 2022, Griqualand West Cooperative, De Villiers Street, Douglas 8730, Northern Cape, e-mail: info@profarmer.co.za telephone: 053 298 8200; <https://cguide.profarmer.co.za/>
- SADC Industrial Development Policy Framework, 2018. <https://www.tralac.org/documents/resources/sadc/1281-sadc-industrial-development-policy-framework-2014/file.html>.
- Slater A, Holtslander C. 2015. Where in the world are GM crops and foods? Canadian Biotechnology Action Network (CBAN).

- Steinberg P, van der Voet H, Goedhart P, Kleter G, Kok E.... 2019. Lack of adverse effects in subchronic and chronic toxicity/carcinogenicity studies on the glyphosate-resistant genetically modified maize NK603 in Wistar Han RCC rats. *Archives of Toxicology* 93:1095–1139.
- Tsatsakis A, Nawaz M, Kouretas D, Balias G,... .2017. Environmental impacts of genetically modified plants: A review. *Environmental Research* 156: 818–833.
- United Nations. 2019. Department of Economic and Social Affairs, Population Division. *World Population Prospects 2019: Data Booklet*. ST/ESA/SER.A/424.
- Van Rensburg J. 2007. First report of field resistance by the stem borer, *Busseola fusca* (Fuller) to Bt-transgenic maize. *South African Journal of Plant and Soil* 24 (3): 147-151.
- Viljoen C, Chetty L. 2011. A case study of GM maize gene flow in South Africa. *Environmental Sciences Europe* 23 (8).
- Wanjala C, Mbugi J, Ototo E, Gesuge M, Afrane Y, Atieli H, 2015. Pyrethroid and DDT Resistance and Organophosphate Susceptibility among *Anopheles* spp. Mosquitoes, Western Kenya. *Emerging Infectious Diseases* 21 (12): 2178-2181. <http://www.weedscience.org/Graphs/GeoChart.aspx> accessed 25 November 2019
- WHO 2014. https://www.who.int/foodsafety/areas_work/food-technology/faq-genetically-modified-food/en/ accessed 22 October 2019.
- Zambia Reports. 2019. <https://zambiareports.com/2019/07/09/nba-grants-permit-4-companies-gmo-imports/>
- Zambia Reports. 2019. <https://zambiareports.com/2019/07/09/nba-grants-permit-4-companies-gmo-imports/>

Appendix A: Newspaper Clippings

Vrydag 24 Januarie 2020

Republiekein

SPORT 3

SA pensioengeld 'weg' in SME-plundering

Catherino Sasman

'n Bedrag van N\$60 miljoen, wat aan die Nasionale Voorsorgingsraad van die Suid-Afrikaanse Munisipale Werkersunie (Samwu) behoort, is maar net een van die sogenaamde beleggings wat in die SME Bank-plundering verdwyn het.

Die *Mail & Guardian* het onlangs berig die bedrag van N\$60 miljoen se by 'n SME-rekening inbetaal onder die pensioenfonds se skriftelike toestemming volgens bedrieglike verklaarings voor die hooggeregshof afgedien, het die oormalige uitvoerende hoof van die SME Bank, die Zimbabweër mnr. Tawanda Muvumba, op 12 Oktober 2016 die besturende direkteur van JM Busha Asset Managers aangese om die N\$60 miljoen na die Namibiese bank oor te plaas. Dié bedrag is die volgende dag na die SME Bank vanuit 'n VBS Mutual Bank-rekening oorgeplaas. Op dieselfde dag het Muvumba 'n promesse in JM Busha se guns oorteken, en sodanig aanspreeklikheid vir die N\$60 miljoen vir die SME Bank aan JM Busha Asset Managers geskep.

BELEGGING' GEWAARBORG

Die *Mail & Guardian* het berig dat, om spyte van die SME Bank wat nou gelikwadeer word, mnr. Joseph Busha, die beleggingsfonds se uitvoerende hoof, volhou die N\$60 miljoen kan weer met rente verhoel word.

Busha het in 'n onderhoud met die SME Bank se lidwidda's, mrs. David Bryant en Jan McLaren, verlede jaar in Oktober aangevoer tydens deur Muvumba bedrieg. Die hoofbeantpe van die Suid-Afrikaanse pensioenfonds, mnr. Themba Mfeka, het aan die *Mail & Guardian* gesê deel van die fonds se "belegging" in die SME Bank is deur die Namibiese regering en fondsdirekteurs gewaarborg. Mfeka het verder beweer daar is regsaksie teen die SME Bank-direkteurs.

'n Anonieme bron het Mfeka se aantygings ontken.

Me. Tania Pearson, die SME Bank se voormalige regverteenwoordiger wat met die likwidasiestrasie in die hof moes help, sê in hofstukke die JM Busha-"belegging" en ander was pogings om die plundering weg te steek.

Finansiële rekords en bankstate toon minstens N\$247,6 miljoen is tussen Desember 2013 en Januarie 2017 uit die SME Bank se rekening by Standard Bank en FNB Namibia aan die voormede oortrekkers van die gesteelde geld betaald.

Dale van hierdie onwettige betalings is as fondse vir rekenaarhardeware of boukoste verdoes, en toe buitengewoon groot bedrae moeilik getraek het om te verduidelik, het die bedriegers dit "beleggings" begin noem. In Junie 2016 het die SME Bank se ekssterne ouditeur (BDO Namibia) 'n aantal transaksies bevestig en afskrifte van onder meer beleggingsooreenkomste versoek. Die goewerneur van die Bank of Namibia (BoN), mnr. Tshipu Shiluvu, het ook raad begin vermoed, maar het geen terugvoering van die SME Bank ontvang nie. Toenemende druk op die bank het Muvumba, mnr. Marwane Kotane van Mampepe Capital (waarin SME Bank glo 'n aantal beleggings gemaak het), en mnr. Avolle Ramachunga, uitvoerende hoof van die South African VBS Mutual Bank, 'n komplot laat bedink om bedrae elders te kry om terug in die SME Bank se rekening te betaal.

Sê is onder meer die Suid-Afrikaanse Munisipale Pensioenfonds vir N\$60 miljoen gemaak. Pearson het gesê Muvumba, Kotane en Ramachunga het in November 2016 dokumente opgestel om aan "allerlei" voor te lê dat N\$57 miljoen aan die SME Bank uit 'n sogenaamde "belegging" terugbetaal is, wat die SME Bank by VBS Mutual Bank gehou het.

"In werklikheid het hulle verdere aanspreeklikheid vir die SME Bank geskep, naamlik 'n bedrag van N\$60 miljoen wat aan JM Busha Asset Managers verskuif is," het Pearson gesê.

-catherino@republiekain.com.za



Die nou verlate SME Bank. FOTO: MARC SPRENGER



Reusepeste slaan toe

Die herfskmandowurm rig gewasse grootskaalse skade aan. FOTO: IFAO

» Noordelike, sentrale gebiede geraak

Kommandowurms, wat boere se weiding en gewasse kan verwoes, het in van die streke kop uitgesteek.

» Elvira Hartingh

Die herfskmandowurm het sy opswaai in Namibië se Zambezi-, Oshana- en Oshikotostreek gemaak. Terselfdertyd het die Afrika-kommandowurm in die Oshana-, Oshikotot- en Omusati-streek uitgebroei. Die ministerie van landbou, water en bosbou het gister 'n mediaverklaring uitgereik om die uitbreking van die peste te bevestig en voeg by verskeie hektaar grond is reeds geraak.

Intussen sê kommersiële boere in die Mieliedriehoek hulle loop reeds drie weke lank onder die herfskmandowurm deur, terwyl gerugte die ronde doen dat klein groepe van die Afrika-kommandowurm ook in die Outjo- en Otavi-gebied uitgebroei het.

Volgens die ministerie is die herfskmandowurm op 6 Januarie op Sibbinda, Kongola, Njombe, Bukalo, Itomba, Nsundwa, Kashebe en Atusanga in die Zambezi-streek opgemerk. "Die gewasse is tans in verskillende groeistadiums, van die ontkieming- tot blomstadium, en is uiters kwesbaar vir peste soos die herfs- en Afrika-kommandowurm," het die ministerie se uitvoerende direkteur, mnr. Percy Misika, in die verklaring gesê.

Misika het ook bevestig streke wat posdoeders beskikbaar het, moet met



bespuitingsprogramme begin sodat die peste nie verder moet versprei nie. Intussen het mnr. Klaas Malan, 'n akkerboer uit Namibië se Mieliedriehoek, bevestig boere in dié omgewing voer drie weke lank al 'n stryd teen die herfskmandowurm.

Hy sê hulle moet voortdurend posdoeders spuit en hy doen dit snags wanneer voordelige insekte soos bye nie rondvlieg nie, sodat hy hulle nie ook doodmaak nie. Malan sê die herfskmandowurm beskuldig mieliekoppe, wat beteken boere kry swakker graadings vir mielies wat hulle wil verkoop.

Die ministerie verduidelik die herfskmandowurm, wat sy oorsprong in Amerika het, is die eerste keer in die 2016-'17-seisoen in Namibië opgemerk. Boonop het dié pes die ongewone vermoë om vinnig te versprei en is

SA se dr. Gerhard Verdoorn onlangs gewaarsku het dié twee peste kan in Namibië uitbreek ná Namibië onder erge motplae gebuk gegaan het.

Hy en sy kollegas het gesê die motte is moontlik dié van die Afrika-kommandowurm.

Verdoorn het toe gesê weerstoelende is ideaal vir die uitbroei van peste, met die warm tropiese, vogtige lug wat uit Wes-Afrika, die Kongo en Zaire oor Suider-Afrika beweeg. "Die Afrika-kommandowurm is 'n baie ernstige pes wat jou veld groot skade kan aanrig. Dit teer hoofsaaklik op weiding en sal soms ook gewasse invaas," het hy gesê.

Hy sê die Afrika-kommandowurm is endemies aan Suider-Afrika, wat beteken dit kom natuurlik hier voor, maar rein en vog veroorsaak dat dit 'n groot getalle uitbreei.

-elvira@republiekain.com.za

Reons Tours
TOER VIR 2020

| | |
|---|--|
| 20 - 22 MAART WATERBERG & ETOSHA TOER | 11 - 18 SEPTEMBER TOER VIR VOLWASSENES EN PENSIOENARISSE NA S.A. (TUINDOEF) |
| 2 - 11 MEI TOER NA THAILAND | 18 - 19 DESEMBER FAMILIE TOER NA WES-KAAP (WESKUS & KAAPSTAD) |
| 8 - 12 JULIE TOER NA SUIDE VAN NAMIBIA (AI-ANS & LUHERITZ) | 19 DESEMBER TOER VIR JONG VOLWASSENES TUSSEN 21 TOT 35 JAAR (THIN ROETE & KAAPSTAD) |
| 15 - 28 AUGUSTUS OPVOEDKUNDE TOER NA WES-KAAP (GR.6 - 9) | 9 - 14 DESEMBER BOOTHRES NA DIE PORTUGESE EILANDE (NABU & NAMIBIE) |
| 18 - 28 AUGUSTUS OPVOEDKUNDE TOER NA WES-KAAP (GR.10 - 12) | 22 - 31 DESEMBER TOER NA VICTORIA VALLE (ZIMBABWE) |

KONTAK REON VAN WYK BY 081 274 0499
OF E-POS: INFO@REONSTOURS.COM

Zambezi farmers face fall armyworm outbreak

📅 2019-02-26 👤 John Muyamba

RUNDU - The Ministry of Agriculture, Water and Forestry has confirmed there is an outbreak of fall armyworms in the Zambezi Region after it was reported on February 14 to have attacked crops in various areas – and considering the poor rain prospects this will affect the harvest immensely. According to the ministry the outbreak poses a significant threat to smallholder farmers, mainly maize farmers, and has become a threat to food security.

In the 2016/2017 cropping season approximately 50 000 hectares of maize and millet were estimated to have been damaged by these worms that adversely affected 27 000 households. The fall armyworms were spotted in Sachona, Kongola, Ngoma, Bukalo, Kasheshe and Musanga. "Following these reports the Ministry of Agriculture, Water and Forestry took assessment missions in the affected areas on 18 and 19 February which confirmed that over 100 hectares of farmland are adversely affected by the worms. Crops in these areas are at different growth stages, from vegetative to flowering stage which is highly susceptible to fall armyworms," Margaret Kalo, spokesperson for the agriculture ministry said.

Research shows that the fall armyworms prefer maize, but can also feed on more than 80 additional species of crops, including rice, sorghum, millet, sugarcane, vegetable crops and cotton.

Fall armyworms were initially detected in Central and Western Africa in early 2016 and they quickly spread across virtually all of Sub-Saharan Africa. In July 2018 it was also confirmed in India and Yemen. Because of trade and the moth's strong flying ability, it has the potential to spread further.

Farmers will need great support through integrated pest management to sustainably manage the pest in their cropping systems. The lifespan of the fall armyworm from egg to larva to moth lasts from one to three months, and it is during the larva stage that it creates the most crop damage. Research also shows the moth can fly up to 100 km per night and the female moth can lay up to a total of 1 000 eggs in her lifetime

This reporter has learned on the website of the Food and Agriculture Organization (FAO) of the United Nations that FAO have developed a mobile phone app which can aid farmers to monitor fall armyworms in their crop fields here in Africa and farmers can research it and see how it can assist them.

There are a number of ways to try to manage this pest in maize and other crops, but because it is a new pest to Africa, none of them are guaranteed to be effective and research is going on to develop more effective solutions.


However, there are some cultural and manual practices that can help reduce their effectiveness like the use of intercropping, and crop rotation with non-grass species such as cassava can reduce crop damage.

Handpick and destroy egg masses and larvae, or collect and drop larvae in hot water. Killing one caterpillar prevents the appearance of more than 1500-2000 new caterpillars within less than four weeks, while using good quality seeds can increase plant vigour and potentially reduce damage, farmers are advised.

📅 2019-02-26 👤 John Muyamba

Worms engulf Namibia

[Home](#) / [Worms engulf Namibia](#)



By [Southern Times](#) -- @Jan24, 2020 -- 0 Comments

[+](#) [🗨️](#) [📷](#)

Timo Shihepo

Windhoek - After a prolonged period of drought, Namibian farmers have yet another problem to tackle in the form of fall army worms and African army worms which are currently attacking their crops.

The areas most affected by the fall army worms and African army worms outbreak are in the Zambezi, Oshana and Oshikoto regions.

"Following these reports, this confirmed the several hectares of farmland are adversely affected by the worms. The crops in these areas are at different growth stages, from germination, vegetative to flowering stages, which is highly susceptible to fall army worms and African army worms," said Ministry of Agriculture, Water and Forestry executive director, Percy Misika.

The fall army worm is a pervasive agricultural pest native to south and central America that ruthlessly worked its way across the African continent, after arriving in West Africa and hitting Nigeria in 2016.

The agricultural ministry said the worms then made their way to south of the Sahara and into Namibia in 2016/2017 cropping season.

"The fall army worm has spread quickly due to its short reproductive cycle and ability to travel long distance quickly during its adult (moth) stage," said Misika.

The fall army worm's lifespan from egg to larva to moth lasts from one to three months. It is during the larva stage that it creates the most crop damage. The agricultural ministry said that the fall army worm is difficult to control because it reproduces fast and in larger numbers and can migrate long distances, hides within growing leaves and is also resistant to several pesticides.

"Okapuka (worms) is in abundance here. I think we might need to buy seeds and replant again because they are really causing havoc," Kambwali, a farmer in Oshikoto region, told *The Southern Times*.

In an effort to help the farmers, the Ministry of Agriculture, Water and Forestry has directed the regions with pesticides to start the spraying programme with immediate effect in order to contain the pests in areas where they have been reported.

"In addition, all the crop growing regions were directed to intensify awareness campaigns through the local radio service in order to educate farmers on the identification, ecology and control of the pest. The Ministry will continue with surveillance, using pheromone traps in areas which are not yet reported to be affected by the pest to ensure timely control of the fall army worm and African worm outbreak," said Misika.

Appendix B: Cost Guide Figures for Bt Maize and non-GMO maize

Non-GM White Maize Dryland

| Cost Guide Figures | | | | | | profarmer | |
|--|------------------|--------------|--------------|-----------------------------|----------------|-----------|---------|
| Sub Crop | | Publications | Pricing Date | White Maize (GM-free) (Ton) | | | |
| Maize (White - GM-free) - Dryland | | Winter 2023 | 2023-07-15 | 5.50 | | | |
| Income | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| White Maize (GM-free) | 5.50 | Ton | 3 420.00 | R/ton | R18 810 | | |
| Gross Production Value | | | | | R18 810 | | |
| Expenses | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| Seed (White Maize GM-free) | 25 000.00 | Kernels/ha | 0.05 | R/pip | R1 250 | | |
| Fertiliser - Macro elements | 1.00 | Ha | 4 630.26 | R/ha | R4 630 | | |
| Fertiliser - Micro elements | 1.00 | Ha | 126.50 | R/ha | R127 | | |
| Fuel (Diesel) | 63.15 | L/ha | 19.51 | R/l | R1 232 | | |
| Herbicide | 1.00 | Ha | 699.19 | R/ha | R699 | | |
| Insecticide | 1.00 | Ha | 2 376.94 | R/ha | R2 377 | | |
| Fungicide | 1.00 | Ha | 1 291.34 | R/ha | R1 291 | | |
| Other Chemicals | 1.00 | Ha | 0.00 | R/ha | R0 | | |
| Aeroplane | 2.00 | Applications | 260.00 | R/ha | R520 | | |
| Insurance - Maize | 18 810.00 | Rand | 2.00 | % | R376 | | |
| Harvester Maize - Dryland | 1.00 | Ha | 675.00 | R/ha | R675 | | |
| Transport | 5.50 | Ton | 165.00 | R/ton | R908 | | |
| Mechanization - Repair and Maintenance | 1.00 | Ha | 749.78 | R/ha | R750 | | |
| Safex Hedging Cost | 5.50 | Ton | 3.00 | R/ton | R17 | | |
| Total Direct Cost | | | | | R14 851 | | |
| Interest | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| Interest | 6 188.04 | Rand | 12.00 | % | R743 | | |
| Total Production Cost | | | | | R15 594 | | |
| Margin Above Cost | | | | | R3 216 | | |
| Breakeven Yield/Ha | | | | | 4.56 | | |
| Breakeven Price/Ton | | | | | R2 835 | | |
| Sensitivity Analysis | | | | | | | |
| Crop Yield (t/ha) | R3 120 | R3 220 | R3 320 | R3 420 | R3 520 | R3 620 | R3 720 |
| 2.50 | -R7 794 | -R7 544 | -R7 294 | -R7 044 | -R6 794 | -R6 544 | -R6 294 |
| 3.50 | -R4 674 | -R4 324 | -R3 974 | -R3 624 | -R3 274 | -R2 924 | -R2 574 |
| 4.50 | -R1 554 | -R1 104 | -R654 | -R204 | R246 | R696 | R1 146 |
| 5.50 | R1 566 | R2 116 | R2 666 | R3 216 | R3 766 | R4 316 | R4 866 |
| 6.50 | R4 686 | R5 336 | R5 986 | R6 636 | R7 286 | R7 936 | R8 586 |
| 7.50 | R7 806 | R8 556 | R9 306 | R10 056 | R10 806 | R11 556 | R12 306 |
| 8.50 | R10 926 | R11 776 | R12 626 | R13 476 | R14 326 | R15 176 | R16 026 |
| Fertiliser (Macro Elements) | | | | | | | |
| N | 107.5 | | | Kg/ha | | | |
| P | 21.5 | | | Kg/ha | | | |
| K | 20.0 | | | Kg/ha | | | |
| Ca | 3.4 | | | Kg/ha | | | |
| Mg | 6.3 | | | Kg/ha | | | |
| S | 12.0 | | | Kg/ha | | | |

GM White Maize Dryland

| Cost Guide Figures | | | | | | profarmer | |
|--|------------------|--------------|----------|--------------|----------------|-------------------|---------|
| Sub Crop | | Publications | | Pricing Date | | White Maize (Ton) | |
| Maize (White - GM) - Dryland | | Winter 2023 | | 2023-07-15 | | 5.50 | |
| Income | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| White Maize | 5.50 | Ton | 3 420.00 | R/ton | R18 810 | | |
| Gross Production Value | | | | | R18 810 | | |
| Expenses | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| Seed (White maize) | 25 000.00 | Kernels/ha | 0.07 | R/ptp | R1 750 | | |
| Fertiliser - Macro elements | 1.00 | Ha | 4 630.26 | R/ha | R4 630 | | |
| Fertiliser - Micro elements | 1.00 | Ha | 126.50 | R/ha | R127 | | |
| Fuel (Diesel) | 63.15 | L/ha | 19.51 | R/l | R1 232 | | |
| Herbicide | 1.00 | Ha | 699.19 | R/ha | R699 | | |
| Insecticide | 1.00 | Ha | 1 802.59 | R/ha | R1 803 | | |
| Fungicide | 1.00 | Ha | 1 291.34 | R/ha | R1 291 | | |
| Other Chemicals | 1.00 | Ha | 0.00 | R/ha | R0 | | |
| Aeroplane | 2.00 | Applications | 260.00 | R/ha | R520 | | |
| Insurance - Maize | 18 810.00 | Rand | 2.00 | % | R376 | | |
| Harvester Maize - Dryland | 1.00 | Ha | 675.00 | R/ha | R675 | | |
| Transport | 5.50 | Ton | 165.00 | R/ton | R908 | | |
| Mechanization - Repair and Maintenance | 1.00 | Ha | 749.78 | R/ha | R750 | | |
| Safex Hedging Cost | 5.50 | Ton | 3.00 | R/ton | R17 | | |
| Total Direct Cost | | | | | R14 777 | | |
| Interest | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| Interest | 6 157.06 | Rand | 12.00 | % | R739 | | |
| Total Production Cost | | | | | R15 516 | | |
| Margin Above Cost | | | | | R3 294 | | |
| Breakeven Yield/Ha | | | | | 4.54 | | |
| Breakeven Price/Ton | | | | | R2 821 | | |
| Sensitivity Analysis | | | | | | | |
| Crop Yield (t/ha) | R3 120 | R3 220 | R3 320 | R3 420 | R3 520 | R3 620 | R3 720 |
| 2.50 | -R7 716 | -R7 466 | -R7 216 | -R6 966 | -R6 716 | -R6 466 | -R6 216 |
| 3.50 | -R4 596 | -R4 246 | -R3 896 | -R3 546 | -R3 196 | -R2 846 | -R2 496 |
| 4.50 | -R1 476 | -R1 026 | -R576 | -R126 | R324 | R774 | R1 224 |
| 5.50 | R1 644 | R2 194 | R2 744 | R3 294 | R3 844 | R4 394 | R4 944 |
| 6.50 | R4 764 | R5 414 | R6 064 | R6 714 | R7 364 | R8 014 | R8 664 |
| 7.50 | R7 884 | R8 634 | R9 384 | R10 134 | R10 884 | R11 634 | R12 384 |
| 8.50 | R11 004 | R11 854 | R12 704 | R13 554 | R14 404 | R15 254 | R16 104 |
| Fertiliser (Macro Elements) | | | | | | | |
| N | 107.5 | | | Kg/ha | | | |
| P | 21.5 | | | Kg/ha | | | |
| K | 20.0 | | | Kg/ha | | | |
| Ca | 3.4 | | | Kg/ha | | | |
| Mg | 6.3 | | | Kg/ha | | | |
| S | 12.0 | | | Kg/ha | | | |

Non-GM White Maize Irrigation

| Cost Guide Figures | | | | | | profarmer | |
|--|------------------|--------------|--------------|-----------------------------|----------------|-----------|---------|
| Sub Crop | | Publications | Pricing Date | White Maize (GM-free) (Ton) | | | |
| Maize (White - GM-free) | | Winter 2023 | 2023-07-15 | 13.00 | | | |
| Income | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| White Maize (GM-free) | 13.00 | Ton | 3 420.00 | R/ton | R44 460 | | |
| Gross Production Value | | | | | R44 460 | | |
| Expenses | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| Seed (White Maize GM-free) | 90 000.00 | Kernels/ha | 0.05 | R/pp | R4 500 | | |
| Fertiliser - Macro elements | 1.00 | Ha | 12 448.02 | R/ha | R12 448 | | |
| Fertiliser - Micro elements | 1.00 | Ha | 1 205.39 | R/ha | R1 205 | | |
| Fuel (Diesel) | 69.15 | L/ha | 19.51 | R/l | R1 349 | | |
| Herbicide | 1.00 | Ha | 909.98 | R/ha | R910 | | |
| Insecticide | 1.00 | Ha | 3 027.79 | R/ha | R3 028 | | |
| Fungicide | 1.00 | Ha | 1 291.34 | R/ha | R1 291 | | |
| Other Chemicals | 1.00 | Ha | 89.00 | R/ha | R89 | | |
| Aeroplane | 2.00 | Applications | 260.00 | R/ha | R520 | | |
| Insurance - Maize | 44 460.00 | Rand | 2.00 | % | R889 | | |
| Harvester Maize | 1.00 | Ha | 1 350.00 | R/ha | R1 350 | | |
| Transport | 13.00 | Ton | 165.00 | R/ton | R2 145 | | |
| Irrigation - Escom | 690.00 | mm/ha | 8.09 | R/mm | R5 582 | | |
| Irrigation - Water Board | 690.00 | mm/ha | 1.99 | R/mm | R1 373 | | |
| Irrigation - Scheduling | 1.00 | Ha | 114.00 | R/ha | R114 | | |
| Mechanization - Repair and Maintenance | 1.00 | Ha | 787.89 | R/ha | R788 | | |
| Pivot Cost - Repair and Maintenance | 1.00 | Ha | 1 133.00 | R/ha | R1 133 | | |
| Safex Hedging Cost | 13.00 | Ton | 3.00 | R/ton | R39 | | |
| Total Direct Cost | | | | | R38 754 | | |
| Interest | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| Interest | 16 147.49 | Rand | 12.00 | % | R1 938 | | |
| Total Production Cost | | | | | R40 692 | | |
| Margin Above Cost | | | | | R3 768 | | |
| Breakeven Yield/Ha | | | | | 11.50 | | |
| Breakeven Price/Ton | | | | | R3 130 | | |
| Sensitivity Analysis | | | | | | | |
| Crop Yield (t/ha) | R3 120 | R3 220 | R3 320 | R3 420 | R3 520 | R3 620 | R3 720 |
| 10.00 | -R9 492 | -R8 492 | -R7 492 | -R6 492 | -R5 492 | -R4 492 | -R3 492 |
| 11.00 | -R6 372 | -R5 272 | -R4 172 | -R3 072 | -R1 972 | -R872 | R228 |
| 12.00 | -R3 252 | -R2 052 | -R852 | R348 | R1 548 | R2 748 | R3 948 |
| 13.00 | -R132 | R1 168 | R2 468 | R3 768 | R5 068 | R6 368 | R7 668 |
| 14.00 | R2 988 | R4 388 | R5 788 | R7 188 | R8 588 | R9 988 | R11 388 |
| 15.00 | R6 108 | R7 608 | R9 108 | R10 608 | R12 108 | R13 608 | R15 108 |
| 16.00 | R9 228 | R10 828 | R12 428 | R14 028 | R15 628 | R17 228 | R18 828 |
| Fertiliser (Macro Elements) | | | | | | | |
| N | 286.0 | | | Kg/ha | | | |
| P | 52.0 | | | Kg/ha | | | |
| K | 80.0 | | | Kg/ha | | | |
| Ca | 10.0 | | | Kg/ha | | | |
| Mg | 10.0 | | | Kg/ha | | | |
| S | 25.0 | | | Kg/ha | | | |

GM White Maize Irrigation

| Cost Guide Figures | | | | | | profarmer | |
|--|------------------|--------------|-----------|--------------|----------------|-------------------|---------|
| Sub Crop | | Publications | | Pricing Date | | White Maize (Ton) | |
| Maize (White - GM) | | Winter 2023 | | 2023-07-15 | | 13.00 | |
| Income | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| White Maize | 13.00 | Ton | 3 420.00 | R/ton | R44 460 | | |
| Gross Production Value | | | | | R44 460 | | |
| Expenses | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| Seed (White maize) | 90 000.00 | Kernels/ha | 0.07 | R/kip | R6 300 | | |
| Fertiliser - Macro elements | 1.00 | Ha | 12 448.02 | R/ha | R12 448 | | |
| Fertiliser - Micro elements | 1.00 | Ha | 1 205.39 | R/ha | R1 205 | | |
| Fuel (Diesel) | 69.15 | L/ha | 19.51 | R/l | R1 349 | | |
| Herbicide | 1.00 | Ha | 909.98 | R/ha | R910 | | |
| Insecticide | 1.00 | Ha | 2 453.44 | R/ha | R2 453 | | |
| Fungicide | 1.00 | Ha | 1 291.34 | R/ha | R1 291 | | |
| Other Chemicals | 1.00 | Ha | 89.00 | R/ha | R89 | | |
| Aeroplane | 2.00 | Applications | 260.00 | R/ha | R520 | | |
| Insurance - Maize | 44 460.00 | Rand | 2.00 | % | R889 | | |
| Harvester Maize | 1.00 | Ha | 1 350.00 | R/ha | R1 350 | | |
| Transport | 13.00 | Ton | 165.00 | R/ton | R2 145 | | |
| Irrigation - Escom | 690.00 | mm/ha | 8.09 | R/mm | R5 582 | | |
| Irrigation - Water Board | 690.00 | mm/ha | 1.99 | R/mm | R1 373 | | |
| Irrigation - Scheduling | 1.00 | Ha | 114.00 | R/ha | R114 | | |
| Mechanization - Repair and Maintenance | 1.00 | Ha | 787.89 | R/ha | R788 | | |
| Pivot Cost - Repair and Maintenance | 1.00 | Ha | 1 133.00 | R/ha | R1 133 | | |
| Safex Hedging Cost | 13.00 | Ton | 3.00 | R/ton | R39 | | |
| Total Direct Cost | | | | | R39 980 | | |
| Interest | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| Interest | 16 658.17 | Rand | 12.00 | % | R1 999 | | |
| Total Production Cost | | | | | R41 979 | | |
| Margin Above Cost | | | | | R2 481 | | |
| Breakeven Yield/Ha | | | | | 12.27 | | |
| Breakeven Price/Ton | | | | | R3 229 | | |
| Sensitivity Analysis | | | | | | | |
| Crop Yield (t/ha) | R3 120 | R3 220 | R3 320 | R3 420 | R3 520 | R3 620 | R3 720 |
| 10.00 | -R10 779 | -R9 779 | -R8 779 | -R7 779 | -R6 779 | -R5 779 | -R4 779 |
| 11.00 | -R7 659 | -R6 559 | -R5 459 | -R4 359 | -R3 259 | -R2 159 | -R1 059 |
| 12.00 | -R4 539 | -R3 339 | -R2 139 | -R939 | R261 | R1 461 | R2 661 |
| 13.00 | -R1 419 | -R119 | R1 181 | R2 481 | R3 781 | R5 081 | R6 381 |
| 14.00 | R1 701 | R3 101 | R4 501 | R5 901 | R7 301 | R8 701 | R10 101 |
| 15.00 | R4 821 | R6 321 | R7 821 | R9 321 | R10 821 | R12 321 | R13 821 |
| 16.00 | R7 941 | R9 541 | R11 141 | R12 741 | R14 341 | R15 941 | R17 541 |
| Fertiliser (Macro Elements) | | | | | | | |
| N | 266.0 | | | Kg/ha | | | |
| P | 52.0 | | | Kg/ha | | | |
| K | 80.0 | | | Kg/ha | | | |
| Ca | 10.0 | | | Kg/ha | | | |
| Mg | 10.0 | | | Kg/ha | | | |
| S | 25.0 | | | Kg/ha | | | |

Non-GM Yellow Maize Dryland

| Cost Guide Figures | | | | | | profarmer | |
|--|------------------|--------------|--------------|---------------------------------|----------------|-----------|---------|
| Sub Crop | | Publications | Pricing Date | Maize (BT-free) - Dryland (Ton) | | | |
| Maize (Yellow - GM-free) - Dryland | | Winter 2023 | 2023-07-15 | 5.50 | | | |
| Income | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| Maize (BT-free) - Dryland | 5.50 | Ton | 3 450.00 | R/ton | R18 975 | | |
| Gross Production Value | | | | | R18 975 | | |
| Expenses | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| Seed (Maize BT-free) | 25 000.00 | Kemels/ha | 0.05 | R/kip | R1 250 | | |
| Fertiliser - Macro elements | 1.00 | Ha | 4 630.26 | R/ha | R4 630 | | |
| Fertiliser - Micro elements | 1.00 | Ha | 126.50 | R/ha | R127 | | |
| Fuel (Diesel) | 63.15 | L/ha | 19.51 | R/l | R1 232 | | |
| Herbicide | 1.00 | Ha | 699.19 | R/ha | R699 | | |
| Insecticide | 1.00 | Ha | 2 376.94 | R/ha | R2 377 | | |
| Fungicide | 1.00 | Ha | 1 291.34 | R/ha | R1 291 | | |
| Other Chemicals | 1.00 | Ha | 0.00 | R/ha | R0 | | |
| Aeroplane | 2.00 | Applications | 260.00 | R/ha | R520 | | |
| Insurance - Maize | 18 975.00 | Rand | 2.00 | % | R380 | | |
| Harvester Maize - Dryland | 1.00 | Ha | 675.00 | R/ha | R675 | | |
| Transport | 5.50 | Ton | 165.00 | R/ton | R908 | | |
| Mechanization - Repair and Maintenance | 1.00 | Ha | 749.78 | R/ha | R750 | | |
| Safex Hedging Cost | 5.50 | Ton | 3.00 | R/ton | R17 | | |
| Total Direct Cost | | | | | R14 856 | | |
| Interest | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| Interest | 6 189.42 | Rand | 12.00 | % | R743 | | |
| Total Production Cost | | | | | R15 597 | | |
| Margin Above Cost | | | | | R3 378 | | |
| Breakeven Yield/Ha | | | | | 4.52 | | |
| Breakeven Price/Ton | | | | | R2 836 | | |
| Sensitivity Analysis | | | | | | | |
| Crop Yield (t/ha) | R3 150 | R3 250 | R3 350 | R3 450 | R3 550 | R3 650 | R3 750 |
| 2.50 | -R7 722 | -R7 472 | -R7 222 | -R6 972 | -R6 722 | -R6 472 | -R6 222 |
| 3.50 | -R4 572 | -R4 222 | -R3 872 | -R3 522 | -R3 172 | -R2 822 | -R2 472 |
| 4.50 | -R1 422 | -R972 | -R522 | -R72 | R378 | R828 | R1 278 |
| 5.50 | R1 728 | R2 278 | R2 828 | R3 378 | R3 928 | R4 478 | R5 028 |
| 6.50 | R4 878 | R5 528 | R6 178 | R6 828 | R7 478 | R8 128 | R8 778 |
| 7.50 | R8 028 | R8 778 | R9 528 | R10 278 | R11 028 | R11 778 | R12 528 |
| 8.50 | R11 178 | R12 028 | R12 878 | R13 728 | R14 578 | R15 428 | R16 278 |
| Fertiliser (Macro Elements) | | | | | | | |
| N | 107.5 | | | Kg/ha | | | |
| P | 21.5 | | | Kg/ha | | | |
| K | 20.0 | | | Kg/ha | | | |
| Ca | 3.4 | | | Kg/ha | | | |
| Mg | 6.3 | | | Kg/ha | | | |
| S | 12.0 | | | Kg/ha | | | |

GM Yellow Maize Dryland

| Cost Guide Figures | | | | | | profarmer | |
|--|------------------|--------------|--------------|----------------------------|----------------|-----------|---------|
| Sub Crop | | Publications | Pricing Date | Maize (BT) - Dryland (Ton) | | | |
| Maize (Yellow - GM) - Dryland | | Winter 2023 | 2023-07-15 | 5.50 | | | |
| Income | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| Maize (BT) - Dryland | 5.50 | Ton | 3 450.00 | R/ton | R18 975 | | |
| Gross Production Value | | | | | R18 975 | | |
| Expenses | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| Seed (Maize BT) | 25 000.00 | Kemels/ha | 0.07 | R/kip | R1 750 | | |
| Fertiliser - Macro elements | 1.00 | Ha | 4 630.26 | R/ha | R4 630 | | |
| Fertiliser - Micro elements | 1.00 | Ha | 126.50 | R/ha | R127 | | |
| Fuel (Diesel) | 63.15 | L/ha | 19.51 | R/l | R1 232 | | |
| Herbicide | 1.00 | Ha | 699.19 | R/ha | R699 | | |
| Insecticide | 1.00 | Ha | 1 802.59 | R/ha | R1 803 | | |
| Fungicide | 1.00 | Ha | 1 291.34 | R/ha | R1 291 | | |
| Other Chemicals | 1.00 | Ha | 0.00 | R/ha | R0 | | |
| Aeroplane | 2.00 | Applications | 260.00 | R/ha | R520 | | |
| Insurance - Maize | 18 975.00 | Rand | 2.00 | % | R380 | | |
| Harvester Maize - Dryland | 1.00 | Ha | 675.00 | R/ha | R675 | | |
| Transport | 5.50 | Ton | 165.00 | R/ton | R908 | | |
| Mechanization - Repair and Maintenance | 1.00 | Ha | 749.78 | R/ha | R750 | | |
| Safex Hedging Cost | 5.50 | Ton | 3.00 | R/ton | R17 | | |
| Total Direct Cost | | | | | R14 780 | | |
| Interest | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| Interest | 6 158.44 | Rand | 12.00 | % | R739 | | |
| Total Production Cost | | | | | R15 519 | | |
| Margin Above Cost | | | | | R3 456 | | |
| Breakeven Yield/Ha | | | | | 4.50 | | |
| Breakeven Price/Ton | | | | | R2 822 | | |
| Sensitivity Analysis | | | | | | | |
| Crop Yield (t/ha) | R3 150 | R3 250 | R3 350 | R3 450 | R3 550 | R3 650 | R3 750 |
| 2.50 | -R7 644 | -R7 394 | -R7 144 | -R6 894 | -R6 644 | -R6 394 | -R6 144 |
| 3.50 | -R4 494 | -R4 144 | -R3 794 | -R3 444 | -R3 094 | -R2 744 | -R2 394 |
| 4.50 | -R1 344 | -R894 | -R444 | R6 | R456 | R906 | R1 356 |
| 5.50 | R1 806 | R2 356 | R2 906 | R3 456 | R4 006 | R4 556 | R5 106 |
| 6.50 | R4 956 | R5 606 | R6 256 | R6 906 | R7 556 | R8 206 | R8 856 |
| 7.50 | R8 106 | R8 856 | R9 606 | R10 356 | R11 106 | R11 856 | R12 606 |
| 8.50 | R11 256 | R12 106 | R12 956 | R13 806 | R14 656 | R15 506 | R16 356 |
| Fertiliser (Macro Elements) | | | | | | | |
| N | 107.5 | | | Kg/ha | | | |
| P | 21.5 | | | Kg/ha | | | |
| K | 20.0 | | | Kg/ha | | | |
| Ca | 3.4 | | | Kg/ha | | | |
| Mg | 6.3 | | | Kg/ha | | | |
| S | 12.0 | | | Kg/ha | | | |

Non-GM Yellow Maize Irrigation

| Cost Guide Figures | | | | | | | profarmer | |
|--|------------------|--------------|-----------|--------------|----------------|-----------------------|-----------|-------|
| Sub Crop | | Publications | | Pricing Date | | Maize (BT-free) (Ton) | | |
| Maize (Yellow - GM-free) | | Winter 2023 | | 2023-07-15 | | 13.00 | | |
| Income | | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | | |
| Maize (BT-free) | 13.00 | Ton | 3 450.00 | R/ton | R44 850 | | | |
| Gross Production Value | | | | | R44 850 | | | |
| Expenses | | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | | |
| Seed (Maize BT-free) | 90 000.00 | Kernels/ha | 0.05 | R/kip | R4 500 | | | |
| Fertiliser - Macro elements | 1.00 | Ha | 12 448.02 | R/ha | R12 448 | | | |
| Fertiliser - Micro elements | 1.00 | Ha | 1 205.39 | R/ha | R1 205 | | | |
| Fuel (Diesel) | 69.15 | L/ha | 19.51 | R/l | R1 349 | | | |
| Herbicide | 1.00 | Ha | 909.98 | R/ha | R910 | | | |
| Insecticide | 1.00 | Ha | 3 027.79 | R/ha | R3 028 | | | |
| Fungicide | 1.00 | Ha | 1 291.34 | R/ha | R1 291 | | | |
| Other Chemicals | 1.00 | Ha | 89.00 | R/ha | R89 | | | |
| Aeroplane | 2.00 | Applications | 260.00 | R/ha | R520 | | | |
| Insurance - Maize | 44 850.00 | Rand | 2.00 | % | R897 | | | |
| Harvester Maize | 1.00 | Ha | 1 350.00 | R/ha | R1 350 | | | |
| Transport | 13.00 | Ton | 165.00 | R/ton | R2 145 | | | |
| Irrigation - Escom | 690.00 | mm/ha | 8.09 | R/mm | R5 582 | | | |
| Irrigation - Water Board | 690.00 | mm/ha | 1.99 | R/mm | R1 373 | | | |
| Irrigation - Scheduling | 1.00 | Ha | 114.00 | R/ha | R114 | | | |
| Mechanization - Repair and Maintenance | 1.00 | Ha | 787.89 | R/ha | R788 | | | |
| Pivot Cost - Repair and Maintenance | 1.00 | Ha | 1 133.00 | R/ha | R1 133 | | | |
| Safex Hedging Cost | 13.00 | Ton | 3.00 | R/ton | R39 | | | |
| Total Direct Cost | | | | | R38 762 | | | |
| Interest | | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | | |
| Interest | 16 150.74 | Rand | 12.00 | % | R1 938 | | | |
| Total Production Cost | | | | | R40 700 | | | |
| Margin Above Cost | | | | | R4 150 | | | |
| Breakeven Yield/Ha | | | | | 11.80 | | | |
| Breakeven Price/Ton | | | | | R3 131 | | | |
| Sensitivity Analysis | | | | | | | | |
| Crop Yield (t/ha) | R3 150 | R3 250 | R3 350 | R3 450 | R3 550 | R3 650 | R3 750 | |
| 10.00 | -R9 200 | -R8 200 | -R7 200 | -R6 200 | -R5 200 | -R4 200 | -R3 200 | |
| 11.00 | -R6 050 | -R4 950 | -R3 850 | -R2 750 | -R1 650 | -R550 | R550 | |
| 12.00 | -R2 900 | -R1 700 | -R500 | R700 | R1 900 | R3 100 | R4 300 | |
| 13.00 | R250 | R1 550 | R2 850 | R4 150 | R5 450 | R6 750 | R8 050 | |
| 14.00 | R3 400 | R4 800 | R6 200 | R7 600 | R9 000 | R10 400 | R11 800 | |
| 15.00 | R6 550 | R8 050 | R9 550 | R11 050 | R12 550 | R14 050 | R15 550 | |
| 16.00 | R9 700 | R11 300 | R12 900 | R14 500 | R16 100 | R17 700 | R19 300 | |
| Fertiliser (Macro Elements) | | | | | | | | |
| N | 266.0 | | | | | | | Kg/ha |
| P | 52.0 | | | | | | | Kg/ha |
| K | 80.0 | | | | | | | Kg/ha |
| Ca | 10.0 | | | | | | | Kg/ha |
| Mg | 10.0 | | | | | | | Kg/ha |
| S | 25.0 | | | | | | | Kg/ha |

GM Yellow Maize Irrigation

| Cost Guide Figures | | | | | | | profarmer | |
|--|------------------|--------------|-----------|--------------|----------------|------------------|-----------|--|
| Sub Crop | | Publications | | Pricing Date | | Maize (BT) (Ton) | | |
| Maize (Yellow - GM) | | Winter 2023 | | 2023-07-15 | | 13.50 | | |
| Income | | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | | |
| Maize (BT) | 13.50 | Ton | 3 450.00 | R/ton | R46 575 | | | |
| Gross Production Value | | | | | R46 575 | | | |
| Expenses | | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | | |
| Seed (Maize BT) | 90 000.00 | Kernels/ha | 0.07 | R/pip | R6 300 | | | |
| Fertiliser - Macro elements | 1.00 | Ha | 12 449.69 | R/ha | R12 450 | | | |
| Fertiliser - Micro elements | 1.00 | Ha | 1 205.39 | R/ha | R1 205 | | | |
| Fuel (Diesel) | 69.15 | L/ha | 19.51 | R/l | R1 349 | | | |
| Herbicide | 1.00 | Ha | 909.98 | R/ha | R910 | | | |
| Insecticide | 1.00 | Ha | 2 453.44 | R/ha | R2 453 | | | |
| Fungicide | 1.00 | Ha | 1 291.34 | R/ha | R1 291 | | | |
| Other Chemicals | 1.00 | Ha | 89.00 | R/ha | R89 | | | |
| Aeroplane | 2.00 | Applications | 260.00 | R/ha | R520 | | | |
| Insurance - Maize | 46 575.00 | Rand | 2.00 | % | R932 | | | |
| Harvester Maize | 1.00 | Ha | 1 350.00 | R/ha | R1 350 | | | |
| Transport | 13.50 | Ton | 165.00 | R/ton | R2 228 | | | |
| Irrigation - Escom | 690.00 | mm/ha | 8.09 | R/mm | R5 582 | | | |
| Irrigation - Water Board | 690.00 | mm/ha | 1.99 | R/mm | R1 373 | | | |
| Irrigation - Scheduling | 1.00 | Ha | 114.00 | R/ha | R114 | | | |
| Mechanization - Repair and Maintenance | 1.00 | Ha | 787.89 | R/ha | R788 | | | |
| Pivot Cost - Repair and Maintenance | 1.00 | Ha | 1 133.00 | R/ha | R1 133 | | | |
| Safex Hedging Cost | 13.50 | Ton | 3.00 | R/ton | R41 | | | |
| Total Direct Cost | | | | | R40 108 | | | |
| Interest | | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | | |
| Interest | 16 711.50 | Rand | 12.00 | % | R2 005 | | | |
| Total Production Cost | | | | | R42 113 | | | |
| Margin Above Cost | | | | | R4 462 | | | |
| Breakeven Yield/Ha | | | | | 12.21 | | | |
| Breakeven Price/Ton | | | | | R3 119 | | | |
| Sensitivity Analysis | | | | | | | | |
| Crop Yield (t/ha) | R3 150 | R3 250 | R3 350 | R3 450 | R3 550 | R3 650 | R3 750 | |
| 10.50 | -R9 038 | -R7 988 | -R6 938 | -R5 888 | -R4 838 | -R3 788 | -R2 738 | |
| 11.50 | -R5 888 | -R4 738 | -R3 588 | -R2 438 | -R1 288 | -R138 | R1 012 | |
| 12.50 | -R2 738 | -R1 488 | -R238 | R1 012 | R2 262 | R3 512 | R4 762 | |
| 13.50 | R412 | R1 762 | R3 112 | R4 462 | R5 812 | R7 162 | R8 512 | |
| 14.50 | R3 562 | R5 012 | R6 462 | R7 912 | R9 362 | R10 812 | R12 262 | |
| 15.50 | R6 712 | R8 262 | R9 812 | R11 362 | R12 912 | R14 462 | R16 012 | |
| 16.50 | R9 862 | R11 512 | R13 162 | R14 812 | R16 462 | R18 112 | R19 762 | |
| Fertiliser (Macro Elements) | | | | | | | | |
| N | 286.1 | | | Kg/ha | | | | |
| P | 52.0 | | | Kg/ha | | | | |
| K | 80.1 | | | Kg/ha | | | | |
| Ca | 10.0 | | | Kg/ha | | | | |
| Mg | 10.0 | | | Kg/ha | | | | |
| S | 25.0 | | | Kg/ha | | | | |

Appendix C: Cost Guide Figures for GM Cotton: Dry-land vs. Irrigated

GM Cotton Dry-land – Summer

| Sub Crop | | Publications | Pricing Date | Cotton - Dryland (Ton) | | | |
|---|------------------|----------------|--------------|------------------------|----------------|---------|---------|
| Cotton - Dryland | | Summer 2022/23 | 2022-10-15 | 1.50 | | | |
| Income | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| Cotton - Dryland | 1.50 | Ton | 11 870.00 | R/ton | R17 805 | | |
| Gross Production Value | | | | | R17 805 | | |
| Expenses | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| Seed (Cotton) | 5.00 | Kg/ha | 138.07 | R/kg | R690 | | |
| Fertiliser - Macro elements | 1.00 | Ha | 2 669.25 | R/ha | R2 669 | | |
| Fertiliser - Micro elements | 1.00 | Ha | 344.45 | R/ha | R344 | | |
| Fuel (Diesel) | 69.59 | L/ha | 23.62 | R/l | R1 644 | | |
| Diesel (Picker) | 10.00 | L/ha | 23.62 | R/l | R236 | | |
| Herbicide | 1.00 | Ha | 528.00 | R/ha | R528 | | |
| Insecticide | 1.00 | Ha | 755.40 | R/ha | R755 | | |
| Fungicide | 1.00 | Ha | 0.00 | R/ha | R0 | | |
| Other Chemicals | 1.00 | Ha | 248.50 | R/ha | R249 | | |
| High Boy sprayer | 6.00 | Applications | 200.00 | R/ha | R1 200 | | |
| Insurance - Cotton | 17 805.00 | Rand | 9.00 | % | R1 602 | | |
| Cotton picker - Dryland | 1.00 | Ha | 3 750.00 | R/ha | R3 750 | | |
| Packaging - Plastic (Cotton) | 0.65 | Bales | 680.00 | R/bale | R439 | | |
| Transport Cotton | 1.50 | Ton | 310.00 | R/ton | R465 | | |
| Mechanization - Repair and Maintenance | 1.00 | Ha | 1 293.71 | R/ha | R1 294 | | |
| Marketing Cost Cotton - statutorily & Nakpo | 1.50 | Ton | 110.00 | R/ton | R165 | | |
| Total Direct Cost | | | | | R16 031 | | |
| Interest | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| Interest | 6 679.43 | Rand | 11.00 | % | R735 | | |
| Total Production Cost | | | | | R16 765 | | |
| Margin Above Cost | | | | | R1 040 | | |
| Breakeven Yield/Ha | | | | | 1.41 | | |
| Breakeven Price/Ton | | | | | R11 177 | | |
| Sensitivity Analysis | | | | | | | |
| Crop Yield (t/ha) | R11 120 | R11 370 | R11 620 | R11 870 | R12 120 | R12 370 | R12 620 |
| 0.75 | -R8 425 | -R8 238 | -R8 050 | -R7 863 | -R7 675 | -R7 488 | -R7 300 |
| 1.00 | -R5 645 | -R5 395 | -R5 145 | -R4 895 | -R4 645 | -R4 395 | -R4 145 |
| 1.25 | -R2 865 | -R2 553 | -R2 240 | -R1 928 | -R1 615 | -R1 303 | -R990 |
| 1.50 | -R85 | R290 | R665 | R1 040 | R1 415 | R1 790 | R2 165 |
| 1.75 | R2 695 | R3 132 | R3 570 | R4 007 | R4 445 | R4 882 | R5 320 |
| 2.00 | R5 475 | R5 975 | R6 475 | R6 975 | R7 475 | R7 975 | R8 475 |
| 2.25 | R8 255 | R8 817 | R9 380 | R9 942 | R10 505 | R11 067 | R11 630 |
| Fertiliser (Macro Elements) | | | | | | | |
| N | 31.0 | | Kg/ha | | | | |
| P | 12.0 | | Kg/ha | | | | |
| K | 16.0 | | Kg/ha | | | | |
| Ca | 0.0 | | Kg/ha | | | | |
| Mg | 0.0 | | Kg/ha | | | | |
| S | 2.3 | | Kg/ha | | | | |

GM Cotton Dry-land -Winter

| Sub Crop | | Publications | Pricing Date | Cotton - Dryland (Ton) | |
|------------------|--|--------------|--------------|------------------------|--|
| Cotton - Dryland | | Winter 2023 | 2023-04-15 | 1.50 | |

| Income | | | | | |
|-------------------------------|------------------|--------------|-----------|--------------|----------------|
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost |
| Cotton - Dryland | 1.50 | Ton | 11 950.00 | R/ton | R17 925 |
| Gross Production Value | | | | | R17 925 |

| Expenses | | | | | |
|---|------------------|--------------|----------|--------------|----------------|
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost |
| Seed (Cotton) | 5.00 | Kg/ha | 138.07 | R/kg | R690 |
| Fertiliser - Macro elements | 1.00 | Ha | 2 172.63 | R/ha | R2 173 |
| Fertiliser - Micro elements | 1.00 | Ha | 323.23 | R/ha | R323 |
| Fuel (Diesel) | 69.59 | L/ha | 20.81 | R/l | R1 448 |
| Diesel (Picker) | 10.00 | L/ha | 20.81 | R/l | R208 |
| Herbicide | 1.00 | Ha | 528.00 | R/ha | R528 |
| Insecticide | 1.00 | Ha | 755.40 | R/ha | R755 |
| Fungicide | 1.00 | Ha | 0.00 | R/ha | R0 |
| Other Chemicals | 1.00 | Ha | 248.50 | R/ha | R249 |
| High Boy sprayer | 6.00 | Applications | 200.00 | R/ha | R1 200 |
| Insurance - Cotton | 17 925.00 | Rand | 11.00 | % | R1 972 |
| Cotton picker - Dryland | 1.00 | Ha | 2 125.00 | R/ha | R2 125 |
| Packaging - Plastic (Cotton) | 0.65 | Bales | 680.00 | R/bale | R439 |
| Transport Cotton | 1.50 | Ton | 290.00 | R/ton | R435 |
| Mechanization - Repair and Maintenance | 1.00 | Ha | 1 404.08 | R/ha | R1 404 |
| Marketing Cost Cotton - statutorily & Nakpo | 1.50 | Ton | 110.00 | R/ton | R165 |
| Total Direct Cost | | | | | R14 114 |

| Interest | | | | | |
|--------------|------------------|--------------|-------|--------------|--------------|
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost |
| Interest | 5 880.76 | Rand | 12.00 | % | R706 |

| | |
|------------------------------|----------------|
| Total Production Cost | R14 820 |
| Margin Above Cost | R3 105 |
| Breakeven Yield/Ha | 1.24 |
| Breakeven Price/Ton | R9 880 |

| Sensitivity Analysis | | | | | | | | |
|----------------------|---------|---------|---------|---------|---------|---------|---------|--|
| Crop Yield (t/ha) | R11 200 | R11 450 | R11 700 | R11 950 | R12 200 | R12 450 | R12 700 | |
| 0.75 | -R6 420 | -R6 232 | -R6 045 | -R5 857 | -R5 670 | -R5 482 | -R5 295 | |
| 1.00 | -R3 620 | -R3 370 | -R3 120 | -R2 870 | -R2 620 | -R2 370 | -R2 120 | |
| 1.25 | -R820 | -R507 | -R195 | R118 | R430 | R743 | R1 055 | |
| 1.50 | R1 980 | R2 355 | R2 730 | R3 105 | R3 480 | R3 855 | R4 230 | |
| 1.75 | R4 780 | R5 218 | R5 655 | R6 093 | R6 530 | R6 968 | R7 405 | |
| 2.00 | R7 580 | R8 080 | R8 580 | R9 080 | R9 580 | R10 080 | R10 580 | |
| 2.25 | R10 380 | R10 943 | R11 505 | R12 068 | R12 630 | R13 193 | R13 755 | |

| Fertiliser (Macro Elements) | |
|-----------------------------|------|
| N | 31.0 |
| P | 12.0 |
| K | 16.0 |
| Ca | 0.0 |
| Mg | 0.0 |
| S | 2.3 |

GM Cotton Irrigated - Summer

| Cost Guide Figures | | | | | | | profarmer | |
|---|-------------------------|---------------------|----------------|---------------------|---------------------|---------------------|----------------|--|
| Sub Crop | | Publications | | Pricing Date | | Cotton (Ton) | | |
| Cotton | | Summer 2022/23 | | 2022-10-15 | | 5.50 | | |
| Income | | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | | |
| Cotton | 5.50 | Ton | 11 870.00 | R/ton | R65 285 | | | |
| Gross Production Value | | | | | R65 285 | | | |
| Expenses | | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | | |
| Seed (Cotton) | 12.00 | Kg/ha | 138.07 | R/kg | R1 657 | | | |
| Fertiliser - Macro elements | 1.00 | Ha | 16 512.36 | R/ha | R16 512 | | | |
| Fertiliser - Micro elements | 1.00 | Ha | 785.90 | R/ha | R786 | | | |
| Fuel (Diesel) | 72.43 | L/ha | 23.62 | R/l | R1 711 | | | |
| Diesel (Picker) | 40.00 | L/ha | 23.62 | R/l | R945 | | | |
| Herbicide | 1.00 | Ha | 528.00 | R/ha | R528 | | | |
| Insecticide | 1.00 | Ha | 4 157.40 | R/ha | R4 157 | | | |
| Fungicide | 1.00 | Ha | 0.00 | R/ha | R0 | | | |
| Other Chemicals | 1.00 | Ha | 1 468.25 | R/ha | R1 468 | | | |
| High Boy sprayer | 6.00 | Applications | 200.00 | R/ha | R1 200 | | | |
| Insurance - Cotton | 65 285.00 | Rand | 9.00 | % | R5 876 | | | |
| Cotton picker | 1.00 | Ha | 4 500.00 | R/ha | R4 500 | | | |
| Packaging - Plastic (Cotton) | 2.37 | Bales | 680.00 | R/bale | R1 608 | | | |
| Transport Cotton | 5.50 | Ton | 310.00 | R/ton | R1 705 | | | |
| Irrigation - Escom | 800.00 | mm/ha | 6.82 | R/mm | R5 456 | | | |
| Irrigation - Water Board | 800.00 | mm/ha | 1.91 | R/mm | R1 528 | | | |
| Irrigation - Scheduling | 1.00 | Ha | 114.00 | R/ha | R114 | | | |
| Mechanization - Repair and Maintenance | 1.00 | Ha | 1 284.69 | R/ha | R1 285 | | | |
| Pivot Cost - Repair and Maintenance | 1.00 | Ha | 771.00 | R/ha | R771 | | | |
| Marketing Cost Cotton - statutorily & Nakpo | 5.50 | Ton | 110.00 | R/ton | R605 | | | |
| Total Direct Cost | | | | | R52 412 | | | |
| Interest | | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | | |
| Interest | 26 205.93 | Rand | 11.00 | % | R2 883 | | | |
| Total Production Cost | | | | | R55 295 | | | |
| Margin Above Cost | | | | | R9 990 | | | |
| Breakeven Yield/Ha | | | | | 4.66 | | | |
| Breakeven Price/Ton | | | | | R10 054 | | | |
| Sensitivity Analysis | | | | | | | | |
| Crop Yield (t/ha) | R11 120 | R11 370 | R11 620 | R11 870 | R12 120 | R12 370 | R12 620 | |
| 4.00 | -R10 815 | -R9 815 | -R8 815 | -R7 815 | -R6 815 | -R5 815 | -R4 815 | |
| 4.50 | -R5 255 | -R4 130 | -R3 005 | -R1 880 | -R755 | R370 | R1 495 | |
| 5.00 | R305 | R1 555 | R2 805 | R4 055 | R5 305 | R6 555 | R7 805 | |
| 5.50 | R5 865 | R7 240 | R8 615 | R9 990 | R11 365 | R12 740 | R14 115 | |
| 6.00 | R11 425 | R12 925 | R14 425 | R15 925 | R17 425 | R18 925 | R20 425 | |
| 6.50 | R16 985 | R18 610 | R20 235 | R21 860 | R23 485 | R25 110 | R26 735 | |
| 7.00 | R22 545 | R24 295 | R26 045 | R27 795 | R29 545 | R31 295 | R33 045 | |
| Fertiliser (Macro Elements) | | | | | | | | |
| N | 180.0 | | | Kg/ha | | | | |
| P | 50.0 | | | Kg/ha | | | | |
| K | 110.0 | | | Kg/ha | | | | |
| Ca | 18.0 | | | Kg/ha | | | | |
| Mg | 15.0 | | | Kg/ha | | | | |
| S | 24.0 | | | Kg/ha | | | | |

GM Cotton Irrigated Winter

| Cost Guide Figures | | | | profarmer | | | |
|---|-------------------------|---------------------|---------------------|---------------------|---------------------|----------------|----------------|
| Sub Crop | Publications | Pricing Date | Cotton (Ton) | | | | |
| Cotton | Winter 2023 | 2023-04-15 | 5.50 | | | | |
| Income | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| Cotton | 5.50 | Ton | 11 950.00 | R/ton | R65 725 | | |
| Gross Production Value | | | | | R65 725 | | |
| Expenses | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| Seed (Cotton) | 12.00 | Kg/ha | 138.07 | R/kg | R1 657 | | |
| Fertiliser - Macro elements | 1.00 | Ha | 13 527.69 | R/ha | R13 528 | | |
| Fertiliser - Micro elements | 1.00 | Ha | 765.78 | R/ha | R766 | | |
| Fuel (Diesel) | 72.43 | L/ha | 20.81 | R/l | R1 507 | | |
| Diesel (Picker) | 40.00 | L/ha | 20.81 | R/l | R832 | | |
| Herbicide | 1.00 | Ha | 528.00 | R/ha | R528 | | |
| Insecticide | 1.00 | Ha | 4 157.40 | R/ha | R4 157 | | |
| Fungicide | 1.00 | Ha | 0.00 | R/ha | R0 | | |
| Other Chemicals | 1.00 | Ha | 1 468.25 | R/ha | R1 468 | | |
| High Boy sprayer | 6.00 | Applications | 200.00 | R/ha | R1 200 | | |
| Insurance - Cotton | 65 725.00 | Rand | 11.00 | % | R7 230 | | |
| Cotton picker | 1.00 | Ha | 4 250.00 | R/ha | R4 250 | | |
| Packaging - Plastic (Cotton) | 2.37 | Bales | 680.00 | R/bale | R1 608 | | |
| Transport Cotton | 5.50 | Ton | 290.00 | R/ton | R1 595 | | |
| Irrigation - Escom | 800.00 | mm/ha | 8.09 | R/mm | R6 472 | | |
| Irrigation - Water Board | 800.00 | mm/ha | 1.93 | R/mm | R1 544 | | |
| Irrigation - Scheduling | 1.00 | Ha | 114.00 | R/ha | R114 | | |
| Mechanization - Repair and Maintenance | 1.00 | Ha | 1 392.95 | R/ha | R1 393 | | |
| Pivot Cost - Repair and Maintenance | 1.00 | Ha | 1 133.00 | R/ha | R1 133 | | |
| Marketing Cost Cotton - statutorily & Nakpo | 5.50 | Ton | 110.00 | R/ton | R605 | | |
| Total Direct Cost | | | | | R51 588 | | |
| Interest | | | | | | | |
| Product Name | Product Quantity | Measure Unit | Price | Measure Unit | Product Cost | | |
| Interest | 25 793.75 | Rand | 12.00 | % | R3 095 | | |
| Total Production Cost | | | | | R54 683 | | |
| Margin Above Cost | | | | | R11 042 | | |
| Breakeven Yield/Ha | | | | | 4.58 | | |
| Breakeven Price/Ton | | | | | R9 942 | | |
| Sensitivity Analysis | | | | | | | |
| Crop Yield (t/ha) | R11 200 | R11 450 | R11 700 | R11 950 | R12 200 | R12 450 | R12 700 |
| 4.00 | -R9 883 | -R8 883 | -R7 883 | -R6 883 | -R5 883 | -R4 883 | -R3 883 |
| 4.50 | -R4 283 | -R3 158 | -R2 033 | -R908 | R217 | R1 342 | R2 467 |
| 5.00 | R1 317 | R2 567 | R3 817 | R5 067 | R6 317 | R7 567 | R8 817 |
| 5.50 | R6 917 | R8 292 | R9 667 | R11 042 | R12 417 | R13 792 | R15 167 |
| 6.00 | R12 517 | R14 017 | R15 517 | R17 017 | R18 517 | R20 017 | R21 517 |
| 6.50 | R18 117 | R19 742 | R21 367 | R22 992 | R24 617 | R26 242 | R27 867 |
| 7.00 | R23 717 | R25 467 | R27 217 | R28 967 | R30 717 | R32 467 | R34 217 |
| Fertiliser (Macro Elements) | | | | | | | |
| N | 180.0 | | | Kg/ha | | | |
| P | 50.0 | | | Kg/ha | | | |
| K | 110.0 | | | Kg/ha | | | |
| Ca | 18.0 | | | Kg/ha | | | |
| Mg | 15.0 | | | Kg/ha | | | |
| S | 24.0 | | | Kg/ha | | | |

Appendix D: Consultant's Curriculum Vitae

ENVIRONMENTAL SCIENTIST**André Faul**

André entered the environmental assessment profession at the beginning of 2013 and since then has worked on more than 190 Environmental Impact Assessments including assessments of the petroleum industry, harbour expansions, irrigation schemes, township establishment and power generation and transmission. André's post graduate studies focussed on zoological and ecological sciences and he holds a M.Sc. in Conservation Ecology and a Ph.D. in Medical Bioscience. His expertise is in ecotoxicological related studies focussing specifically on endocrine disrupting chemicals. His Ph.D. thesis title was The Assessment of Namibian Water Resources for Endocrine Disruptors. Before joining the environmental assessment profession he worked for 12 years in the Environmental Section of the Department of Biological Sciences at the University of Namibia, first as laboratory technician and then as lecturer in biological and ecological sciences.

CURRICULUM VITAE ANDRÉ FAUL

| | | |
|-------------------|---|--|
| Name of Firm | : | Geo Pollution Technologies (Pty) Ltd. |
| Name of Staff | : | ANDRÉ FAUL |
| Profession | : | Environmental Scientist |
| Years' Experience | : | 22 |
| Nationality | : | Namibian |
| Position | : | Environmental Scientist |
| Specialisation | : | Environmental Toxicology |
| Languages | : | Afrikaans – speaking, reading, writing – excellent English – speaking, reading, writing – excellent |

EDUCATION AND PROFESSIONAL STATUS:

| | | |
|------------------------------|---|--------------------------------------|
| B.Sc. Zoology/Biochemistry | : | University of Stellenbosch, 1999 |
| B.Sc. (Hons.) Zoology | : | University of Stellenbosch, 2000 |
| M.Sc. (Conservation Ecology) | : | University of Stellenbosch, 2005 |
| Ph.D. (Medical Bioscience) | : | University of the Western Cape, 2018 |

| | |
|---------------------|---------------------------|
| First Aid Class A | EMTSS, 2017; OSH-Med 2022 |
| Basic Fire Fighting | EMTSS, 2017; OSH-Med 2022 |

PROFESSIONAL SOCIETY AFFILIATION:

Environmental Assessment Professionals of Namibia (Practitioner)

AREAS OF EXPERTISE:

Knowledge and expertise in:

- ◆ Water Sampling, Extractions and Analysis
- ◆ Biomonitoring and Bioassays
- ◆ Biodiversity Assessment
- ◆ Toxicology
- ◆ Restoration Ecology

EMPLOYMENT:

| | | |
|-----------|---|--|
| 2013-Date | : | Geo Pollution Technologies – Environmental Scientist |
| 2005-2012 | : | Lecturer, University of Namibia |
| 2001-2004 | : | Laboratory Technician, University of Namibia |

PUBLICATIONS:

| | |
|-----------------------------|------|
| Publications: | 5 |
| Contract Reports | +190 |
| Research Reports & Manuals: | 5 |
| Conference Presentations: | 1 |

Appendix D: Tree Information

Trees recorded in quarter degree squares 19118AA (Curtis & Mannheimer, 2005)

| Name | Common_Name | Notes |
|---|-----------------------|---|
| <i>Acacia ataxacantha</i> | Flame-thorn | 0 |
| <i>Acacia erioloba</i> | Camel-thorn | Protected by forestry legislation |
| <i>Acacia fleckii</i> | Sand-veld Acacia | 0 |
| <i>Acacia hebeclada</i> subsp <i>hebeclada</i> | Candle-pod Acacia | 0 |
| <i>Acacia hereroensis</i> | Mountain-thorn | 0 |
| <i>Acacia karroo</i> | Sweet-thorn | 0 |
| <i>Acacia kirkii</i> subsp <i>kirkii</i> var <i>kirkii</i> | Floodplain Acacia | May be declining in Etosha and north-west |
| <i>Acacia luederitzii</i> var <i>luederitzii</i> | Kalahari Acacia | 0 |
| <i>Acacia mellifera</i> subsp <i>detinens</i> | Blue-thorn Acacia | Aggressive Invasive |
| <i>Acacia nilotica</i> subsp <i>kraussiana</i> | Scented-pod Acacia | 0 |
| <i>Acacia reficiens</i> subsp <i>reficiens</i> | Red-thorn | Very Aggressive Invader |
| <i>Acacia tortilis</i> | 0 | 0 |
| <i>Acacia tortilis</i> subsp <i>spirocarpa</i> | Umbrella-thorn | 0 |
| <i>Albizia anthelmintica</i> | Worm-cure Albizia;Aru | The low numbers of young trees recorded is a concern, as is the number of dead trees in some areas. It is Protected by forestry legislation |
| <i>Aloe littoralis</i> | Windhoek Aloe | Potentially threatened by pachycaul trade. Protected by the Nature Conservation Ordinance and listed in CITES Appendix II. |
| <i>Artemisia afra</i> | 0 | 0 |

| | | |
|---|--|---|
| <i>Berchemia discolor</i> | Bird Plum | This species is Protected by forestry legislation, as well as by traditional Owambo cultures for its fruit and shade. The population does not appear to be in any real danger at the moment, but communities could be encouraged to plant this species. |
| <i>Boscia albitrunca</i> | Shepherd's Tree | Although widespread and hardy, it is heavily utilised by people and animals. The difficulty that young plants have in becoming established is a concern, but fortunately there appears to be a health and widespread population of young plants. Protected by forestry legislation. |
| <i>Boscia foetida</i> subsp <i>foetida</i> | Smelly Shepherd's-bush | 0 |
| <i>Caesalpinia rubra</i> | 0 | 0 |
| <i>Catophractes alexandri</i> | Trumpet-thorn;Rattlepod | Invasive in some areas |
| <i>Combretum apiculatum</i> subsp <i>apiculatum</i> | Kudu-bush | 0 |
| <i>Combretum hereroense</i> subsp <i>hereroense</i> | Mouse-eared Combretum | 0 |
| <i>Combretum imberbe</i> | Leadwood | Although heavily utilized by people regrowth is good and growth of young trees is vigorous. Because of its religious importance and many uses, it is protected locally. Old specimens warrant protection as monuments. Protected by forestry legislation |
| <i>Commiphora africana</i> | Hairy Corkwood;Poison-grub Commiphora | 0 |

| | | |
|--|--|---|
| <i>Commiphora angolensis</i> | Sand Corkwood | 0 |
| <i>Commiphora glandulosa</i> | Tall Common Corkwood | 0 |
| <i>Commiphora glaucescens</i> | Blue-leaved Corkwood | 0 |
| <i>Commiphora mollis</i> | Velvet Corkwood | 0 |
| <i>Commiphora pyracanthoides</i> | Fire Thorn Corkwood; Small Common Corkwood | 0 |
| <i>Commiphora tenuipetiolata</i> | Satin-bark Corkwood | 0 |
| <i>Croton gratissimus</i> | Lavender Croton | 0 |
| <i>Croton gratissimus var subgratissimus</i> | 0 | 0 |
| <i>Croton menyhartii</i> | Rough-leaved Croton | 0 |
| <i>Datura spp</i> | 0 | 0 |
| <i>Dichrostachys cinerea subsp africana</i> | Kalahari Christmas Tree; Sickle-bush | Of concern because of its effects on other species (invasive) |
| <i>Dombeya rotundifolia</i> | Wild Pear | Two varieties rotundifolia and velutina. Velutina is endemic and classified as least concern |
| <i>Ehretia alba</i> | White-puzzle Bush | 0 |
| <i>Elaeodendron transvaalense</i> | Transvaal Saffron; Bushveld Saffron | 0 |
| <i>Elephantorrhiza suffruticosa</i> | Skew-leaved Elephant Root | 0 |
| <i>Erythrina decora</i> | Namib Coral-tree | Endemic to Namibia and very uncommon throughout its range Worthy of protection very few young trees. Protected by forestry legislation. |
| <i>Euclea undulata var myrtina</i> | Common Guarri; Mountain Ebony | 0 |
| <i>Ficus thonningii</i> | 0 | 0 |
| <i>Flueggea virosa subsp virosa</i> | White-berry Bush | 0 |

| | | |
|---|--|--|
| <i>Grewia bicolor</i> var <i>bicolor</i> | Two-coloured Raisin-bush | 0 |
| <i>Grewia flava</i> | Velvet Raisin | 0 |
| <i>Grewia flavescens</i> | Sandpaper Raisin | 0 |
| <i>Grewia villosa</i> var <i>villosa</i> | Mallow Raisin | 0 |
| <i>Gymnosporia senegalensis</i> | Confetti Spikethorn | 0 |
| <i>Helinus integrifolius</i> | 0 | 0 |
| <i>Hyphaene petersiana</i> | Makalani Palm | Should be monitored due to extensive harvesting. Protected by forestry legislation |
| <i>Ipomoea adenioides</i> | 0 | 0 |
| <i>Kirkia acuminata</i> | Common Kirkia | 0 |
| <i>Maerua schinzii</i> | Ringwood Tree | Increasingly impacted by humans and giraffes. Protected by forestry legislation |
| <i>Montinia caryophyllacea</i> | Wild Clove-bush | 0 |
| <i>Mundulea sericea</i> | Silverbush | 0 |
| <i>Olea europaea</i> subsp <i>cuspidata</i> | Wild Olive | 0 |
| <i>Opilia campestris</i> var <i>campestris</i> | 0 | 0 |
| <i>Ozoroa paniculosa</i> | Common Resin-bush | 0 |
| <i>Pavetta zeyheri</i> | Small-leaved Bride's-bush | May be declining |
| <i>Peltophorum africanum</i> | Muparara | 0 |
| <i>Philenoptera nelsii</i> subsp <i>nelsii</i> | Kalahari Omupanda; Kalahari Apple-leaf | 0 |
| <i>Rhigozum brevispinosum</i> | Simple-leaved Rhigozum | 0 |
| <i>Searsia marlothii</i> | Bitter Karee | 0 |
| <i>Searsia tenuinervis</i> var <i>tenuinervis</i> | Kalahari Currant | 0 |

| | | |
|---|-----------------------|---|
| <i>Sclerocarya birrea</i> | Marula | Protected locally by communities that use them. Protected by forestry legislation |
| <i>Spirostachys africana</i> | Tambooti | Protected by forestry legislation |
| <i>Tarchonanthus camphoratus</i> | Camphor Bush | 0 |
| <i>Terminalia prunioides</i> | Purple-pod Terminalia | 0 |
| <i>Terminalia sericea</i> | Silver Cluster-leave | 0 |
| <i>Tinnea rhodesiana</i> | Maroon Bells | May be overlooked. |
| <i>Ximania americana</i> var <i>microphylla</i> | Blue Sourplum | 0 |
| <i>Ximania caffra</i> var <i>caffra</i> | Large Sourplum | 0 |
| <i>Ziziphus mucronata</i> | Buffalo-thorn | Protected by forestry legislation |

Appendix E: Proof of Public Consultation

Notified IAPs

| Name | Organisation |
|--------------|---|
| M. Kombanola | Farm Cork FMB/00696, Deal FMB/00698 and Colombo FMB/00694 |
| G. Henning | Farm Cooktown FMB/00695 |
| C. Coetzee | Farm Brisbane Noord FMB/1432/00001 and Bombay FMB/670 |
| J. Murangi | Namwater |
| | Oshikoto Regional Council |

IAPs Notified by Hand Delivered Letter



Public Participation Notification: Environmental Assessment
 Irrigation Based Agriculture Activities on the Farm Cuxhaven Oos No 1278, Oshikoto Region

| Name & Surname | Organisation/Address | Tel / Mobile | Email | Signature | |
|---------------------|----------------------|--------------|-------|-----------|--|
| Mr Martin Kalimunda | Privacy Block | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |

Geo Pollution Technologies
 Irrigation Based Agriculture Activities on Farm Cuxhaven Oos No 1278

 March 2023

Notification Letter



TEL.: (+264-61) 257411 ♦ FAX.: (+264) 88626368
 CELL.: (+264-81) 1220082
 PO BOX 11073 ♦ WINDHOEK ♦ NAMIBIA
 E-MAIL: gpt@thenamib.com

To: Interested and / or Affected Party 16 March 2023
Re: Environmental Assessment and Environmental Management Plan for Irrigation Based Agriculture Activities on Farm Cuxhaven Oos, Oshikoto Region

Dear Sir/Madam

Geo Pollution Technologies (Pty) Ltd was appointed by L M Potgieter to undertake an environmental assessment for irrigation based activities on farm Cuxhaven Oos FMB/1278, in the Oshikoto Region (see location map on page 2). The assessment will be conducted according to the Environmental Management Act of 2007 and its regulations as published in 2012.

Project: Environmental Scoping Assessment and Environmental Management Plan for Irrigation Based Agriculture Activities on Cuxhaven Oos, Oshikoto Region

Proponent: L M Potgieter

Environmental Assessment Practitioner: Geo Pollution Technologies (Pty) Ltd

The Proponent has collectively cleared approximately 76 ha for irrigation by means of centre pivot and drip irrigation systems. The main crops cultivated are vegetables, maize and citrus. Irrigation is from three production boreholes in line with a water abstraction permits as issued by the Ministry of Agriculture, Water and Land Reform. Pending the outcome of the environmental assessment and hydrogeological specialist assessment, an application to increase the total water abstraction per year, may be submitted.

The environmental scoping assessment will include all infrastructure and operational activities associated with the agricultural activities on the farm. This include land clearing, soil preparation, planting, pest control and fertilizer use, harvesting and support services such as electricity supply, staff accommodation and fuel storage.

Interested and affected parties or neighbours are invited to register with the environmental consultant, to receive further documentation and communication regarding the project, or to provide comments related to the project, for inclusion in the assessment. Please register or submit comments at:

Fax: 088-62-6368 or **E-Mail:** cuxhaven@thenamib.com

Should you require any additional information please contact Geo Pollution Technologies at telephone 061-257411.

Sincerely,


Geo Pollution Technologies

Quzette Bosman

Social and Environmental Assessment Practitioner

Directors:

Page 1 of 2
 P. Botja (B.Sc. Hons. Hydrogeology) (Managing)



Project Location

Background Information Document

**ENVIRONMENTAL SCOPING ASSESSMENT AND ENVIRONMENTAL
MANAGEMENT PLAN FOR
IRRIGATION BASED AGRICULTURE ACTIVITIES ON FARM CUXHAVEN
OOS NO 1278, OSHIKOTO REGION
BACKGROUND INFORMATION DOCUMENT**



Prepared by:



Prepared for:

L M Potgieter

March 2023

1 INTRODUCTION

Geo Pollution Technologies (Pty) Ltd was appointed by L M Potgieter (the Proponent) to undertake an environmental assessment for irrigation activities on the farm Cuxhaven Oos FMB/1278 in the Oshikoto Region (Figure 1-1). Currently the Proponent irrigates a combined area of 76 ha from production boreholes on the two farm. Pending the outcome of an environmental assessment, the total water abstraction per year, may be increased. Crop cultivation will focus on maize, vegetable and citrus production which will mainly be irrigated by means of centre pivot, drip and micro sprayer systems.

An environmental clearance certificate (ECC) for the operations is required as per the Environmental Management Act No. 7 of 2007 (EMA). A scoping environmental assessment report (SR) and an environmental management plan (EMP) are proposed to be submitted to the Ministry of Environment, Forestry and Tourism's Department of Environmental Affairs (DEA) in support of an application for an ECC. The environmental assessment will include all operational activities associated with the agricultural activities of the Proponent.




Figure 1-1 Project location

2 PURPOSE OF THE BID

With this background information document (BID), GPT aims to provide interested and affected parties (IAPs) with information about the project and interact with them regarding it. All IAPs are therefore invited to register with GPT for the project in order to:

- ◆ Provide GPT with additional information which should be taken into account in the assessment of impacts;
- ◆ Share any comments, issues or concerns related to the project; and
- ◆ Review and comment on the reports (SR and EMP).

3 PROJECT DESCRIPTION

Activities associated with the project have been divided into the following phases: planning, maintenance/construction, operational and the decommissioning phase. A brief outline of expected activities for each phase is detailed below.

3.1 PLANNING PHASE

While planning for operations, construction / maintenance activities and decommissioning of the farm, it is the responsibility of the Proponent to ensure they are and remain compliant with all legal requirements. The Proponent must also ensure that all required management measures are in place prior to and during all phases, to ensure potential impacts and risks are minimised. Typical planning activities include:

- ◆ Obtain permits and approvals from local and national authorities including Ministry of Agriculture, Water and Land Reform.
- ◆ Make provisions to have a health, safety and environmental coordinator to implement the EMP.
- ◆ Ensure provisions for a fund to cater for environmental incidents (e.g. pollution) and ecological restoration are made.
- ◆ Ensure all appointed contractors and employees enter into agreements which include the EMP.
- ◆ Establish and/or maintain a reporting system to report on aspects of construction activities, operations and decommissioning as outlined in the EMP.

3.2 CONSTRUCTION AND MAINTENANCE PHASE

Some construction activities will form part of the development of the farm. Maintenance continues on a daily basis and may also include some construction activities. Maintenance include minor repairs to infrastructure, general upkeep of buildings, servicing of vehicles, etc.

3.3 OPERATIONAL PHASE

The main operational activities pertains to land clearing and preparation, planting, water abstraction and irrigation, pest control and harvesting. High value crops are produced in the form of vegetables, maize and citrus. Historically, areas of the farm were cleared for a combination of dryland and irrigation-based crop cultivation. However, current areas being irrigated, by means of centre pivot systems, amount to approximately 76 ha with the potential of further cultivation on 25 ha. Historically cleared areas for crop cultivation, excluding the existing and potential irrigation areas, amount to approximately 172 ha. Active rangeland improvement was and is continually conducted over the northern portions of the farm. All areas which have undergone rangeland improvement (through bush-thinning) amount to approximately 89 ha. Water for irrigation is abstracted from production boreholes located on the farm. Operations also include activities typical of farming, inclusive of management of support services such as electricity supply, waste handling and domestic effluent disposal.

3.4 DECOMMISSIONING PHASE

Decommissioning is not foreseen during the validity of the ECC. Decommissioning will however be assessed. Should decommissioning occur at any stage, rehabilitation of the area may be required. Decommissioning will entail the complete removal of all infrastructure including buildings and underground infrastructure. Pollution present on the site, if any, must then be remediated.

3.5 PRELIMINARY IDENTIFIED IMPACTS

During the environmental assessment all components of the environment will be considered, however only those components which are being impacted on significantly, or are deemed to be sensitive, will be assessed. These include the following:

- ◆ Health and safety risks
- ◆ Soil and groundwater pollution
- ◆ Over abstraction of groundwater
- ◆ Fire risks

- ◆ Waste and effluent generation and disposal
- ◆ Traffic
- ◆ Noise
- ◆ Visual impact
- ◆ Ecosystem and biodiversity impacts
- ◆ Socio-economic contributions

4 PUBLIC CONSULTATION

GPT invites all IAPs to provide in writing, any issues and suggestions regarding the development. This correspondence must include:

- ◆ Name and surname,
- ◆ Organization represented or private interest,
- ◆ Position in the organization,
- ◆ Contact details, and
- ◆ Any direct business, financial, personal or other interest which you may have in the approval or refusal of the application.

All contributions become public knowledge and will be circulated along with the reports as per the EMA requirements.

The comments, inputs and suggestions will also be submitted to the DEA along with how any issues have been addressed in the SR. The public participation process will remain ongoing during the environmental assessment.

The project team may be contacted on the contact details below



Geo Pollution Technologies (Pty) Ltd.

Telephone: (+264-61) 257411

Fax: (+264) 88626368

E-mail: cuxhaven@thenamib.com

Your Rights as an IAP according to the Environmental Management Act, No7 of 2007, Government Notice No. 30 (Environmental Impact Assessment Regulations)

Section 23. (1) A registered interested or affected party is entitled to comment in writing, on all written submissions made to the Environmental Commissioner by the applicant responsible for the application, and to bring to the attention of the Environmental Commissioner any issues which that party, believes may be of significance to the consideration of the application, as long as -

- (a) comments are submitted within 7 days of notification of an application or receiving access to a scoping report or an assessment report;*
 - (b) the interested and affected party discloses any direct business, financial, personal or other interest which that party may have in the approval or refusal of the application.*
- (2) Before the applicant submits a report compiled in terms of these regulations to the Environmental Commissioner, the applicant must give registered interested and affected parties access to, and an opportunity to comment in writing on the report.*
- (3) Reports referred to in sub regulation (2) include (a) scoping reports; (b) scoping reports amended and resubmitted; (c) assessment reports; and (d) assessment reports amended and resubmitted.*
- (4) Any written comments received by the applicant from a registered interested or affected party must accompany the report when the report is submitted to the Environmental Commissioner.*
- (5) A registered interested or affected party may comment on any final report that is submitted by a specialist reviewer for the purposes of these regulations where the report contains substantive information which has not previously been made available to a registered interested or affected party.*

Section 24: The applicant responsible for an application must ensure that the comments of interested and affected parties are recorded in reports submitted to the Environmental Commissioner in terms of these regulations, and comments by interested and affected parties on a report which is to be submitted to the Environmental Commissioner may be attached to the report without recording those comments in the report itself.

NEWS IN SHORT

Hippo meat aids vulnerable community

A poached hippo was discovered in the Myunke area of the Zambezi Region last Sunday. Its meat was subsequently distributed to a vulnerable community.

According to spokesperson Romeo Mnyanda, the environment ministry responded to a report of a dead hippo, and upon closer inspection by staff members, it was determined that the hippo had been shot. Mnyanda said the meat from the hippo was distributed to a community in the Kyarwan village, about 25 kilometres west of Katima Mulilo.

"The community there is in a destitute situation. They are a vulnerable community, since they are part of the San community. Over 90% of them do not work and therefore it was deemed necessary that we assist them."



CONSERVATION INITIATIVE: A report on the first-ever synchronised and coordinated aerial survey of Africa's largest savannah elephant populations will be made public by May/June. PHOTO: KAZANKE ZAMBEZI TRANSFRONTIER CONSERVATION AREA

KAZA elephant survey report public in June

A poached hippo was discovered in the Myunke area of the Zambezi Region last Sunday. Its meat was subsequently distributed to a vulnerable community.

According to spokesperson Romeo Mnyanda, the environment ministry responded to a report of a dead hippo, and upon closer inspection by staff members, it was determined that the hippo had been shot. Mnyanda said the meat from the hippo was distributed to a community in the Kyarwan village, about 25 kilometres west of Katima Mulilo.

"The community there is in a destitute situation. They are a vulnerable community, since they are part of the San community. Over 90% of them do not work and therefore it was deemed necessary that we assist them."

Sun

BLASTS 'ARCHAIC' LAW

Lobby group pushes for decriminalisation of sodomy

The LGBTQI umbrella network hosted a media information-sharing workshop last weekend.

JEMMAH NDEBELE WINDHOEK

The Diversity Alliance of Namibia (DAN) has reiterated its stance on what it called the "archaic" sodomy law.

Sodomy is criminalised by the Roman-Dutch common law of Namibia, which was inherited from South Africa at independence.

While the sentence for those found guilty under this law is not clear, currently, the legislation - along with the common law crime of "unnatural sexual offences" - criminalises consensual sexual acts between men. DAN, an umbrella network of over 10 LGBTQI human rights organisations in Namibia, made its stance clear at a media information-sharing workshop held at Heja Lodge last weekend.

A report detailing considerations from a panel discussion held early last year was also launched at the two-day media convening.



THE TIME IS NOW: The Diversity Alliance of Namibia launched a report on a discussion it held last year on the sodomy law. PHOTO: JEMMAH NDEBELE

"Decriminalisation is not only for the repeal of the sodomy offence, but also to grant legal footing to LGBTQI persons to engage with different sectors, including economic empowerment, access to legal services beyond health, voting and civic participation in politics - especially when the person does not conform to the particulars given on their identity document.

"Discrimination [against these persons occurs] daily and is unrelenting, across services and sectors in all aspects of life" the report read.

'The time is now'

Programme facilitator and activist Abigail Solomons said the time is now for media practitioners to join hands with the community and push for the decriminalisation of sodomy in the country.

"The lives of our gay children are at stake here."

"The time is now' is not just the launch of a report, but of an era. The time for change is the present. This is an important year for the community. We are living in history. Moreover, the workshop saw members of the LGBTQI commu-

nity airing their disdain about the lack of media coverage, further urging media practitioners to be more mindful about the way stories regarding the community are covered, and their use of language when doing so.

Additionally, the role the media plays in the decriminalisation of the sodomy law was also highlighted.

Economic exclusion

Members of the LGBTQI community emphasised the fact that they are economically excluded.

"We should understand that we cannot have a healthy, equal, equitable economy within the country if we do not have a healthy society - if we live in a society that does not want to unlearn their modes of oppression.

"So, this is why I think Namibia should strive for economic inclusion. When I speak about economic inclusion, I mean ensuring there is structural change that allows LGBTQI persons to be able to fully, equitably and equally participate in the economy.

"We are all serious about this country," one of the community members, who preferred to remain anonymous, said.



HOUSING: The construction of 70 affordable National Housing Enterprise homes is underway at Okahao.



PROGRESS: Okahao mayor Cornelia Iyambula. PHOTOS: ENZO AMBELE

OKAHAO MAYOR APPLAUDS DEVELOPMENT

ENZO AMBELE OKAHAO

Okahao mayor Cornelia Iyambula says she is satisfied with the influx of commercial entities at the town.

The town council will make land more affordable to attract investors in order to speed up development, she added.

Okahao was proclaimed a town in 2005 and the council started operations in 2006, with the mandate of providing services to residents.

In an interview with Namibian Sun this week, Iyambula said development in the town is progressing, fuelled by council, corporate institutions and the business community.

The town has about seven developmental projects underway, which includes the upgrading of a section of Fidel Castro Street from gravel to bitumen stand-

ard - to the tune of N\$4.2 million.

The recent upgrade of the town council's office added additional office space for a NaTis centre, which enabled the town to provide vehicle registration and licence renewal services.

"Our NaTis office will start offering testing for learners' licences within the coming two weeks as we have just received equipment," Iyambula said.

She added that the council is set to add new vehicles to its operational fleet to improve service delivery.

Boosting economy

According to the mayor, the council recently allocated land for the construction of the International University of Management's (IUM) Omusati Region campus.

"This campus will boost the economy of the town as students will now be based in Okahao for ter-

tiary education," she said.

"We have taken all our council staff members for training to ensure that we have a common understanding of the system in terms of organisational strategies and objectives," Iyambula noted.

Meanwhile, the town council is planning on revising the Elhao Cultural Expo, which could not take place for the past three years due to the Covid-19 pandemic.

The expo, which - according to council - has the potential to drive raise funds and create a platform for the youth to showcase their skills or sell their products, is scheduled to take place from 26 June to 2 July.

Junias Jakob, the town's CEO, called on the youth to participate in the expo as it will enable them to expand their existing businesses and also network with other entrepreneurs.

PUBLIC PARTICIPATION NOTICE

ENVIRONMENTAL SCOPING ASSESSMENT: BROADBAND WIRELESS AGRICULTURAL ACTIVITIES ON WAKKOS FARM IN ABERHAR, OSHANA REGION

Geo Pollution Technologies (Pty) Ltd. was appointed by Oshana Farming CC, Ltd (Pvt) Ltd, D. Buschhoff and Adriaan Sobot CC to undertake environmental scoping assessments for irrigation related activities on farms Broadway FMB070, Portion 1 of Broadway Noord FMB1432, Cudavas Oos FMB00278, Groot Bontia FMB01343 and Oos Wakkos FMB00225. Additional and location information can be obtained at: <http://www.thesunil.com/nam/en/activities.html>

The environmental assessments will be conducted according to the Environmental Management Act of 2007 and its regulations as published in 2012.

The Proponents engage by means of onsite visits, interviews systems and deep interviews. Water for these purposes is abstracted from production boreholes. All relevant operational activities will be included in the assessments.

All interested and affected parties are invited to register with the environmental consultant. By registering you are provided with the opportunity to share any comments, issues or concerns related to the project, for consideration in the environmental assessment. Additional information can be requested from Geo Pollution Technologies. All written comments and concerns should be submitted to Geo Pollution Technologies by 29 March 2023.

Quetta Bosman
Geo Pollution Technologies
Tel: +264-61-257411
Fax: +264-88020588
E-Mail: abecuda@thesunil.com

NEWS IN SHORT

Namibia 'must interrogate greenwashing'

There is a lack of harmony between the politics and international treaties Namibia has signed and the country's ability to actually reach targets set out by these instruments.

This is according to political scientist and author Professor Andre du Pisani, who spoke at a recent panel discussion on corruption and greenwashing. "We are required to follow and sign off on all these international agreements and we say we will meet these targets, but we know in our hearts of hearts that we would not be able to," he said.

He urged Namibians to move away from Western traditions and towards people-based, sustainable development with a focus on efficiencies.

Meanwhile, anti-corruption activist Professor Chuma Nwankwo said when Namibians look at the topic of greenwashing, they must ask themselves at what point they would assess whether an entity entering their industries is really a green enterprise. Greenwashing occurs when an organisation spends more time and money marketing itself as environmentally-friendly than actually minimising its environmental impact.

- JEMIMA BEUKES

FMD control measures lifted in Zambezi

All control measures for foot-and-mouth disease (FMD) were lifted in the Zambezi Region last week.

According to the agriculture ministry, these restrictive measures were imposed in the region as a result of an outbreak which occurred in Mbalalante in the Karibe South constituency last October. "The last confirmed case was reported on 29 November 2022 and according to Namibia's FMD contingency plan, restrictions in the infected zone can be lifted three months after the last confirmed case," the ministry's chief veterinary officer, Dr Albertina Shilongo, said.

"All FMD restrictive measures that were imposed as a result of the outbreak are therefore lifted with immediate effect."

She added that, to date, a 92% mass vaccination of cattle in the affected area has been achieved.

- ELIANE SMIT

OTHER AIRLINE 'SIMPLY COMPETITIVE'

Etosha hopes to fly by June

Fly Etosha plans to be up and running within a few months.

OGONE TLHAGE WINDHOEK

Fly Etosha says it intends to be airborne by June if all its ducks are in order.

These are the sentiments of representative William Ekandjo, who said the company has every intention of becoming operational soon, "despite an earlier bid by local aviation company FlyNamibia to challenge" its plans. FlyNamibia has since abandoned its objections.

According to Ekandjo, Fly Etosha ticks all the boxes in terms of

the regulatory requirements to become an airline.

FlyNamibia's complaint was simply due to the competitive nature of the aviation sector, he said.

Documents were submitted through FlyNamibia and Westair's attorneys, Ellis and Partners, opposing an application by Fly Etosha for a non-scheduled air transport service licence.

According to the documents, Ekandjo failed dismally to provide sufficient proof that he will be able to render a safe, satisfactory and reliable air service in respect of continuity, frequency, punctuality, reasonable charges and general efficiency.

Focused

"We are focused on what we are



AMBITIOUS: Fly Etosha representative William Ekandjo. PHOTO: NAMIBIA MEDIA SERVICES

doing. This is a field we know very well, so we didn't even bother [with FlyNamibia's opposition] because we are going in for competition," Ekandjo said.

"We satisfy all requirements," he added. "Funds, one could say 'OK, it's not enough, maybe, but what amount [would be enough] because we could raise all the funds."

The team behind Fly Etosha was equally up to task, he said.

"Regulatory requirements, we are top class. We have been running these requirements for so many years. We shall make sure that this airline shall fly all over the world within two years," he added.

Ekandjo admitted that Fly Etosha may not immediately acquire all aircraft types it deemed necessary for its operations, but said this was not a concern for it to get operational approval.

"Equipment, those aeroplanes are available everywhere. Maybe the ones we are crying for the most, maybe they are difficult to get simply because of the demand for those aircraft."

'I coined the name'

Ekandjo further alleged that he initially coined the name FlyNamibia.

"I had the paperwork for FlyNamibia. All by surprise, I heard you have to get away from that name because that name was registered as a trademark and I foresaw that this thing is going to take long."

"If I am really aiming to have an air service licence by this year, an airline that must take off this year, I think I must give that a break," he said.

SARDAROV'S SCHOOL A BEACON OF HOPE

JEMIMA NDESEBE WINDHOEK

Russian billionaire and Marula Game Ranch owner Rashid Sardarov's pledge to build a combined school to the tune of N\$40 million has transfused the lives of farmworkers with hope.

The construction of the school will be privately funded by Sardarov. "My client will spend N\$20 million per year in expenditures, salaries, administration, learners' care, tuition, boarding and clothing, etc," lawyer Sisa Namandje said.

He added that the school will essentially be for the children of farmworkers. The school - which will teach grades 0 to 12 - will carry all costs on behalf of learners.

"We are building the school for 140 workers' children, because it has happened so often that we lose good workers because there is no support for their children who do not live on the property. Currently, we are busy with the architectural work and we hope to start construction at the end of this month

and have it complete by next year," the ranch's general manager, Johan Kotze, said.

Orphan programme

Meanwhile, Namandje and Kotze said Sardarov is looking into an orphan programme, which will take in selected learners from all 14 regions of the country and cover the cost for their welfare and academic expenses.

"At this stage, the school will only be for workers' children and selected orphans. The idea is to break the cycle of workers' children being separated from their parents without the parents having to worry about costs," Namandje said.

Dorobab residents have shown great excitement for the new development, despite the initiative being limited to employees' children.

"We would also like our kids to attend the school and stand a chance at having a future brighter than what we currently have. We hope that this school can bring change to our community" community representative Aitsoab said.

PUBLIC PARTICIPATION NOTICE

ENVIRONMENTAL SCOPING ASSESSMENT: IRRIGATION BASED AGRICULTURAL ACTIVITIES IN VARIOUS FARMS IN ARIKARA, OSHANA AND KARIBE

Geo Pollution Technologies (Pty) Ltd was appointed by Godelstein Farming CC, LM Purgers, D Boshoff and Adressita Sales CC to undertake environmental scoping assessments for irrigation based activities on farms Bonhef FM2670, Portion 1 of Bonhef Noord FM291432, Cuckoo's Nest FMB01278, Great Bonhef FMB01343 and Old Sandfield FMB01233. Additional and location information can be obtained at <http://www.themarb.com/projects/projects.html>

The environmental assessments will be conducted according to the Environmental Management Act of 2007 and its regulations as published in 2012.

The Proponents engage by means of scoping pivot, main sprayer systems and drip irrigation. Water for these purposes is abstracted from production boreholes. All relevant operational activities will be included in the assessments.

All interested and affected parties are invited to register with the environmental consultant. By registering you are provided with the opportunity to share any comments, issues or concerns related to the project, for consideration in the environmental assessment. Additional information can be requested from Geo Pollution Technologies. All written comments and concerns should be submitted to Geo Pollution Technologies by 29 March 2023.

Qazette Swanson
Geo Pollution Technologies
Tel: +264-61-257411
Fax: +264-882-0368
E-Mail: swanson@themarb.com



LiquorShop

WHISKY WEDNESDAY

FINE WHISKY

WEDNESDAY 22 MARCH 2023

199⁹⁹

OFFER VALID ON WEDNESDAY 22 MARCH 2023 ONLY. PRICES APPLY TO 500ML BOTTLES. QUANTITIES LIMITED. WE RESERVE THE RIGHT TO LIMIT QUANTITIES. AND TRADING PLEASE. SELECTED ITEMS MAY NOT BE AVAILABLE AT CERTAIN STORES. VAT INCLUSIVE. STOCKS & CLOSING TIMES ENFORCED. LICENSING ONLY AVAILABLE AT STORES WITH A VALID LICENSING AUTHORITY. CHECK FOR SALE TO PERSONS UNDER THE AGE OF 18 YEARS. WHILE STOCKS LAST.

99% VOLUME BOTTLE

Not for Persons Under the Age of 18. Drink Responsibly.

Press Notice: Die Republiek 15 and 22 March 2023

| | | | | | |
|---|--|---|--|---|--|
| 2 NUUS | | Republiek | | Woensdag 15 Maart 2023 | |
| KONTAKPERSONE | | GENL. MURTALA MUHAMMEDRYLAAN, POSBUS 3436, WINDHOEK TEL: 061 297 2000 VOLG ONS OP: | | WEER | |
| REDAKTOER Frank Steffen 081 24 0882 061 297 2316 fsteffen@az.com.na | | HOUDELIKREDAKTOER Henriette Lamprecht 081 350 3801 061 297 2035 henriette@republiek.com.na | | SPORTNUUS Andrew Poolman 081 247 2837 061 297 2011 andrew@republiek.com.na | |
| REDAKTOER Carasso Steiner 081 239 7654 061 297 2002 carasso@spoorj.com.na | | LEWISOORD Dita Fieck 081 299 1281 diti@wonga.com.na Faks: 064 403 451 Dorena Gases 081 550 0815 | | STREKKE Gushetha: Elvira Hartweg 081 737 3235 Baka: Teyoboo Haidala 081 338 3812 Dehagwa: Erico Amala 081 569 8675 Henda: Kanya Kambova 081 724 1044 | |
| WEBWERFkode: 8329 | | REPUBLIEK-REPUBLICOM.NA 1 Publikasie van ANDREA METZ HOLLINGS (Pty) Ltd, gedruk deur NEWSPRINT NAMIBIA (Pty) Ltd. | | BINNELAND: Somrig en warm in die suidweste. Eiers gedroeflik bewolk en warm tot baie warm met enkele donderbuie. Geseleerde donderbuie sal in die ooste en suidoste voorkom. Windrige toestande sal voortduur. KUS: Gedroeflik bewolk en koel tot warm. GETYE BY WALVISBAAI: H: 09:01 L: 15:15 H: 22:05 | |

Outjo

VAN BL. 1

"Ous het dit op 'n mooi manier probeer doen, deur vergaderings en samewerkingsoostel-le, maar daar het niks van gekom nie. "Toe ons begin moeilik raak omdat niks gebeur nie en al die gemeenskap se werk in hul gewig betragte word, toe word dit 'n politieke kwessie. Dit een rede waarom ek my intussen begin ontroek het."

HULP GEWEIER

"Ous het met 'Droom vir Outjo' se gebl' n professionele hidrolog aangestel, wat vir die munisipaliteit gewys het waar boorgate gesink behoort te word. Die eerste van die boorgate lewer 'n vullinhoudre 60 000 liter per uur. Hulre rig dit egter net nie in nie," se Verster. "Ous het aangeblid dat die gemeenskap die pompe koop en die koste word van die



Freddy

VAN BL. 1

Tog is sy nuwe ruete minder gunstig, wat beteken dit sal nie weer kan versterk nie en sal hopelik heeltemal uitroei.

Freddy het egter alle versigtige outroefas die langsligste trapas sikkoon op rekord, wat teen gister reeds 38 dae lank was.

Dit is ook een van net vier sikkone wat die hele Indiese Oseaan van wes na oos lank deurkruis en boonop 'n U-draai teen die Mosambiese Ius kon maak en na die Mosambiek-kanaal terugbeweeg het.

Hoërfhof lewer uitspraak

Roofvoëls bly by N/a'an ku sê

Die voëls sluit in 'n Kaapse aasvoël-kuiken en sy ouers, twee roofarende, 'n bateleurarend, 'n Afrika-grootjagarend, 'n reusooruil en 'n gevlekte ooruil.

Kristien Kruger

Die N/a'an ku sê-stigting is nou die permanente tuista van noge hedrigle roofvoëls in-gevolge 'n beslissing van die Windhoekse hoërfhof dat die voëls in hul song moet bly.

Dit volg naulaf N/a'an ku sê op 18 Julie 2022 'n dringende aansoek teen die ministerie van omgewing, bosbou en toerisme ingedien het met die versoek dat die voëls vanaf die Hare and Endangered Species Trust (Best) na die stigting se wildreservaat geneem en daar versorg word.

PHOENIX EN KIE

Toe hierdie hofsaak begin het, het die Kaapse aasvoël-broeijaar wat by best aangehou was, een eier gehad wat intussen uitgebroei het. Die negre roofvoëls sluit daarom 'n Kaapse aasvoël-kuiken in, met die naam Phoenix.

Phoenix se ouers is tans die enigste broeijaar in aanhouding in die land. "Phoenix se ouerling is noodsaaklik vir die wêreldbewaring van hierdie spesie," het N/a'an ku sê in November in 'n mediaverklaring gesê.

Die ander voëls sluit in Phoenix se ouers (Halle en Nesher), twee roofarende (Tom Tom en Aquila), 'n bateleurarend (Olle), 'n Afrika-grootjagarend (Phabo), 'n reusooruil (Die Boy) en 'n gevlekte ooruil (Poppy).

WETSARTIKEL VAL

N/a'an ku sê is van die mening dat die wek van natuurbeouders blemmer word deur die onredelike optrede van die ministerie.

"N/a'an ku sê is daar toe verbind om hierre lous te verbeter om Namibie 'n wêreldleier in dienselwing en-bewaring te maak," lous die mediaverklaring.

N/a'an ku sê was suksesvol in sy grondwetlike uithaling van artikel 80(C) van die Natuurbewarings-ordnansie, nr. 4 van 1975. Die artikel het gestel dat die Minister nie verplig word om enige roede te verskaf vir sy weiering van 'n permit, lisensie, registrasie, goedkeuring, toetsing of vrystelling nie. Die artikel is gevolglik ongondwettig verklaar. - kristien@republiek.com.na



Die Kaapse aasvoëlkuiken met die naam Phoenix. 1502/ARRE

PUBLIC PARTICIPATION NOTICE

ENVIRONMENTAL SCOPING ASSESSMENT IRRIGATION BASED AGRICULTURAL ACTIVITIES ON KARIBO'S FARMS IN ASENAR, OSHANA REGION.

Geo Pollution Technologies (Pty) Ltd was appointed by Grubben Farming CC, LM Projector, D Boschhoff and Akonduu Salihi CC, to undertake environmental scoping assessments for irrigation based activities on farms Bushby FMB0167, Paratu 1 of Brisbane Noord FMB01452, Cordatus Oos FMB01278, Great Bourse FMB01341 and Oid Sankhfeld FMB01231. Additional scoping and location information can be obtained at: <http://www.thegrowth.com/projects/projects.html>

The environmental assessments will be conducted according to the Environmental Management Act of 2007 and its regulations as published in 2012.

The Proponents invite by means of public post, micro speaker systems and drip irrigation. Water for these purposes is abstracted from production boreholes. All relevant operational activities will be included in the assessments.

All interested and affected parties are invited to register with the environmental consultant. By registering you are provided with the opportunity to share any comments, issues or concerns related to the project, for consideration in the environmental assessment. Additional information can be requested from Geo Pollution Technologies. All written comments and concerns should be submitted to Geo Pollution Technologies by 29 March 2023.

Quazile Bosman
 Geo Pollution Technologies
 Tel: +264-61-257411
 Fax: +264-68263108
 E-Mail: quazile@thegrowth.com

TOT SIENS, LA NIÑA

Die La Niña-toestande in die tropiese Stille Oseaan het intussen ook tot 'n einde

Monah

VAN BL. 1

Beeds met haar eerste modevertoning waaraan sy deelgeneem het, is haar skeppings goed ontvang, vertel Monah tuis.

Met haar tweede modevertoning het sy besluit om ook plusgrootte skeppings deel van die looppad te maak.

"Ek glo elke vrou is mooi, ongewoon haar figuur."

Haar pad het met Doreed Valkonen op Facebook gekruis waar lausgeniede 'n beroep op Namibiese ontwerpers gedoen het om ontwerpe voor te lê waaruit sy haar lesse vir die Graft-hanklikheidsbal sou maak.

Monah het vier skutse per e-pus gestuur en twee weke later het Valkonen lous weet sy gaan een van Monah

Creations se skeppings met die Namibiese landskap op inspirasie dra.

"Dis vir my baie belangrik dat my kleinste deel deel van my ontwerp en konsep."

Sedert 2019 wat sy vultyde by Monah Creations betrekke is, verskaf sy werk aan sewe mense en erken "dit is baie werk, jy moet dit goed doen en ten volle toegewyd wees"

Haar gunsteling-kledingstuk om te ontwerp en te maak, is aendrokke, "maar ek doen alles. Dis belangrik dat die klient gemaklik en geïnspireerd voel wanneer hulle my ontwerpe dra."

Sy wil graag aan die Windhoekse Modeweek deelneem en beplan om in die "stil maande" klein wat dra-gereed te maak. Volg haar (Monah Creations) op Facebook en Instagram.

- heri@republiek.com.na

| | | | | |
|--|--|--|--|--|
| | COMMERCIAL PROPERTY TO RENT | | | CONTACT LLEWELLYN KRUGER AT +264 81 285 7686 |
| | Location: 78 Gaf'Porton, Muldenweg Ave East, Windhoek, Namibia | Type of Property: Commercial Property Description: Office Building Block J Occupatio Date: Immediately Office Floor Area: 470 m² | | |

» Wêreldwaterdag is onlangs gevier

Waterprobleme knel Groot Aub stéeds

Die regering is op soek na oplossings vir Groot Aub, waar die vraag na water groter is as wat poorgate kan voorsien, terwyl boerderybedrywighede die beskikbare water besoedel.

Evira Haningh

Die waterkwessies in Groot Aub is tweeledig. Eerstens oorskry die vraag na water verskaffing uit poorgate tot so 'n mate dat daar nie genoeg water beskikbaar is nie, terwyl dit ook toenemend besoedel word as gevolg van menslike aktiviteite.

Dit aldus die minister van landbou, water en grondhervorming, Calle Schlettwein, tydens die viering van Wêreldwaterdag, ook Wêreld-vleiland-dag, wat die afgelope Vrydag op Groot Aub gevier is. "Dit is tyd vir die herstel van vleilande", noek "Verstelling van verandering deur vennootskappe en samewerking" was die onderskeie temas vir die wêreld-dae.

Schlettwein het in sy toespraak gesê, soos voorgesê deur die ministerie se uitvoerende direkteur, Nallyskupi Nghitshwamata – dat die vraag na water by Groot Aub oor die afgelope jare gegroei het vanweë die bevolkinggroei wat in die nedersetting veras word. Mensê het hêself hier som wêreld op soek na behuising en



Calle Schlettwein het in sy toespraak gesê dat die vraag na water by Groot Aub oor die afgelope jare gegroei het.

beter geleentheid.

Hy sê verder studies toon dat die water in die gebied byvoorbeeld nitrate bevat, wat met die reën versenselwig word.

"Ons moet seker maak dat die heperkte water wat beskikbaar is, skoon gelou word deur maatskaps in te stel sodat ons die water kan bestuur."

"Dit beteken ons moet gebiede albei vir die aanbou van diere en afdal bestuur sodat dit nie in ons ondergrondse water beland nie.

"Ons moet sanitasiedienste verbeter om 'bostoilette' te vermy en persoonlike higiëne te bevorder..."

Schlettwein het gesê hy en die minister van stedelike en landelike ontwikkeling, Erastus Untch, het Groot Aub in Februarie besoek om vas te stel wat die probleem met die water is en moontlike oplossings te ondersoek.

"Ons wou dat Groot Aub nou deel van die Windhoek-gebied is, en terselfdertyd word die gemeenskap

deur waterverskaffings- en sanitasieweessies in die gasig gestaat."

Hy sê die regering en die Windhoekse munisipaliteit kan nie albees probeer om die verskaffing met soortgelyke waterkwessies te help nie, maar benodig vennote in die private sektor.

Intussen het Alka Bhatia, plaaslike verteenwoordiger van die Verenigde Nasies se Ontwikkelingsprogram (UNDP), daareop geuus dat Namibiê die droogte land suid van die Sahara

is en terselfdertyd ook baie kwesbaar vir klimaatverandering is. Terselfdertyd maak 80% van Namibiê se nywerhede en inwoners op ondergrondse water vir water-voorsiening staat.

Calle Schlettwein
OPGEWINGS-MINISTER

"Ons moet sanitasiedienste verbeter om 'bostoilette' te vermy en persoonlike higiëne te bevorder..."

Sy het bygevoeg dat – om kwesbare gebiede soos Namibiê te help om beter veerkragtigheid teenoor klimaatverandering te toon – ondersteun VN-agentskappe die regering met die volhoubare bestuur van natuurlike hulpbronne, veral as dit by ondergrondse water en vleilande kom.

Sy het na kleiner projekte soos die Cuxhaven-projek verwys – wat weerk doen vir verbetere watersekerheid en gemeenskaplike werkkragtigheid in die aangrensende Cuxhaven- en Namibia-oongrondswaterkomitee.

www.republiekain.com

Ons is moeg, sê Dordabis se inwoners

Jemimah Ntshole

Die meeste inwoners van Dordabis – sowat 80 km oos van Windhoek – word "gedwing om in arm omstandighede te woon".

Die gemeenskap, wat geteister word deur 'n gebrek aan werkgeleentheid, sanitasie en grond, is desperaat vir die regering om in te gryp om hul lewensstandaarde te verbeter. "Ons jongmense doen niks nie en sê hulle is moeg. Daar is geen opleidinggeriewe om hulle vaardighede te leer nie. Ons is moeg. Ons is omring deur private eiendomme en vyurmaskehouer wat ons nie mag gebruik nie," sê die 78-jarige gemeenskapverteenvoerder, Alfrus Namah, tydens 'n besoek aan die nedersetting.

"Die grond wat ons het, is nie groot genoeg om in boerdery- en voedingsbehoeftes van die huishouding te voorsien nie."

Volgens hulle het inwoners meer grond nodig en by het gevolglik 'n beroep op die regering gedoen om die omliggende private grond te koop en dit in hervestigingsplase omskep.

Die nedersetting, wat die tuiste van sowat 1 000 mense is, val onder die Windhoek andelike-kiesafdeling.



Inwoners van Dordabis wil hê die regering moet ingryp. FOTO: JEMIMAH NTSHOLE

Daar is net 'n poskantoor, kliniek, polisie-kantoor en een primêre skool. Leerlinge moet noodgedwonge in koshuise in Windhoek, Rehoboth, Hoachanas, Groot Aub of Leonardville gaan bly om hul skoolopleiding voort te set.

"Ons is uitgeleërd, ons het niks meer as

'n graad 7-rapport nie. Ons wil die kennis en vaardighede hê wat ons lewensgehalte soveel as dié van ons kinders sal verbeter. Ons is moeg daarvoor om net op te staan en niks anders te hê om te doen as om by die kroeg te sit en te gaan slaap wanneer die son sak nie," sê 'n 28-jarige ma van drie wat verlies om anoniem te bly.

"Soms sit jy net by die huis en dink daaraan om jouself dood te maak, want daar is niks om te doen nie. Om te drink, te rook en rond te slaap, niks help nie. Politici kom net met verkiesingstyd hienasies en maak beloftes aan ons. Ons wil resultate sien," het sy gesê. "Sy is bang om in gemeenskapvergaderings op te staan en raadslid te bevestigteken uit vrees vir viktimsiering.

"Wanneer ons iets sê, veral jongmense, voel die raadslidde ons respektieer hulle nie en kritiseer ons daarvoor. Dit is sakkig."

Volgens haar wend baie van die jong meisies in die nedersetting hulle foto's te werk en slaap by maats wat uit Windhoek, Rehoboth en Gobabis die nedersetting kom besoek.

Kommentaar van die Windhoek landelike-kiesafdeling was teen druktyd nie beskikbaar nie.

www.republiekain.com

PUBLIC PARTICIPATION NOTICE

ENVIRONMENTAL SCOPING ASSESSMENT: IRRIGATION BASED AGRICULTURAL ACTIVITIES ON WARD 105 FARM IN ARDABIS, OMBUDHO 851005

Geo Pollution Technologies (Pty) Ltd was appointed by Groot Aub Farming CC, 141 Prinseng, D Bisschoff and Aderwale, Salfra CC, to undertake environmental scoping assessments for irrigation based activities on farms Bontata FMB9670, Porton 1 of Bontata Noord FMB1432, Oudman Oos FMB01278, Groot Bontata FMB01543 and Oud Bontata FMB01233. Additional and location information can be obtained at <http://www.thedordabis.com/irrigation-projects.html>

The environmental assessments will be conducted according to the Environmental Management Act of 2007 and its regulations as published in 2012.

The Proposers invite by means of certain pilot, micro-sprayer systems and drip irrigation. Water for these purposes is abstracted from production boreholes. All relevant operational activities will be included in the assessments.

All interested and affected parties are invited to register with the environmental consultant. By registering you are provided with the opportunity to share any comments, issues or concerns related to the project, for consideration in the environmental assessment. Additional information can be requested from Geo Pollution Technologies. All written comments and concerns should be submitted to Geo Pollution Technologies by 29 March 2023.

Quezette Boman
Geo Pollution Technologies
Tel: +264-61-257411
Fax: +264-88263368
E-Mail: qboman@thetanzab.com

GEO
POLLUTION
TECHNOLOGIES

CHEERS TO

WHISKY WEDNESDAY

22 MARCH 2023

White Horse Blended Scotch Whisky

ONE DAY ONLY

199.99 750ml

LiquorShop

TEAM NAMIBIA

Not for Persons Under the Age of 18. Drink Responsibly.

Site Notice



Appendix F: Consultants' Curriculum Vitae

ENVIRONMENTAL ASSESSMENT PRACTITIONER**Quzette Bosman**

Quzette Bosman has 16 years' experience in the Impact Assessment Industry, working as an Environmental Assessment Practitioner and Social Assessment practitioner mainly as per the National Environmental Legislation sets for South Africa and Namibia. Larger projects have been completed in terms of World Bank and IFC requirements. She studied Environmental Management at the Rand Afrikaans University (RAU) and University of Johannesburg (UJ), including various Energy Technology Courses. This has fuelled a passion towards the Energy and Mining Industry with various projects being undertaken for these industries. Courses in Sociology has further enabled her to specialize in Social Impact Assessments and Public Participation. Social Assessments are conducted according to international best practise and guidelines. Work has been conducted in South Africa, Swaziland and Namibia.

CURRICULUM VITAE QUZETTE BOSMAN

| | | |
|-------------------|---|--|
| Name of Firm | : | Geo Pollution Technologies (Pty) Ltd. |
| Name of Staff | : | QUZETTE BOSMAN |
| Profession | : | Social Impact Assessor / Environmental Assessment Practitioner |
| Years' Experience | : | 16 |
| Nationality | : | South African |
| Position | : | Senior Environmental Consultant |
| Specialisation | : | ESIA & ESMP; SIA |
| Languages | : | Afrikaans – speaking, reading, writing – excellent English – speaking, reading, writing – excellent German –speaking, reading - fair |

| | |
|--------------------------------|----------------------------|
| First Aid Class A | EMTSS, 2017 |
| First Aid LSM | OSH-Med International 2022 |
| Basic Fire Fighting | EMTSS, 2017 |
| Basic Industrial Fire Fighting | OSH-Med International 2022 |

EDUCATION AND PROFESSIONAL STATUS:

| | | | |
|----|----------------------------------|---|----------------------------------|
| BA | Geography & Sociology | : | Rand Afrikaans University, 2003 |
| BA | (Hons.) Environmental Management | : | University of Johannesburg, 2004 |

PROFESSIONAL SOCIETY AFFILIATION:

Namibian Environment and Wildlife Society
International Association of Impact Assessors South Africa (IAIA SA)
Member 2007 - 2012
Mpumalanga Branch Treasurer 2008/2009

OTHER AFFILIATIONS

Mkhondo Catchment Management Forum (DWAF): Chairperson 2008-2010
Mkhondo Water Management Task Team (DWAF): Member 2009

AREAS OF EXPERTISE:

Knowledge and expertise in:

- ◆ environmental impact assessments
- ◆ project management
- ◆ social impact assessment and social management planning
- ◆ community liaison and social monitoring
- ◆ public participation / consultation, social risk management
- ◆ water use licensing
- ◆ environmental auditing and compliance
- ◆ environmental monitoring
- ◆ strategic environmental planning

EMPLOYMENT:

| | | |
|----------------|---|--|
| 2015 - Present | : | Geo Pollution Technologies – Senior Environmental Practitioner |
| 2014-2015 | : | Enviro Dynamics – Senior Environmental Manager |
| 2010 - 2012 | : | GCS – Environmental Manager (Mpumalanga Office Manager) |

2007 - 2009 : KSE-uKhozi - Technical Manager: Environmental
2006 -2007 : SEF – Environmental Manager
2004 - 2005 : Ecosat – Environmental Manager

PUBLICATIONS:

Contract reports : +190
Publications : 1