


Orange-breasted Waxbill |
Amandava subflava




The few records from north-east Namibia, including the Okavango River, comprise the southern tail of the distribution of the subspecies *A. s. niethammeri*, which is abundant in tropical central and western Africa (Clancey 1980). It differs therefore from the subspecies common in central Zimbabwe and northern and eastern South Africa (Colahan 1997). Reporting rates for this subspecies were low, averaging just below 5% (Jarvis *et al.* 2001). It may be overlooked in the moist grasslands in which it occurs because it is a ground-feeder. It breeds in previously used bishop and widow nests, so breeding is also rarely recorded. It occupies an area of 2,300 km², of which 53% lies within protected areas. It could be argued that this species may be a conservation priority because of its rare subspecific status and potential targeting as a cage bird. However, there is no current evidence to support this (Goodwin 1982).

**Orange-winged Pytilia
(Golden-backed Pytilia) |**
Pytilia afra

This sister species to the Green-winged Pytilia (Melba Finch) *P. melba* was not recorded in Namibia during the 24-year SABAP1 atlas period (Tree 1997k). However, its irruptive nature explains the sightings made by Koen (1988) in the Zambezi region of a species that is widespread and frequently recorded. It has also been recorded in a number of places around Katima Mulilo, particularly in thickets along the banks of the Zambezi River, and in piles of cleared vegetation from cultivated fields on the edge of the woodlands (CJ Brown pers. obs.). More recently, it has been seen in the Bwabwata National Park, about 15 km west of the Kwandu River (P Funston pers. comm.) and about five kilometres east of Shamvura in the Kavango




© Mark Paxton



East region (Paxton 2010). Other records come from across the Chobe River in Botswana (reviewed in Tree 1997k). Its absence during the atlas period may arise from confusion with the Green-winged Pytilia, or because of its skulking nature in thick grass and bush in broad-leaved woodland (Tree 1997k, CJ Brown pers. obs.). It appears to be resident in Zimbabwe and is expected to be resident in Namibia, too. Perhaps the best guide to its presence is the occurrence of the much more conspicuous Broad-tailed Paradise-Whydah *Vidua obtusa* that brood parasitises this species (Randall *et al.* 1994). Since the paradise-whydah is associated mainly with the Chobe and lower Zambezi rivers in Namibia, the Orange-winged Pytilia should be looked for in these areas. It is not a conservation priority in Namibia given its wide distribution stretching into north-east Africa.

Brown Firefinch |
Lagonosticta nitidula




A relatively widespread species in north-east Namibia, this bird occupies 14,800 km² of dense thickets adjacent to perennial rivers (Nuttall 1997a). It enters the *Rare and Peripheral* category because in southern Africa it is restricted to the Okavango Delta of Botswana, the Okavango, Kwando, Chobe and Zambezi river systems of Namibia and extreme western Zimbabwe. It is a poorly known species whose density, population size and conservation needs are virtually unknown. It breeds from October to April in Zimbabwe (Dean & Payne 2005); only two nests are known from Namibia, with eggs being laid in February (Brown *et al.* 2015). It is the host of the Village Indigobird (Steelblue Widowfinch) *Vidua chalybeata* (Dean & Payne 2005). Some conservation concern for this resident firefinch is expressed because of the high pressure on wetland margins by local people and their livestock. In the Mahango area of the Bwabwata National Park, where it occurs in the thickest undergrowth about 500 m from the Okavango River (M Paxton pers. obs.), thickets are impacted in areas where large numbers of elephants are resident (P Lane pers. obs.). Surveys of this diminutive species would greatly assist in providing population estimates for different areas in Namibia.



© Eckart Demasius

Bronze Mannikin |
Lonchura cucullata
(*Spermestes cucullatus*)



This small species, widespread throughout sub-Saharan Africa, just touches the Zambezi and eastern Kavango regions. While it is inconspicuous, it occurs in flocks and is unlikely to be mistaken for any other species. It occupies edge habitat often near water over an area of 3,700 km², of which 40% occurs within conservation areas of the Nkasa Rupara (Mamili) National Park and the Mahango area of the Bwabwata National Park (Jarvis *et al.* 2001). This explains its presence along the Okavango, Zambezi and Kwando rivers in Namibia (Nuttall 1997b). It has low reporting rate in Namibia of 6%, despite probably being resident here. This species may be spreading as water points become more common. It is not considered a conservation priority because of its wide range and large abundance, as well as its pest status in parts of West Africa (Nuttall 1997b).

Broad-tailed Paradise-Whydah |
Vidua obtusa

Within southern Africa, this exotic-looking *Vidua* is found predominantly but sparsely in Zimbabwe, and spills into the Zambezi region along the Chobe and Zambezi riverine woodlands, where its reporting rate is a mere 4% (Tree 1997l). It brood parasitises the Orange-winged Pytilia (Golden-backed Pytilia) *Pytilia afra*, an erratically occurring species rarely observed in north-east Namibia (Brown 1990). This may explain the bird's sudden appearance in other parts of southern Africa where it has not been seen before, displaying and probably breeding (Randall *et al.* 1994). Breeding habits in the wild are poorly known, with laying months given as February to April (Tarboton 2011). It occupies an area of 2,100 km², of which 14% occurs within conservation areas of the Nkasa Rupara (Mamili)