

CONTENTS

VOLUME 40 (1) 2007

OSBORNE, T.O. Editorial	1
LUDWIG, D. Annual report by the chairman of the bird club	2
DIEKMANN, M. & J. First record of Ruppell's Vulture for Namibia	4
HAMUKWAYA, F. & CUNNINGHAM. Some notes on the breeding of the Crowned Cormorant, <i>Phalacrocorax coronatus</i> , from Bird Rock, Namibia	6
MIDDENDORFF, G. Gocheaganas Outing on 15 October 2006	9
MIDDENDORFF, G. Brustbandprinie (<i>Prinia flavicans</i>) brütet im Garten	10
WARREN, Y. & KAAPEHI, C. Bird observations at farm Christirina in the Kalahari-bushveld region	12
TREE, A.J. & BOORMAN, M. Luderitz colour-ringed Hartlaub's Gull at Walvis Bay	24
BRIDGEFORD, P. & HEINRICH, D. Raptor electrocutions and collisions with power-lines in Namibia	25
FRIEDERICH, T. & G. Wetland birds on Tsutsab Vlei 2005/2006	26
FRIEDERICH, T. & G. Birding weekend on farm Tiefland	29
KOLBERG, H. Summary of the 2005 ringing season in Namibia	35
BIRD OBSERVATIONS AND NOTES	45
Request for assistance	48

Editorial

The trials of the Namibia Bird Club continue with it mainly being a Windhoek orientated organization. The number of articles submitted to its journal the *Lanioturdus* has withered to almost unacceptable levels. In 2006 only one issue of the *Lanioturdus* was published and here we are half way through 2007 and finally the editor has enough articles to publish what probably will be the only issue of 2007.

There is obviously an interest in birding within Namibia since there are now many subgroups specializing in aspects of birds. There is Raptors Namibia who publish an on line newsletter once a month. The Crane Working Group, Raptor Road Counts, Rare and Endangered Species Trust working on vultures, and the Wetland Bird Counts which enumerate all the wetlands twice per year.

I know that there are birders observing birds because of the rare birds which have been reported to the magazine *Birds and Birding* from Namibia.

Once again I appeal to all members to help the Namibia Bird Club. If you want to keep the club viable you must also do your part and try and get new members to join. We are currently down to fewer than 100 members.

Fortunately there is good news and there has been a recent increase in the number of articles submitted and hopefully this trend will continue and we could see another issue published this year.

I hope readers will note the addition to the Namibia list of Ruppell's Vulture. There are probably more juvenile Ruppell's Vultures visiting Namibia than we realize. Observers should keep their eyes open when watching vultures at a kill, you may be able to add a new species to your list. If you do see one of these birds please try to photograph it.

Brustbandprinie (*Prinia flavigans*) brütet im Garten

Gudrun Middendorff

Im vergangenen Sommer bemerkte ich in meinem Garten in Windhoek, Stadtteil Olympia, in einem Schwarzdorn-Silberbusch (*Catophractes alexandri*) ein Nest. Es wurde feinstes Grasmaterial verwendet und es war deutlich zu sehen, dass dort noch fleißig gebaut wurde. Bald war klar, dass ein Pärchen Brustbandprinien die Bauunternehmer dieses Nestes waren.

Über eine Zeitspanne von 10 Tagen konnte ich beobachten wie das Nest ausgebaut und schließlich mit wolligen Pflanzenfusseln gut gepolstert wurde. Bald saßen die Vögel abwechselnd im Nest und brüteten. Es regnete fast jeden Tag und das Nest schrumpfte etwas zusammen. Als eines Tages der Altvogel das Nest verließ, habe ich schnell nachgesehen und 3 Eier gezählt. Am nächsten Tag waren es vier Eier. Nach etwa 13-14 Tagen fing ein Elternteil an Futter zum Nest zu bringen.

Am Anfang hat nur ein Altvogel gefüttert, aber bald waren beide voll damit beschäftigt Motten, Würmer und andere Insekten ins Nest zu schleppen. Man konnte die Köpfe der Jungvögel bald sehen, und wenn die Alten auftauchten, rissen sie ihre Schnäbel weit auf.

Allmählich wurden die Raupen und Falter immer größer und fetter und die Alten stopften und stopften. Bald war für den Altvogel nachts auch kein Platz mehr im Nest und die Jungen wärmen sich gegenseitig. Nach nochmals etwa 13-14 Tagen saßen die Jungvögel im Busch außerhalb des Nestes und mit viel Locken und Vorfliegen übten die Jungen sich im Fliegen.

Die Vogelfamilie blieb noch drei Tage in meinem Garten, hauptsächlich in den Stachelbeersträuchern. Dort war es geschützt und kühl und es gab eine große Auswahl an Insekten zu jagen.

Etwa 6 Wochen waren seit dem Anfang des Nestbaus vergangen und die Prinienfamilie zog aus dem Garten ins offene Feld.

Summary

During the last rainy season I was able to observe a pair of nesting in a rattle tree (*Catophractes alexandri*) and raising four chicks in my garden in Windhoek. Building a nest and laying eggs took about two weeks, about 13-14 days incubation and roughly the same time was necessary for rearing the offspring till adults and fledglings left the garden.



Brustbandprinie/blackcheasted prinia (*Prinia flavigans*)

Foto/Photo: Neil Thomson