
Climate and Near-surface Airflow Over the Central Namib

J. A. Lindesay & P. D. Tyson

Climatology Research Group, Department of Geography, University of the Witwatersrand, Johannesburg, 2050 South Africa

Research on the climatology of the central Namib Desert on the west coast of southern Africa is reviewed, and diurnal and seasonal oscillations of the atmospheric boundary layer over the region are examined in detail. Both the vertical and horizontal structures of the thermo-topographic airflow compare well with similar wind systems occurring elsewhere. Thermo-topographic airflows over the central Namib are found to have a regional significance frequently equalling or exceeding that of the general circulation. The strength, depth and unusually clearly defined diurnal and seasonal oscillations of these winds render the central Namib a unique area for the study of boundary-layer oscillations.

INTRODUCTION

The Namib Desert on the west coast of southern Africa is a complex environment in which a unique ecology has evolved in response to the environmental parameters characterizing the region. An important controlling factor in the evolution of the desert biota and of the present-day physical features of the Namib, as in any desert, is the climate of the area (e.g., Hadley and Szarek, 1981; Crawford and Gosz, 1982). Early climatological studies of the region were confined to the more readily accessible coastal zone (Taljaard and Schumann, 1940; Jackson, 1942), but over the last three decades the recording of climatic parameters has been extended inland, particularly over the central Namib (Fig. 1). Nevertheless, few studies of central Namib climate exist: exceptions include the analyses of single-station weather records by Schulze (1969) and Seely and Stuart (1976), and the more comprehensive work on data from several central Namib stations by Lancaster, Lancaster and Seely (1984).

Of the various climatic parameters available for consideration, moisture variables and wind over the central Namib have received the greatest amount of scientific attention. Two sources of moisture sustain life in the Namib: rainfall and fog. Rainfall over the central Namib occurs mainly in the form of convective summer storms (Sharon, 1981) from which maximum precipitation is received over the Escarpment to the east (Fig. 1), while fog-water precipitation is the dominant moisture source over the western parts of the Desert adjacent to and inland from the coast (Lancaster *et al.*, 1984). Temporal and spatial characteristics of both rainfall (Dyer and Marker, 1978; Nieman, Heyns and Seely, 1978; Gamble, 1980; Sharon, 1981; Pietruszka and Seely, 1985) and fog-water precipitation (Nagel, 1962; Pietruszka and Seely, 1985) have been investigated, but such studies have been limited by the paucity of long-term, continuous and reliable records for the area.

Rather than forming the focus for analysis, central Namib climatic parameters have more frequently been incorporated into studies focused on geological/geomorphological (e.g., Besler, 1972; Marker, 1977; Hattle, 1985; Wilkinson, 1987) or

ecological aspects of the area (e.g., Seely, 1979a; Seely and Louw, 1980). The role of rainfall in the ecology of the central Namib has been investigated (Seely and Louw, 1980), and the importance of fog-water precipitation to vegetation on the gravel plains in the north of the area (Bornman, Botha and Nash, 1973) and on the dunes to the south (Seely, De Vos and Louw, 1977; Louw and Seely, 1980), as well as to the dune fauna (Louw, 1972; Hamilton and Seely, 1976; Seely and Hamilton, 1976; Seely, 1979b; Seely, Lewis, O'Brien and Suttle, 1983), has been shown. Analyses of surface winds over the central Namib have provided an important input to geomorphic studies of the Namib dune field (Ward and von Brunn, 1985), where present and past prevailing winds and seasonal variations in wind directions have been incorporated into studies of dune orientation and movement (e.g., Besler, 1980; Harmse, 1982; Lancaster, 1983, 1985; Ward and von Brunn, 1985; Wilkinson, 1987). Some of the most comprehensive climatological work undertaken in the central Namib is the analysis of surface (Tyson and Seely, 1980) and lower-atmospheric boundary-layer (Lindesay and Tyson, 1990) airflow characteristics over the region, and it is the recent work on the vertical and spatial characteristics of the lower boundary layer over the central Namib that will be considered in this paper.

Local climates of coastal and mountain areas are characterized by the modification of synoptic-scale airflow by thermally and topographically induced diurnal oscillations of the lower boundary layer. Numerous observational studies of the sea/land breezes, local mountain/valley winds and regional mountain/plain winds which constitute these oscillations (Defant, 1958; Flohn, 1969; Atkinson, 1981; Yoshino, 1981; Sturman, 1987) and numerical modelling of these systems (see the reviews of Atkinson, 1981, 1983, and Pielke, 1984) have led to an improved theoretical understanding of individual thermo-topographic wind systems. The complex wind regimes produced by the interactions of topographic and thermal effects on local and regional scales with each other and with the synoptic circulation, however, are not as well documented. Empirical studies in which interactions among various thermo-topographic boundary-layer wind systems and larger-scale




Fig. 1

a: Location map of the central Namib Desert, showing major physical features, with transect line (Figs 6 & 7) shaded. **b:** Section through Southern Africa from northwest to southeast, showing the two coastal margin areas where thermo-topographic winds are known to develop.

airflows are described have been undertaken for areas such as Indonesia (van Bemmelen, 1922), California (Edinger and Kao, 1959; Frenzel, 1962), Israel (Skibin and Hod, 1979; Bitan, 1981; Goldreich, Druyan and Berger, 1986), and the Canterbury Plains of New Zealand (Sturman and Tyson, 1981; McKendry, 1983; McKendry, Sturman and Owens, 1986). Such studies are largely based on surface wind information augmented by limited vertical observations.

In southern Africa the nature and interactions of local and regional thermo-topographic winds over the eastern plateau slopes of Natal have been thoroughly considered in this way (Tyson, 1966, 1967, 1968a, b; Preston-Whyte, 1969, 1974), but local wind regimes over the central Namib Desert are not as well described and understood. Nevertheless, Goldreich and Tyson (1988) have shown that diurnal boundary-layer processes dominate the near-surface windfield there in all seasons. Characteristics of the sea breeze have been documented for this coast (Jackson, 1942, 1954); further inland topographically and thermally induced air movements have been identified only from surface data (Tyson and Seely, 1980). In this paper, diurnal and seasonal oscillations of the boundary layer over the desert of the central Namib will be examined. Both the vertical and horizontal structure of the thermo-topographic airflow will be considered and interactions among the various mesoscale and synoptic wind systems will be discussed.

PHYSICAL SETTING AND GENERAL AIRFLOW CLIMATOLOGY

The hyper-arid central Namib Desert is bounded by the cold Atlantic Ocean to the west and a dissected western plateau slope beneath the Escarpment (which has an average altitude

of 1500 m, rising to 2300 m in places) some 160 to 180 km to the east (Fig. 1a). The deeply incised Kuiseb River valley forms a boundary between the relatively flat gravel plains to the north and the coast-parallel linear dunes of the sand sea to the south. A research station is situated in the Kuiseb valley at Gobabeb (Fig. 1a), which is 56 km inland from the coast and is where the majority of the measurements included in this paper were made. The northwest-southeast trend of the Kuiseb valley, from near the coast to some distance beyond Gobabeb, changes to northeast-southwest nearer to the Escarpment (Fig. 1a).

Physical contrasts between the western and eastern plateau slopes of southern Africa are great (Fig. 1b). The cold waters of the Benguela Current and sparsely vegetated desert on the west coast contrast markedly with the warm waters of the Agulhas Current and well-vegetated coastal margins of Natal, and render the land-sea thermal contrast more distinct for the west than for the east coast. Nevertheless, well-developed thermo-topographic boundary layer oscillations occur over Natal. Although topographic influences on airflow over the west coast are expected to be less marked than those for the east coast, thermally induced boundary-layer airflow should develop more strongly over the central Namib owing to the larger land-sea temperature contrast along the west coast. It remains to be seen to what extent a similar model of local and regional diurnal airflow to that proposed for Natal (Preston-Whyte and Tyson, 1988) will describe conditions in the central Namib.

Surface boundary-layer airflow over the central Namib is distinctive (Tyson and Seely, 1980) (Fig. 2). Daytime south-westerly sea breezes and northwesterly valley and plain-mountain winds dominate the near-surface circulation in summer, when calms are least frequent and regional pressure




Fig. 2

Schematic models of the local components of airflow over the central Namib Desert to show the occurrence of sea breezes, valley winds and plain-mountain winds by day and in summer, and land breezes, mountain winds and mountain-plain winds by night in winter (after Tyson and Seely, 1980).

gradients reinforce the strong land-sea thermal contrast (Fig. 3a–c). Nocturnal northeasterly land breezes and southeasterly mountain and mountain-plain winds associated with cool air drainage occur preferentially in winter. Distinctive changes in nocturnal airflow patterns are evident at Gobabeb between summer and winter (Fig. 3d). Inter-seasonal variations are less obvious during the afternoon-evening period (12h00 to 21h00) when surface heating effects are greatest. At times synoptic disturbances may completely disrupt the diurnal rhythm of boundary-layer airflow over the Namib, the most common such disturbance being the easterly to northeasterly Berg winds resulting from strong pressure gradients normal to the coast. Such disturbances are infrequent, however, and the surface climate of the central Namib is dominated by near-surface thermally and topographically induced boundary-layer oscillations. A more detailed analysis of surface winds at Gobabeb confirms these findings (Lancaster *et al.*, 1984).

THE METHODS

Hourly pilot balloon (pibal) ascents were made from Gobabeb in the central Namib (Fig. 1a) during each of the months July 1986 and January 1988. On several suitable occasions hourly ascents were also made from Rooibank, in the Kuiseb valley 22 km from the coast, and at Zebra Pan, on the gravel plains northeast of Gobabeb and 100 km from the sea (Fig. 1a). Simultaneous measurements enabled the construction of ver-

tical sections of the boundary layer along a transect some 100 km long between the coast and the Escarpment, while continuous hourly pibal measurements at Gobabeb allowed the detailed investigation of the diurnal rhythms and interactions within the boundary layer over a point representative of much of the central Namib area. The programme of field observations provided more than 460 hours of soundings during periods of generally light gradient flow.

Single-theodolite tracking of balloons was used, with cognizance being taken of the errors to which such tracking is susceptible (Reynolds, 1966; Boatman, 1974). Since no appropriate radiosonde data were available against which to verify the pibal data, double-theodolite tracking was used on several occasions during the summer period when the effects of surface heating on lapse rates were greater. Errors in heights and wind speeds were no greater than those found in similar studies elsewhere (e.g., Sturman and Tyson, 1981).

THE OBSERVATIONS

Plain-mountain winds

Surface anemometer observations at Gobabeb show clearly that northwesterly plain-mountain winds predominate in summer (Fig. 3c) and are least frequent in winter. The plain-mountain winds appear to be purely antitriptic winds resulting from the thermal gradient established between the gravel plains to the north and the Escarpment to the east. During summer,

when surface heating effects are at a maximum, the plain-mountain wind oscillates in depth and strength day after day, with little or no disturbance (Fig. 4a, lower). At Gobabeb maximum depths of 1000 m to 1600 m and speeds of 10–15 ms^{-1} occur between 16h00 and 18h00; minimum depths and speeds occur around and after midnight. Although the near-surface flow may be disturbed and may reverse around sunrise (with the occurrence of the reverse-direction mountain-plain wind), at levels near the top of the boundary layer the inland airflow may continue uninterrupted for long periods.

In winter, when nocturnal cooling effects are strongest, the inland-directed plain-mountain wind is a daytime phenomenon only and is seldom as deep as its summertime counterpart (Fig. 4a, upper). The contrast between the summer and winter examples of the plain-mountain wind over the central Namib is striking. In summer this wind is a constant determinant of the structure of the boundary layer; in winter the near-surface airflow is much more disturbed and many more local winds develop due to the clear diurnal reversals in low-level temperature gradients between the Escarpment and the coast and within the valleys of the western plateau slopes.

Mountain-plain winds

Just as the boundary-layer structure is dominated by plain-mountain winds in summer, so the winter pattern is characterized by regularly occurring nocturnal southeasterly mountain-plain winds blowing from the Escarpment zone towards the coast (Fig. 4b, upper). These winds develop at Gobabeb around 22h00, reach a maximum depth of about 1000 m and strength of 5–10 ms^{-1} after sunrise, and then decay rapidly between 10h00 and 12h00 the next day. This pattern is repeated day after day unless disturbed by synoptic-scale weather perturbations such as Berg winds, coastal lows or cold fronts. At times, when synoptic disturbances are entirely absent, the stable mountain-plain wind may develop a clear return current in a closed circulation seldom more than 1000 m deep. In summer, mountain-plain winds still occur, but only for a few hours at a time and seldom to a depth of even 500 m (Fig. 4b, lower). It is surprising that they occur at all.

Sea breezes

Whereas the plain-mountain and mountain-plain winds over





Fig. 3

a: Schematic diagram of the three major thermo-topographic winds prevailing in the near-surface circulation over the central Namib Desert. **b:** Mean January and July surface wind-roses for Gobabeb for the period 1976 to 1981 (after Lancaster *et al.*, 1984). **c:** Monthly variation of wind direction frequencies at Gobabeb. **d:** January and July three-hourly surface wind vectors for Gobabeb for the period 1976 to 1981 (adapted after Lancaster *et al.*, 1984).

the central Namib are characterized by marked inter-seasonal differences in occurrence, sea breezes occur regularly throughout the year, with a tendency to show a semi-annual cycle with peaks around the equinoxes (Fig. 3c). The characteristic direction of airflow in the sea breeze in the region is southwesterly, owing to turning of the breeze by the Coriolis effect (Jackson, 1954). On the Namib coast at Walvis Bay the sea breeze is established between 09h00 and 12h00 on most

**Fig. 4**

Time-height sections of hourly winds at Gobabeb for 72-hour periods in winter (upper diagram of each pair) and summer (lower diagram). Plain-mountain winds are shaded in **a**, mountain-plain winds in **b**, and sea breezes in **c**. Flags fly with the wind: one feather represents wind speeds of 2,5–4,9 ms^{-1} , two feathers 5,0–9,9 ms^{-1} , three feathers 10,0–14,9 ms^{-1} .

mornings, attains maximum depths of about 1000 m and then decays until, after sunset, southwesterly winds are infrequent at the coast (Jackson, 1942, 1954). There is little seasonal variation in sea breeze frequency at Walvis Bay, although Jackson (1954) suggests that the winter breeze is shallower than its summer counterpart. In winter the sea breeze penetrates inland to reach Gobabeb around 14h00, ceases at the ground at about 20h00, when it is undercut by the developing mountain-plain wind, and continues blowing at heights of

500–750 m until around or after midnight (Fig. 4c, upper). The system attains its maximum depth over the central Namib (seldom in excess of 1000 m) just before sunset. At the Gamsberg, 170 km inland, the sea breeze reaches the Escarpment at about 18h00 and only lasts an hour or two before ceasing altogether.

In all the winter cases the evening sea breeze at Gobabeb was undercut by the mountain-plain wind. In summer, when the breeze is not as strong as its winter counterpart and has a




Fig. 5

Vertical profiles of the zonal component (westerlies positive) of various thermo-topographic local winds at selected stations in the central Namib Desert.

symmetrical diurnal pattern of growth and decay (Fig. 4c, lower), this undercutting does not occur. The summer sea breeze may exceed 1000 m in depth just before sunset, owing to the enhanced instability of the boundary layer. In summer the sea breeze system over the central Namib decays from the top downward; in winter from the ground upward.

Wind profiles

Velocity profiles of the thermo-topographic winds induced by the thermal gradients and topography of the central Namib, and by the western and eastern boundary conditions imposed by the Benguela Current and the Escarpment respectively, evidence a surprising regularity and symmetry (Fig. 5). Return currents are difficult to measure, although on occasion such currents may be clear (as in Fig. 5a). Jackson (1954) reports return currents several times the depth of the sea breeze system at Walvis Bay, as does Atkinson (1981) for other parts of the world. At Gobabeb the return currents that were observed were always less deep than the sea breezes below them. The nocturnal land breeze is considerably weaker and shallower than its daytime sea breeze counterpart (Fig. 5b).

Plain-mountain winds in summer are deep and strong, and velocity maxima in excess of 10 ms^{-1} at heights of about 500 m are common (Fig. 5c). The wintertime nocturnal mountain-plain winds (Fig. 5d) are considerably weaker (with speeds of up to 3 ms^{-1}), but may reach similar depths to the plain-mountain winds. The zonal velocity maxima in both cases are attained at between 250 m and 400 m, that is approximately one quarter to one third the total depth of the system, although

the maximum is less well defined for the mountain-plain than for the plain-mountain wind case. The sea breeze reaches a maximum velocity normal to the coast at a height that is likewise one quarter to one third the depth of the system.

Airflow between the Escarpment and the coast

The vertical structure and temporal variation of the thermo-topographic boundary-layer airflow over the central Namib have been described. Spatial continuity of this structure and of the oscillations of the boundary layer over time may be determined along transects on an approximately northwest-southeast section from seaward of Rooibank near the coast to east of Zebra Pan some 100 km inland (Fig. 1a). Examples of these transects will be considered.

In summer a typical sequence of the decay of the plain-mountain wind and sea breeze, followed by the onset of the oppositely-directed mountain-plain wind, the decay of this wind and the reestablishment of the plain-mountain wind the following day, is given in Fig. 6. A 1500 m deep late-afternoon plain-mountain wind overlain by a sea breeze some 500 m deep decayed rapidly after sunset, so that by 21h00 it was a skin of air with a depth of approximately 100 m moving slowly inland. From 23h00 until after 01h00 the inland penetration of the plain-mountain flow was prevented by drainage of cool air from the interior in the mountain-plain wind. At times this seaward drift of cool air was obliterated by a surge of warmer air moving inland; at times land breezes developed temporarily on the coast. At 06h00 a thin layer of mountain-plain air draped the entire desert. Two hours later a plain-mountain wind had developed near the coast and penetrated inland beyond Gobabeb. By 10h00 the plain-mountain wind had advanced halfway toward the Escarpment.

A typical winter sequence shows the development of the same local winds as in summer, but with very different emphases (Fig. 7). At 16h00 a strong sea breeze (with a clear and spatially-continuous return current) was displacing a decaying plain-mountain wind as it moved inland toward the Escarpment. The sea breeze began weakening first at the coast; this weakening then progressed inland. By 18h00, and for the following two hours, the sea breeze dominated the low-level airflow over the whole sea-to-Escarpment transect. By 22h00 a complete reversal had occurred and thereafter throughout the night drainage of cool air seaward dominated the local airflow regime to a depth of between 300 m and 500 m. At all times the local flow was most prone to disturbance by synoptically-driven winds the further observations were made away from the Kuiseb River valley. Between 09h00 and 11h00 the following day the mountain-plain flow reversed abruptly to a plain-mountain wind, which then advanced inland as it strengthened throughout the day.

DISCUSSION

The role of interacting thermo-topographic airflows in establishing the characteristics of the boundary layer over both the western (Tyson and Seely, 1980) and eastern (Preston-Whyte and Tyson, 1988) coastal margins of southern Africa has long been recognized. Over the central Namib thermo-topographically induced boundary-layer oscillations clearly dominate the diurnal and seasonal variations of local and regional airflow,




Fig. 6

Time sequence of transects showing interactions of summertime thermo-topographic winds over the central Namib Desert between the coast and the Escarpment. Sea breezes are stippled; mountain-plain and plain-mountain winds are shaded.

confirming the identification of this area of southern Africa as one with particularly high diurnal variability of the near-surface windfield (Goldreich and Tyson, 1988). Disturbance of thermo-topographic airflow in the region by strong synoptic-scale winds is relatively rare (Tyson and Seely, 1980; Lancaster *et al.*, 1984) and the ratio of local to general winds gives it one of the highest mesoscale wind indices for the subcontinent (Goldreich and Tyson, 1988).

The particular characteristics of thermo-topographic regional airflow over the central Namib are the result of seasonally varying interactions among three major boundary-layer wind systems: the sea/land breeze response to the cold ocean/hot desert interface; the valley/mountain winds within the Kuiseb and similar river valleys dissecting the coastal plain; and the plain-mountain and mountain-plain winds resulting from the parallel existence of the desert plains and the Escarpment (Tyson and Seely, 1980).

During undisturbed summer periods when surface heating effects over the central Namib are strongest, the warm, unstable northwesterly plain-mountain wind dominates the airflow within the boundary layer, both by day and by night, for days at a time. Its rhythm is broken only by short-lived punctuations near the ground by sea breezes and mountain-plain winds. By contrast, the shallower east-coast plain-mountain winds, which reach depths of about 1200 m (Tyson, 1966, 1968b;

Tyson and Preston-Whyte, 1972; Preston-Whyte and Tyson, 1988) compared with Namib plain-mountain winds more than 1500 m deep, are commonly replaced by nocturnal mountain-plain flow after midnight in both summer and winter.

The occurrence of cool, stable southeasterly mountain-plain winds over the central Namib is almost entirely limited to winter nights, when cooling effects are best developed. As is the case over Natal, west-coast mountain-plain winds are weaker and shallower than the plain-mountain winds and develop after sunset. The winter mountain-plain wind over the central Namib, however, persists for several hours longer than that over Natal (Tyson, 1968b; Tyson and Preston-Whyte, 1972; Preston-Whyte and Tyson, 1988). It may blow from an hour before sunset to late morning of the following day, is reasonably constant in depth, and evidences surging of the kind reported over Natal (Tyson, 1968c). Both northwesterly plain-mountain and southeasterly mountain-plain winds seldom bear any significant relationship to the near-surface pressure gradients of the general circulation. In their preferred seasons of development diurnal valley and plain-mountain winds and nocturnal mountain and mountain-plain airflows over the western desert coastal margins of southern Africa are better developed than similar regional airflows reported for the European Alps (Burger and Ekhart, 1937), the Rockies of North America (Hawkes, 1945), parts of Mexico (Lauer and Klaus, 1975), the




Fig. 7

Time sequence of transects showing interactions of wintertime thermo-topographic winds over the central Namib Desert between the coast and the Escarpment. Sea breezes and return currents are stippled; mountain-plain and plain-mountain winds are shaded.

Southern Alps of New Zealand (Sturman, Fitzsimons and Holland, 1985) and the coastal and adjacent inland areas of Israel (Goldreich *et al.*, 1986).

Disruption of predominant plain-mountain winds over the central Namib by sea breezes occurs briefly on most summer afternoons. During winter there is a clear diurnal oscillation of the boundary layer set up by the interaction between well-developed nocturnal mountain-plain winds, short-lived plain-mountain winds around midday, and the mid- to late-afternoon occurrence of the sea breeze. Sea breezes at Walvis Bay have been reported to occur with approximately equal frequency throughout the year, but to be most strongly developed during summer (Jackson, 1942). The latest sets of data for the central Namib show strongest sea breezes in winter, a discrepancy attributable to the higher frequency of occurrence and dampening effect of middle- and high-level cloud on local wind development over the central Namib in summer. Sea breezes, like all thermo-topographic winds, are best developed with completely clear skies. The suppressive effects of clouds on sea breezes have recently been documented (Segal, Purdom, Song, Pielke and Mahrer, 1986) and may be considerable, although in New Zealand sea breezes will develop in the presence of cloud (Sturman and Tyson, 1981).

The sea breeze is strong and deep over the Namib coast and the characteristics of the breeze are in agreement with

results obtained in both low latitudes (van Bemmelen, 1922; Roy, 1940; Dekate, 1968) and middle latitudes (Frizzola and Fisher, 1963; Camuffo, Tampieri and Zamboni, 1979; Sturman and Tyson, 1981; Mathews, 1982; McKendry *et al.*, 1986; Prezerakos, 1986). After setting in at the coast at about 09h00, the Namib sea breeze backs to southwest through the day under the influence of the Coriolis force. Similar turning of the sea breeze has been reported for other areas (Haurwitz, 1947; Estoque, 1961, 1962; Yan and Anthes, 1987). By mid-afternoon the sea breeze is southwesterly throughout the boundary layer from the coast to beyond Gobabeb, and by sunset the sea breeze has penetrated almost to the Escarpment. Weakening of the sea breeze at the coast around sunset coincides with maximum southwesterly flow at Gobabeb, is followed several hours later by a sea breeze maximum further inland and then by penetration of the wind to the Gamsberg on the plateau an hour or so before midnight. The sea breeze dominates boundary-layer airflow over the entire central Namib between the coast and the Escarpment for much of the afternoon and early evening period, giving way around midnight at Gobabeb to relatively stable mountain-plain flow advancing seaward from the Escarpment, although the breeze has ceased at the coast several hours before. Unlike its plain-mountain and mountain-plain counterparts the sea breeze may on occasion be associated with a clear return current.

In contrast to the deep, unstable afternoon sea breeze circulation, the nocturnal land breeze of the coastal littoral is a shallow, short-lived system seldom deeper than 300 m. Onset of land-breeze flow on the southern African west coast after 01h00 and cessation of the breeze before 09h00 accords with observations of land breezes in India (Dekate, 1968; Aggarwal, Singal, Kapoor and Adiga, 1980) and North America (Meyer, 1971; Atkinson, 1981) and with theoretical studies (Rotunno, 1983).

The vertical structures of west-coast boundary-layer airflow across the central Namib are similar to those of east-coast Natal thermo-topographic winds (Preston-Whyte and Tyson, 1988). Lagrangian profiles of the wind component normal to each of the coasts show that the maximum velocities of land and sea breezes, plain-mountain and mountain-plain winds are comparable for both regions, although west-coast airflow systems are deeper, particularly in summer. Over the west coast, as in Natal (Tyson, 1966, 1968a-c; Tyson and Preston-Whyte, 1972; Preston-Whyte and Tyson, 1988) maximum-velocity airflow occurs at 250–300 m above the surface. The variation of wind speed with height may be modelled using a parabolic profile in which the height of maximum velocity is half the height of the system (Davidson and Rao, 1963), or using a Prandtl profile in which the height of maximum velocity is one quarter the height of the system (Defant, 1958; Atkinson, 1983; Pielke, 1984). The parabolic profile gives a good representation of mountain, mountain-plain and plain-mountain winds in the Drakensberg region and of the Natal-coast sea breeze (Preston-Whyte and Tyson, 1988), whereas the Prandtl profile best fits mountain winds in the Natal interior (Tyson, 1968c) and Drakensberg valley winds. Central Namib wind profiles of plain-mountain and mountain-plain winds and of the sea breeze, by contrast, conform closely to the Prandtl model. The plain-mountain wind maximum occurs at a height of about 300 m in a system some 1200 m deep, and the sea breeze maximum at 250 m in a 1000 m-deep system. The west-coast land breeze, however, exhibits a parabolic profile, with maximum wind speeds at approximately half the depth of the system.

ACKNOWLEDGEMENTS

The staff of the Desert Ecological Research Unit of Namibia at Gobabeb, particularly Dr M. K. Seely, provided invaluable facilities, assistance and hospitality. Their help is gratefully acknowledged. The study would not have been possible without the assistance of staff and students of the University of the Witwatersrand. Our thanks go to Dr M. J. Wilkinson and Mrs C. H. Vogel for helping to organize and run the fieldtrips. Kathy Barbfiera, Mario Barbfiera, Brynn Bayman, Brenda Beverly, Kim Christie, Lynne Collie, Sue Crouch, Caryn Eccles, Gavin Elliot-Wilson, Gordon Fenwick, Diane Geer, Jenny Gobey, Michael Haynes, Yvonne Hong, Greg Knill, Jerry Lengoasa,

CONCLUSIONS

Surface winds over the central Namib Desert have received more attention than most other climatic parameters for the region. The analysis of winds has now been extended to include the whole boundary layer, and it has been shown that the nature of the thermo-topographic forcing of boundary-layer airflow over the central Namib is similar to that occurring on the east coast of southern Africa. The strong thermal gradients across the central Namib Desert, bounded to the west by the consistently cold ocean and to the east by the Escarpment, strongly heated by day and similarly cooled by night, together with the pronounced diurnal heating and cooling of the desert itself, are responsible for the strength and longevity of the boundary-layer airflows that develop over the region. Throughout the year the boundary layer varies in depth from under 500 m at night to over 1000 m by day as a pulsating mass of air moves alternately inland and thereafter seaward in an oscillation as distinctive as that reported anywhere. The seasonally clearest suite of successional winds is encountered in winter when the plain-mountain winds are weak and short-lived and the boundary layer is dominated by mountain-plain winds by night and sea breezes during the post-noon period. In summer, northwesterly plain-mountain winds are present day and night in a remarkably persistent response to thermal and topographic forcing. These distinctive boundary-layer oscillations are disrupted only infrequently by strong synoptic-scale disturbances. Surface characteristics and vertical structure of the lower atmosphere over the central Namib confirm that, in the region between the coast and the inland plateau, boundary-layer airflow is controlled by surface thermal effects and by both local and regional topography. Thermo-topographic airflows in the region frequently have a regional significance equalling or exceeding that of the general circulation. By virtue of the strength, depth and unusually clearly defined diurnal and seasonal oscillations of the thermo-topographic airflows of the region, the central Namib Desert constitutes a unique laboratory for the study of boundary-layer oscillations.

Paul Maclear, Christine Muller, Charles Nepgen, Sean O'Beirne, Michelle Pearse, Colin Pilkington, Mike Slattery, Belinda Theron, Greg Theron, Jenny Tyson, Anja van Nierop and Ingo Vogt assisted with data collection. They constituted a marvellous team, and without their assistance the study could not have proceeded. Mr P. Stickler drew the diagrams. The work forming the bulk of this paper was presented at the 26th International Geographical Congress of the International Geographical Union, Sydney, 21–26 August 1988. The research was funded in part through the Foundation for Research Development of the C.S.I.R.

REFERENCES

- AGGARWAL, S. K., SINGAL, S. P., KAPOOR, R. K. and ADIGA, B. B., 1980. A study of atmospheric structures using sodar in relation to land and sea breezes. *Boundary Layer Meteorology* 18: 361–371.
- ATKINSON, B. W., 1981. *Meso-scale atmospheric circulations*. Academic Press, London.
- ATKINSON, B. W., 1983. Numerical modelling of thermally-driven,

- mesoscale airflows involving the planetary boundary layer. *Progress in Physical Geography* 7: 177–209.
- BESLER, H., 1972. Klimaverhältnisse und Klimageomorphologische Zonierung der Zentralen Namib (Südwestafrika). *Stuttgarter Geomorphische Studien* 83: 1–209.
- BESLER, H., 1980. Die Dünen Namib: Entstehung und Dynamik eines Ergs. *Stuttgarter Geomorphische Studien* 96: 1–241.
- BITAN, A., 1981. Lake Kinneret (Sea of Galilee) and its exceptional wind system. *Boundary Layer Meteorology* 21: 477–487.
- BOATMAN, J. F., 1974. The effect of tropospheric temperature lapse rates on the ascent rates of pilot balloons. *Journal of Applied Meteorology* 13: 955–961.
- BORNMAN, C. H., BOTHA, C. E. J. and NASH, J., 1973. *Welwitschia mirabilis*: observations on movement of water and assimilates under föhn and fog conditions. *Madoqua* 2: 25–31.
- BURGER, A. and EKHART, E., 1937. Über die tägliche Zirkulation der Atmosphäre im Bereich der Alpen. *Beiträge der Geophysik* 49: 341–367.
- CAMUFFO, D., TAMPIERI, F. and ZAMBON, G., 1979. Local meso-scale circulation over Venice as a result of the mountain-sea interaction. *Boundary Layer Meteorology* 16: 83–92.
- CRAWFORD, C. S. and GOSZ, J. R., 1982. Desert ecosystems: their resources in space and time. *Environmental Conservation* 9: 181–195.
- DAVIDSON, B. and RAO, P. K., 1963. Experimental studies of the valley-plain wind. *International Journal of Air and Water Pollution* 7: 907–921.
- DEFANT, F., 1958. Local wind systems. *Compendium of meteorology*, pp. 662–672. American Meteorological Society, Boston.
- DEKATE, M. V., 1968. Climatological study of sea and land breezes over Bombay. *Indian Journal of Meteorology and Geophysics* 19: 421–442.
- DYER, T. G. J. and MARKER, M. E., 1978. On the variation of rainfall over South West Africa. *South African Geographical Journal* 60: 144–149.
- EDINGER, J. G. and KAO, S. K., 1959. The influence of terrain and thermal stratification in the surface and planetary boundary layers. *Final Report, AF 19(604)-2424, AFCRC-TR*, pp. 59–401, UCLA Department of Meteorology, 59–401.
- ESTOQUE, M. A., 1961. A theoretical investigation of the sea breeze. *Quarterly Journal of the Royal Meteorological Society* 87: 136–146.
- ESTOQUE, M. A., 1962. The sea breeze as a function of the prevailing synoptic situation. *Journal of the Atmospheric Sciences* 19: 244–250.
- FLOHN, H., 1969. Local wind systems. *World survey of climatology* Vol. 2, pp. 139–171.
- FRENZEL, C. W., 1962. Diurnal wind variations in central California. *Journal of Applied Meteorology* 1: 405–412.
- FRIZZOLA, J. A. and FISHER, E. L., 1963. A series of sea-breeze observations in the New York City area. *Journal of Applied Meteorology* 2: 722–739.
- GAMBLE, F. M., 1980. Rainfall in the Namib Desert Park. *Madoqua* 12: 175–180.
- GOLDREICH, Y., DRUYAN, L. M. and BERGER, H., 1986. The interaction of valley/mountain winds with a diurnally veering sea/land breeze. *Journal of Climatology* 6: 551–562.
- GOLDREICH, Y. and TYSON, P. D., 1988. Diurnal and inter-diurnal variations in large-scale atmospheric turbulence over southern Africa. *South African Geographical Journal* 70: 48–56.
- HADLEY, N. F. and SZAREK, S. R., 1981. Productivity of desert ecosystems. *Bio-Science* 31: 747–753.
- HAMILTON, W. J. III and SEELY, M. K., 1976. Fog basking by the Namib desert beetle, *Onymacris unguicularis*. *Nature* 262: 284–285.
- HARMSE, J. T., 1982. Geomorphologically effective winds in the northern part of the Namib Sand Desert. *South African Geographer* 10: 43–52.
- HATTLE, A., 1985. Surface water hydrology of the Kuiseb. In: HUNTLEY, B. J., ed., *The Kuiseb environment: the development of a monitoring baseline*, pp. 27–32. South African National Scientific Programmes Report No. 106, F.R.D.
- HAURWITZ, B., 1947. Comments on the sea-breeze circulation. *Journal of Meteorology* 4: 1–8.
- HAWKES, H. B., 1945. Mountain and valley winds. *Weather Division, U.S. Army Signal Corps, Report No. 982*, 44 pp.
- JACKSON, S. P., 1942. Weather on the coasts of South Africa. *Meteorological Services of the Royal Navy and South African Air Force Parts 1–5*, 372 pp.
- JACKSON, S. P., 1954. Sea breezes in South Africa. *South African Geographical Journal* 36: 13–23.
- LANCASTER, I. N., 1983. Controls of dune morphology in the Namib sand sea. In: BROOKFIELD, M. E. and AHLBRANDT T. S., eds, *Eolian sediments and processes*, pp. 261–289. Elsevier, Amsterdam.
- LANCASTER, I. N., 1985. Winds and sand movements in the Namib sand sea. *Earth Surface Processes and Landforms* 10: 607–619.
- LANCASTER, J., LANCASTER, N. and SEELY, M. K., 1984. Climate of the central Namib Desert. *Madoqua* 14: 5–61.
- LAUER, W. and KLAUS, D., 1975. The thermal circulation of the central Mexican meseta region within the influence of the trade winds. *Archiv für Meteorologie, Geophysik und Bioklimatologie, Ser. A* 23: 343–366.
- LINDESAY, J. A. and TYSON, P. D., 1990. Thermo-topographically induced boundary layer oscillations over the central Namib. *Journal of Climatology* 10: 63–77.
- LOUW, G. N., 1972. The role of advective fog in the water economy of certain Namib Desert animals. *Symposium of the Zoological Society of London* No. 31: 297–314.
- LOUW, G. N. and SEELY, M. K., 1980. Exploitation of fog water by a perennial Namib dune grass, *Stipogrostis sabulicola*. *South African Journal of Science* 76: 38–39.
- MCKENDRY, I. G., 1983. Spatial and temporal aspects of the surface wind regime on the Canterbury Plains, New Zealand. *Journal of Climatology* 3: 155–166.
- MCKENDRY, I. G., STURMAN, A. P. and OWENS, I. F., 1986. A study of interacting multi-scale wind systems, Canterbury Plains, New Zealand. *Meteorology and Atmospheric Physics* 35: 242–252.
- MARKER, M. E., 1977. A long-return geomorphic event in the Namib Desert, South West Africa. *Area* 9: 209–213.
- MATHEWS, J. H., 1982. The sea breeze front – forecasting aspects. *Australian Meteorological Magazine* 30: 205–209.
- MEYER, J. H., 1971. Radar observations of land breeze fronts. *Journal of Applied Meteorology* 10: 1224–1232.
- NAGEL, J. F., 1962. Fog precipitation measurements on Africa's southwest coast. *Notas* 11: 51–60.
- NIEMAN, W. A., HEYNS, C. and SEELY, M. K., 1978. A note on precipitation at Swakopmund. *Madoqua* 11: 69–73.
- PIELKE, R. A., 1984. *Mesoscale meteorological modeling*. Academic Press, London.
- PIETRUSZKA, R. D. and SEELY, M. K., 1985. Predictability of two moisture sources in the Namib Desert. *South African Journal of Science* 81: 682–685.
- PRESTON-WHYTE, R. A., 1969. Sea breeze studies in Natal. *South African Geographical Journal* 51: 38–49.
- PRESTON-WHYTE, R. A., 1974. Land breezes and mountain-plain winds over the Natal coast. *South African Geographical Journal* 56: 27–35.
- PRESTON-WHYTE, R. A. and TYSON, P. D., 1988. *The atmosphere and weather of southern Africa*. Oxford University Press, Cape Town.
- PREZERAKOS, N. G., 1986. Characteristics of the sea breeze in Attica, Greece. *Boundary Layer Meteorology* 36: 245–266.
- REYNOLDS, R. D., 1966. The effect of atmospheric lapse rate on balloon ascent rates. *Journal of Applied Meteorology* 5: 537–541.
- ROTUNNO, R., 1983. On the linear theory of the land and sea breeze. *Journal of the Atmospheric Sciences* 40: 1999–2009.
- ROY, A. K., 1940. The sea-breeze at Madras. *Scientific Notes of the Indian Meteorology Department* 8: 138–146.
- SCHULZE, B. R., 1969. The climate of Gobabeb. *Scientific Papers of the Namib Desert Research Station* 38: 5–12.
- SEELY, M. K., 1979a. Ecology of a living desert: twenty years of research in the Namib. *South African Journal of Science* 75: 298–303.
- SEELY, M. K., 1979b. Irregular fog as a water source for desert dune beetles. *Oecologia (Berlin)* 42: 213–227.
- SEELY, M. K., DE VOS, M. P. and LOUW, G. N., 1977. Fog imbibation, satellite fauna and unusual leaf structure in a Namib Desert dune plant, *Trianthema hereroensis*. *South African Journal of Science* 73: 169–172.
- SEELY, M. K. and HAMILTON, W. J. III, 1976. Fog catchment sand

- trenches constructed by tenebrionid beetles, *Lepidochora*, from the Namib Desert. *Science* **193**: 484–486.
- SEELY, M. K., LEWIS, C. K., O'BRIEN, K. A. and SUTTLE, A. E., 1983. Fog response of tenebrionid beetles in the Namib Desert. *Journal of Arid Environments* **6**: 135–143.
- SEELY, M. K. and LOUW, G. N., 1980. First approximation of the effects of rainfall on the ecology and energetics of a Namib Desert dune ecosystem. *Journal of Arid Environments* **3**: 25–54.
- SEELY, M. K. and STUART, P., 1976. Namib climate: 2. The climate of Gobabeb, ten year summary 1962–1972. *Namib Bulletin* **1**: 7–9.
- SEGAL, M., PURDOM, J. F. W., SONG, J. L., PIELKE, R. A. and MAHRER, Y., 1986. Evaluation of cloud shading effects on the generation and modification of mesoscale circulations. *Monthly Weather Review* **114**: 1201–1212.
- SHARON, D., 1981. The distribution in space of local rainfall in the Namib desert. *Journal of Climatology* **1**: 69–75.
- SKIBIN, D. and HOD, A., 1979. Subjective analysis of mesoscale flow patterns in northern Israel. *Journal of Applied Meteorology* **18**: 329–338.
- STURMAN, A. P., 1987. Thermal influences on airflow in mountainous terrain. *Progress in Physical Geography* **11**: 183–206.
- STURMAN, A. P., FITZSIMONS, S. J. and HOLLAND, L. M., 1985. Local winds in the Southern Alps, New Zealand. *Journal of Climatology* **5**: 145–160.
- STURMAN, A. P. and TYSON, P. D., 1981. Sea breezes along the Canterbury coast in the vicinity of Christchurch, New Zealand. *Journal of Climatology* **1**: 203–219.
- TALJAARD, J. J. and SCHUMANN, T. E. W., 1940. Upper air temperatures and humidities at Walvis Bay, South West Africa. *Bulletin of the American Meteorological Society* **21**: 293–296.
- TYSON, P. D., 1966. Examples of local air circulations over Cato Ridge during July 1965. *South African Geographical Journal* **48**: 13–31.
- TYSON, P. D., 1967. Some characteristics of the mountain wind over Pietermaritzburg. *Proceedings of the Jubilee Conference of the South African Geographical Society*, Durban, pp. 103–128.
- TYSON, P. D., 1968a. Nocturnal local winds in a Drakensberg valley. *South African Geographical Journal* **50**: 15–32.
- TYSON, P. D., 1968b. Southeasterly winds over Natal. *Journal for Geography* **3**: 237–246.
- TYSON, P. D., 1968c. Velocity fluctuations in the mountain wind. *Journal of the Atmospheric Sciences* **25**: 381–384.
- TYSON, P. D. and PRESTON-WHYTE, R. A., 1972. Observations of regional topographically-induced wind systems in Natal. *Journal of Applied Meteorology* **11**: 643–650.
- TYSON, P. D. and SEELY, M. K., 1980. Local winds over the central Namib. *South African Geographical Journal* **62**: 135–150.
- VAN BEMMELEN, W., 1922. Land -und Seebrise über Batavia. *Beiträge zur Physik der Atmosphäre* **10**: 169–177.
- WARD, J. D. and VON BRUNN, V., 1985. Sand dynamics along the lower Kuiseb River. In: HUNTLEY, B. J., ed., *The Kuiseb environment: the development of a monitoring baseline*, pp. 51–72. South African National Scientific Programmes Report No. 106, F.R.D.
- WILKINSON, M. J., 1987. A Late-Cenozoic geomorphic history of the Tumas drainage basin in the central Namib Desert. Unpublished Ph.D. thesis, University of Chicago.
- YAN, H. and ANTHES, R. A., 1987. The effect of latitude on the sea breeze. *Monthly Weather Review* **115**: 936–956.
- YOSHINO, M. M., 1981. Orographically-induced atmospheric circulations. *Progress in Physical Geography* **5**: 76–98.