

LANIOTURDUS

Newsletter of the SWA/Namibia Bird Club

Mitteilungen des SWA/Namibia Vogelklubs

Vol./Jg. 23, No. 1

1987

SWA/NAMIBIA VOGELKLUB
eine Zweigstelle der
SWA Wissenschaftlichen Gesellschaft
und der
Southern African
Ornithological Society

Beiträge bitte an den Redakteur p.A. SWA Wissenschaftliche Gesellschaft, Postfach 67, Windhoek 9000, SWA senden.

Mitgliedsbeiträge für die S.A.O.S. und die Wissenschaftliche Gesellschaft für 1987 sind wie folgt:

Privatperson: S.A.O.S. + SWA Wissenschaftliche Gesellschaft (mit Ostrich)	R 58.00
Privatperson: S.A.O.S. +SWA Wissenschaftliche Gesellschaft (ohne Ostrich)	R 50.00
Schüler und Studenten unter 24 Jahren: (mit Ostrich)	R 38.00
Schüler und Studenten unter 24 Jahren: (ohne Ostrich)	R 30.00
Privatperson: nur SWA Wissenschaftliche Gesellschaft	R 30.00
Schüler und Studenten: nur SWA Wissenschaftliche Gesellschaft	R 20.00
Abonnement nur für den LANIOTURDUS	R 10.00

Bemerkungen an Mitarbeiter

Manuskripte sollten mit Schreibmaschine (oder in gut leserlicher Handschrift) geschrieben sein und zwar mit doppeltem Zeilenabstand auf A4 (30 x 21 cm) Format. Skizzen, Karten und Tabellen sollten auf weißem Qualitätspapier mit schwarzer Tinte gezeichnet werden. Klare Schwarz-weiss-Photographien (15 x 20 cm) können eingereicht werden, um die Arbeit zu illustrieren.

CONTENTS

INHALTSANGABE

EDITORIAL/REDAKTION	2
ARTICLES & REPORTS/ARTIKEL & BERICHTE:	
Coastal Bird Counts along the Namib Coast, Cape Cross to Lüderitz, November and December 1985	3
NOTES and NEWS/BEMERKUNGEN & MITTEILUNGEN:	
The feeding pattern of some birds in a Suiderhof garden ..	9
Bemerkungen zu den Rotbraunen Webern in Windhoek	10
Fair Warning	10
PROJECTS/PROJEKTE:	
Look out for yellow waders this spring	12
Bustard Alert	12
BIRD CLUB ACTIVITIES/AKTIVITÄTEN DES VOGELKLUBS:	
Jahresbericht des SWA/Namibia Vogelklubs 1986	14
Income and Expenditure Account for the year ending 31.12.1987	15
Walvis Bay Bird Count	16
Birdslide-Archives	20
NOTICES/BEKANNTMACHUNGEN:	
XX International Ornithological Congress 1990	22
Ornithologische Literatur zu kaufen	22

English and scientific names	Count areas				Total
	1	2	3	4	
Bank Cormorant <u>Phalacrocorax neglectus</u>	5	4	137	0	146
Cape Cormorant <u>Phalacrocorax capensis</u>	1	2	20	0	23
Crowned Cormorant <u>Phalacrocorax coronatus</u>	17	0	53	0	70
Grey Heron <u>Ardea cinereus</u>	1	0	1	0	2
Black Oystercatcher <u>Haematopus moquini</u>	96	68	74	5	243
Whitefronted Plover <u>Charadrius marginatus</u>	26	14	23	0	63
Kelp Gull <u>Larus dominicanus</u>	18	0	48	63	129
Hartlaub's Gull <u>Larus hartlaubii</u>	0	0	4	0	4
Swift Tern <u>Sterna bergii</u>	0	5	64	0	69
Damara Tern <u>Sterna balaenarum</u>	0	0	6+	0	6+
Total: resident species	174	93	462	68	797

Table 2(II)

Palaeartic migrant species

Grey Plover <u>Pluvialis squatarola</u>	10	0	38	2	50
Turnstone <u>Arenaria interpres</u>	41	9	117	0	167
Common Sandpiper <u>Tringa hypoleucos</u>	1	0	2	0	3
Curlew Sandpiper <u>Calidris ferruginea</u>	7	0	4	0	11
Sanderling <u>Calidris alba</u>	4	0	26	0	30
Ruff <u>Philomachus pugnax</u>	1	0	0	0	1
Bartailed Godwit <u>Limosa lapponica</u>	1	0	0	0	1
Whimbrel <u>Numenius phaeopus</u>	5	0	4	0	9
Common Tern <u>Sterna hirundo</u>	0	0	4	0	4
Sandwich Tern <u>Sterna sandvicensis</u>	0	1	0	0	1
Total : Palaeartic migrants	70	10	195	2	277
Grand total number of birds	244	103	657	70	1,074

Count areas are: 1) Lüderitz - near Danger Rocks; 2) Flamingo Island; 3) Aurus to Wreck Point; and 4) Spencer Bay.

.....oooooOOOOOooooo.....

Notes and News

Bemerkungen & Mitteilungen

THE FEEDING PATTERN OF SOME BIRDS IN A SUIDERHOF GARDEN

During November and December 1986 I established that some of the common Windhoek birds (mostly seed-eaters) followed a fairly consistent pattern of arrival and feeding at the feeding site in my garden in Suiderhof. Food is set out once-daily, usually in the afternoon.

Very little activity takes place until about 17h30, when the first birds to arrive are the Blackthroated Canaries Serinus atrogularis. Within a few minutes the first Laughing Doves Streptopelia senegalensis arrive, but continue to come and go throughout the feeding 'session'. The doves are usually followed closely by small numbers of Paradise and Shafttailed Whydas Vidua paradisea and V. regia. Minutes later House Sparrows Passer domesticus (in varying numbers) and the first few Chestnut Weavers Ploceus rubiginosus arrive. By 17h45 the main group of Chestnut Weavers arrives, at which stage the Blackthroated Canaries leave. Lastly, a few Redbilled Queleas Quelea quelea arrive. Two or three White-backed Mousebirds Colius colius usually arrive during the feeding session, but avoid the 'rabble' and feed on the 'dubbeltjie' (Morgensstern) plants in the garden.

Although the timing of arrival varies from day to day, there is usually very little variation in the sequence of species arriving to feed.

SUZANN MALLET-VEALE, P.O. Box 3787, Windhoek, 9000

HÄGAR the Horrible

®

By Dik Browne

