

# GAME COUNTS IN NORTH-WEST NAMIBIA

## Conservancies north of the veterinary fence


May 2021

### Total Population Estimates

Species	Population estimate	Lower 95% CL	Upper 95% CL	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Gemsbok (U)	435	167	1,135																					
Kudu (U)	468	188	1,161																					
Ostrich (U)	2,233	1,491	3,344																					
Springbok (U)	45,788	24,030	87,245																					
Steenbok (U)	2,205	1,329	3,659																					
Hartmann's Zebra (U)	2,604	1,533	4,421																					

All above estimates are derived using DISTANCE analysis. This takes account of drop off in detection with distance from the transect line. They are conservative estimates as 35% of the count area is not sampled (due to inaccessibility) and is consequently assumed to hold no animals. Model selection: U = uniform key; HN = half normal

### Total number of animals seen each year




### Animals seen during this count and minimum estimates

Species	Total	Anabeb	Ehi-Rovipukta	Marienfluss	Okangunduma	Okondjombo	Omatendeka	Ombujokangundi	Ongongo	Oropembe	Orupupa	Otjambangu	Otjikondavirongo	Otjiu-west	Otzemba	Ozondunu	Puros	Sanitas	Sesfontein
Total Route km	3,247	233	273	281	139	192	211	158	64	292	159	56	135	88	102	77	328	151	307
Total area (km <sup>2</sup> )	28,288	1,636	1,979	3,034	1,130	1,643	1,613	657	619	2,616	1,775	348	1,067	1,208	741	743	3,564	1,446	2,469
Number of routes	74	5	5	6	3	4	4	3	5	6	3	1	4	2	3	2	6	4	8
% area excluded	35	51	28	28	29	16	48	26	53	10	44	74	58	71	26	55	31	28	42

Values without brackets are numbers of animals seen along transects. Values inside brackets are minimum estimates assuming all animals within 500m on each side of the transect line are detected i.e. there is no adjustment for drop off in detection with distance from the transect line. In addition, for springbok, gemsbok and giraffe, large groups were excluded from extrapolations and added afterwards.



The sum of these values will be significantly lower than the totals indicated in the top left table as the total estimates take account of species detection curves.

### Rainfall




The rainfall season is from July to June and values are an average for the whole area. The year represents the season immediately prior to the count.

### Vegetation



NDVI is a measure of the density of chlorophyll in vegetation cover. It can be used as an indicator of the amount of biomass available to wildlife. The map shows the NDVI status in the current year (Feb-Apr) and the trend indicates the average deviation of northern area conservancies from the long-term mean, in each year.

### Mortalities



Total number of carcasses from all species recorded in the Event Book