

W1504

D.R.F.N.	: REPRINT	89
REFERENCE:	507	
LIBRARY	: WIK	

JOURNAL 48
Namibia Wissenschaftliche Gesellschaft / Namibia Scientific Society
Windhoek, Namibia 2000
ISSN: 1018 - 7677

LONG-TERM ECOLOGICAL RESEARCH AT GOBABEB: GAINING AND APPLYING KNOWLEDGE ABOUT A HIGHLY VARIABLE ENVIRONMENT

JOH R. HENSCHER, MARY K. SEELY and JULIANE ZEIDLER

Desert Research Foundation of Namibia, Gobabeb, P.O.Box 20232, Windhoek, Namibia
e-mail: drfn@drfn.org.na

Abstract

Long-term ecological research (LTER) is valuable for understanding environmental processes in arid countries like Namibia. It is also essential for effective natural resource management. Using a wealth of knowledge accumulated over the years in the Central Namib Desert and, more recently, in rural communal areas in western Namibia, the Desert Research Foundation of Namibia (DRFN) is using LTER to connect basic and applied research in an appropriate manner. Various projects concern climate, the biophysical environment, plants, animals, and people. Examples are given of how these projects are interrelated and how they serve many social and environmental functions in rural Namibia. The formation of a national Namibian LTER network, and Namibia's membership in the International Long Term Ecological Research Network affect the future of LTER conducted by DRFN and other Namibian organisations.

Introduction

The Desert Research Foundation of Namibia at the Gobabeb Training and Research Centre, situated in the Central Namib, has set itself the goal of increasing understanding of a hyperarid environment, especially its variability, that can be applied to other variable environments to help reduce the latter's vulnerability to loss of sustainable livelihoods (SEELY *et al.*, this volume). One of the major means of increasing understanding, is through Long Term Ecological Research (LTER), continuous or periodic monitoring combined with detailed short-term studies of numerous environmental aspects. By conducting LTER in the Namib Desert, a hyperarid region, it is possible to understand the effects of extreme hydrological variability, and to apply this knowledge appropriately in line with the above-stated goal.

Understanding natural environmental variability is important for environmental management (LANDRES *et al.*, 1999). Tropical arid and semi-arid regions like those characterising much of Na-

Namibia experience extreme temporal and spatial variation of rainfall (TYSON, 1986; NICHOLS and WONG, 1990) due to the negative correlation between rainfall variability and aridity (TYSON and DYER, 1975; Fig.1). In arid regions, brief episodic events affect long-term processes (CRAWFORD and GOSZ, 1982; WEATHERHEAD, 1986). In particular, heavy rainfall events elicit dramatic responses (SEELY and LOUW, 1980; NOY-MEIR, 1981), have effects lasting for decades (WALTER, 1971; SOUTHGATE *et al.*, 1996; HENSCHHEL, SEELY and POLIS, 1998), and are thus prominent ecosystem drivers. Furthermore, there is high spatial variability of rainfall in arid regions (DEAN and MILTON, 1999; WARD and ELLIS, 1999; Fig. 1). This leads to spatial concentration of resource production, and rich patches can affect large resource-poor areas, i.e. populations in rich patches are sources for surrounding sink populations that are sustained by them (PULLIAM, 1988; WATSON and SUTHERLAND, 1995; DIAS, 1996). Rich patches arise, expand, shift and contract dynamically at variable rates (JELTSCH *et al.*, 1998; MILTON, 1995; REYNOLDS *et al.*, 1999). Biotic responses to temporal and spatial variability may depend either on pulses and sources, i.e. irruption, or on tolerance of reserve and sink conditions, i.e. persistence. These responses affect and are affected by people, and it is therefore important to understand natural variability in order to facilitate appropriate environmental management (LANDRES *et al.*, 1999).

Long-term Ecological Research (LTER) enables one to recognise and elucidate natural patterns of temporal and spatial variability and changes in such patterns (RISSER, 1995). LTER integrates the findings of continuous or periodic environmental monitoring with short, experimental studies. LTER focuses on time spans of decades to a century, and encompasses historic records inasmuch as they are based on measures that are repeatable (LTER, 2000). While paleoclimatological and archaeological research makes inferences on environmental and anthropogenic changes over time spans of centuries to millions of years, more direct measurements are required to understand the complex patterns and factors underlying environmental variability and changes during present times and the past century. During the latter time span, it is not only possible to elucidate complexity, but also possible to influence the outcome, as this time span encompasses funding cycles of projects, legislative periods of governments, and the generations and lifetimes of people. LTER covers this time span and it can thus provide researchers, resource managers, and policy makers with data needed to detect, quantify, locate, understand and respond to changes in terrestrial ecosystems so as to support sustainable development (GTOS, 2000). SEELY (1998), SEELY and JACOBSON (1994) and SEELY *et al.* (submitted) outline how this connection between science and development serves creative problem solving in Namibia.

LTER provides baseline data on long-term environmental processes, including responses to episodic events. It facilitates the interpretation of short-term studies, and it provides the connecting framework between different studies. LTER can record environmental changes caused by human activities and elucidate factors affecting people, combining biophysical with socio-economic data. These data do not only support a theoretical framework, but are an invaluable source of information for generating environmental awareness, influencing decision-makers, and training students in environmental fields. LTER is therefore an invaluable tool to guide the development process in Namibia.

In August 1999, Namibia formed a national LTER network (Na-LTER, 2000) and became the first country in Africa to join the International Long-Term Network, ILTER (HENSCHTEL, 1999). ILTER is a growing global network comprising countries around the world that co-ordinate and share research efforts and results through this network (FRANKLIN *et al.*, 1990; RISSER, 1995; GOSZ, 1996; ILTER, 1999, 2000). Allied to ILTER, the Global Terrestrial Observation System (GTOS, 2000) also endeavours to improve the sharing of information and data on sites being monitored globally on a long-term basis. Na-LTER is connected to these international networks so as to facilitate the sharing of data, information and experience in both directions, especially so as to achieve value-addition through such collaboration.

The purpose of the Namibian Na-LTER network is that its partners have increased capacity to provide, access, understand and use long-term ecological data and information in Namibia (Na-LTER, 2000). The Na-LTER network comprises institutions and researchers willing to contribute to its goals, purpose and objectives. Na-LTER endeavours to identify, promote and facilitate the appropriate operation of sites where long-term environmental monitoring and analyses are conducted. A core facility of Na-LTER is planned to be a Metadatabase, comprising information on data, namely what the data concern, where, when, and by whom they were obtained, how and where others can obtain copies, conditions of use, and other details.

Gobabeb is a national Na-LTER and international ILTER site, where the LTER programme, conducted in collaboration with many institutions, forms part of the institutional programme of the Desert Research Foundation of Namibia (SEELY *et al.*, this volume). The premise for the joint venture agreement on the Gobabeb Training and Research Centre between MET and DRFN provides this site with the collaborative spirit that is central to LTER. Other Namibian LTER sites are part of monitoring programmes in the Etosha National Park (e.g. Etosha Ecological Institute), and at agricultural, forestry and university research stations. Yet another type are LTER sites that are not directly affiliated to on-site research stations, e.g., parks, reserves, and conservancies, weather stations, hydrological monitoring points, coastal and other wetlands, river catchments, the Eastern National Water Carrier, Inselbergs, desertification programme study sites, Sardep agricultural study sites, Rehoboth Acacia Forest, Polytechnic field sites, and those of foreign programmes, such as BIOTA. A further type are other regional or site-specific data shared through the Na-LTER network, such as socio-economic monitoring by NEPRU (Namibian Economic Policy Research Unit), livestock censuses by Veterinary Services, wildlife censuses by Directorate of Specialist Support Services, as well as runoff and groundwater monitoring by Department of Water Affairs.

In this paper we describe the LTER programme co-ordinated by the DRFN at Gobabeb. We also show how LTER is being extended and applied in different ways since Independence (SEELY *et al.*, submitted). Thirty-eight years of research in the Namib (since 1962) has enabled the DRFN to gain significant knowledge on environmental processes under arid conditions (Henschel *et al.*, 2000). Since Independence, this knowledge is being used to further environmental awareness and to promote the capacity to manage the Namibian environment appropriately. This is addressed through research, information and training programmes that focus environmental fields such as climatic variability and

its influence on the environment, ephemeral rivers and their catchments, and desertification. LTER is fundamental to these programmes, and this paper focuses on its biophysical and socio-economic aspects. Besides strong applications for training, we emphasise the importance of involving people, not only to raise awareness, but also to facilitate the process by which rural communities become an integral part of monitoring and managing the environment in Namibia.

This paper does not attempt to provide a comprehensive review of the application of the knowledge gained through LTER at Gobabeb, nor of research in the Namib where numerous other organisations are involved. Besides describing the Gobabeb LTER programme, we provide examples of how the DRFN and its associated students and scientists are applying information derived from it, as well as setting up monitoring sites in the areas of application.

Study Areas and General Methods

LTER around the Gobabeb Training and Research Centre (23°33'S; 15°02'E) encompasses several sites in the Central Namib Desert Study sites are in the three major habitats characterising the Central Namib, on the gravel plains, in the Kuiseb riverbed and in the dunefield of the Great Sand Sea. The general climatic, geological, geomorphological and ecological conditions were described e.g. by BESLER (1972a), GOUDIE (1972), ROBINSON (1978), ROBINSON and SEELY (1980), LANCASTER *et al.* (1984) and SEELY (1990a). Study sites are monitored at regular intervals, varying between projects from daily to monthly, seasonally, annually, or at intervals of several years. Some projects were initiated by short-term research projects when the methods and monitoring protocol were established. Monitoring then continued after the publication of initial results identified the potential value of long-term data.

The sites of the Namibian Programme to Combat Desertification (NAPCOD) where DRFN has initiated studies that can continue with LTER are situated in the communal and commercial farming areas of the former Damaraland in north-western Namibia (Kunene Region). This and other farming areas of Namibia, not described here, form part of the NAPCOD-III programme (ZEIDLER, 2000). The north-west sites include farms with a background of commercial, communal and alternating commercial/communal land tenure that depended on changing land tenure policies under different governments (KAMBATUKU, 1996; ZEIDLER, 1999). In the current paper we use one of the communal farms, Olifantputs (20°26'S; 14°96'E), as example. LTER at the NAPCOD sites follows the guiding principles of the Convention to Combat Desertification (CCD), such as conducting participatory, community-based and community-relevant research (UNEP, 1995). Besides the establishment of ecological monitoring programmes focusing on climatic, vegetative, soil-related and biotic components of the environment, it is important to monitor aspects relating to people, such as the history of land use practices (KAMBATUKU, 1996), population dynamics and migratory patterns of people and livestock, changes in the use of natural resources, and lifestyles (ZEIDLER *et al.*, 1998; ZEIDLER, 1999). The local farmers are actively involved with the DRFN in the collection, management, and interpretation of data.

The Projects

Table 1 summarises the different research foci and trajectories referred to in this section.

	Conservation areas	Agricultural areas	ELSEWHERE IN NAMIBIA (since 1992)
Climate	• Nature reserves	• Rangelands	• Rainfall records by farmers
Biophysical environment	• Restricted areas	• Cultivated fields	• Fog harvesting
Plants • natural vegetation • crops	• Communal farmers		• Soil processes
Animals • vertebrates & invertebrates			• Ephemeral river ecosystems
People • demography • land uses • livestock			• Grass production
			• Community composition
			• Repeat photography
			• Game
			• Invertebrate biodiversity
			• Livestock populations
			• Household demographics
			• Natural resource use
			• Land use practices

	NAMIB (Gobabeb)	YEAR started
Existing long-term record from desert area		
Climate	• Climate	• 1962
Biophysical environment	• Fog	• 1970
Plants	• Renewable energy & groundwater	• 1996
Animals	• Dune dynamics	• 1970
People	• Weathering	• 1976
	• Off-road vehicle impact on surfaces	• 1978
	• Kuiseb River ecosystem	• 1962
	• Welwitschia and Inara	• 1985
	• Grass production	• 1974
	• Plant community dynamics	• 1976
	• Fixed point photography	• 1997
	• Game, small mammal, lizard & invertebrate populations	• 1968
	• Tenebrionid beetle populations	• 1968
	• Livestock along Kuiseb	• 1976
	• Topnaar natural resource management, esp. Nara & water	• 1976


LTER projects being carried out by the Desert Research Foundation of Namibia.

Climate

Long-term weather data are fundamental to the understanding of a highly variable environment such as the Namib. At Gobabeb, weather has been recorded continuously since 1962 at the First Order Weather Station of the Namibian Meteorological Services. Furthermore, DRFN operates a network of autographic weather stations covering an area of 10 000 km². Of the 45 stations, there are eight main ones that are operated continuously and cover the width of the Namib from near the coast (Radio Station "33" 10 km east of Walvis Bay, at an altitude of 40 m above mean sea level) up to above the Great Escarpment (Farm Weissenfels, 10 km east of the Gamsberg, altitude 1800 m above mean sea level). These weather stations are fitted with electronic data loggers and software that transcribe data at hourly intervals and are monthly copied to computer (Mike Cotton Systems, Steenberg, South Africa). Other organisations that also operate weather stations in the Namib include the Namibian Meteorological Services (several manually and automatic stations along the entire Namib coast) and the University of Cologne (several weather stations between Swakopmund and Usakos).

The surface weather records of the DRFN have been used to characterise Namib climate (LANCASTER *et al.*, 1984). They have documented episodic events, such as rare, heavy rainfall, or a poor fog season, heavy sand storm, or heat wave. For example, the sources of surface water at Gobabeb differ in their variability. Fog from the Atlantic Ocean is the most regular source (mean = 39 mm per annum; CV = 43%), while rain is sporadic (21.2 mm; CV = 122%). Runoff in the ephemeral Kuiseb river from rainfall in the inland catchment in the Khomashochland occurs nearly every year, but differs in duration (18 days; CV = 150%). Variability in rainfall differs across the Namib (Fig.1), and this is very important for understanding the composition of biotic communities, the distribution of organisms and their life history patterns (HOLM, 1970; ROBINSON, 1978; SEELY, 1973, 1978a, b, 1990b; SEELY and LOUW, 1980; NEL, 1983; HAMILTON, 1985, 1986; TILSON and HENSCHEL, 1986; YEATON, 1988; BOYER, 1989; BERRY and SIEGFRIED, 1991; GÜNSTER, 1993, 1995; BRAIN, 1993; KILIAN, 1995; KOK and NEL, 1996; SOUTHGATE *et al.*, 1996; HACHFELD, 1996; JACOBSON, 1996, 1997a; BURKE, 1997; JACOBSON and JACOBSON, 1998; HENSCHEL, SEELY and POLIS, 1998; HENSCHEL and SEELY, in press). Micro-climate varies between habitats and this has been monitored at several sites around Gobabeb. Most of the research and training at Gobabeb incorporates one or the other climatic or micro-climatic aspect, and it is seen as the common thread that links most fields. Climate is among the fundamental factors required for effective planning development in the region (BENDER, 1999).

The many instruments and data bank at Gobabeb are invaluable tools for training. The diversity of instruments illustrate fundamental climatic principles, while the data bank allows climatologists, ecologists, geomorphologists, appropriate technologists, and students to understand these principles in the context of the spatial and temporal variability of climate.



Coefficient of variation (CV) versus mean annual rainfall (mm) at five places across the Namibia Desert and eastwards. Numbers in brackets indicate distance (km) for the coast. Data are from Meteorological Services of Namibia (n = 20-108 years)

Since the beginning of 1997, rainfall data are collected at sites in southern Kunene region by communal farmers in a participatory manner, i.e. farmers and other community members monitor the rain gauges and keep records. Data analyses and interpretation are collaborative with NAPCOD field staff. Discussions facilitate the understanding of the spatial and temporal variability of rainfall, which assist farmers in planning sustainable resource management.

As an important climatic feature of the Namib, fog has been studied over 60 years (e.g., WALTER, 1937; BOSS, 1941; LANCASTER *et al.*, 1984). In the last three years monitoring has been intensified with the aim of collecting this water source for domestic use in the western Namib Desert (HENSCHEL, MTULENI *et al.*, 1998), including gardens in Swakopmund (COETZEE and MULDER, pers.comm.). The possibility is being explored to utilise sun and wind energy for small solar stills to purify substandard water and to provide electricity to remote centres, such as Gobabeb (SHANYENGANA, 1997a, b). The LTER data series are invaluable for planning these applications.

Biophysical Environment

Dune movements - Namib dune dynamics have been monitored since 1970. Such measurements were established to monitor the geomorphological processes of dune movements *per se* as well as the relation thereof to the biotic system. It has been suggested that the high diversity of certain taxa such as tenebrionids in the Namib Desert can be explained by the mobility and changing configuration of sand dunes (ENDRÖDY-YOUNGA, 1982; PENRITH, 1986; IRISH, 1990). Populations of psammophilous animals become geographically isolated from conspecifics when their dune moves. Conversely, two dunes that were not previously connected may connect and their satellite organisms become sympatric. The Namib Great Sand Sea has many types of dunes with different configurations and dynamics changing along an east-west gradient (LANCASTER, 1989). The middle of the dunefield, including the area south of Gobabeb, is dominated by long linear dunes. Since BESLER's (1970, 1972b) initial study, DRFN has monitored the physical dynamics of one such linear dune on a monthly basis. Here, trainees can learn environmental principles such as the relationship between dune configuration, microhabitats, and associated animals and plants. Students who have been sensitised in this way are better able to recognise similar relationships in more complex environments.

Weathering - The Central Namib Desert represents a hostile environment for rock, with salt, temperature changes, fog and lichens, all known to be important weathering agents. Over time, weathering produces debris (including silt-sized material) and leaves eroded rock. This process is being studied by monitoring weathering rates and processes across an East-West transect through the Central Namib Desert over a period of many years. Changes in the micro-environmental conditions are being measured so as to elucidate their effects (GOUDIE, 1972; GOUDIE *et al.*, 1997; GOUDIE and PARKER, 1998), and assist planning of development. The deployed blocks of rock at Gobabeb is very useful for demonstrating the principles of weathering to environmental trainees.

Off-road vehicles (ORV) - The impact of ORVs on the desert surface has been monitored since 1978 (SEELY and HAMILTON, 1978; DANEEL, 1992a, b). Vehicles change the properties of soil, inclu-

ding the desert pavement, calcrete and gypsum crusts, as well as the biota, particularly lichens and dwarf shrubs. For example, ORVs reduce the microtopography and push stones and lichens into the soil, thereby reducing surfaces for lichens to grow on and impeding their recovery; a fresh ORV track destroys 80% of the lichens (DANEEL, 1992a). These properties take a long time to recover (ECKARDT, 1996) and long-term studies are required to monitor recovery. SEELY and HAMILTON (1978) and DANEEL (1992a), respectively, placed control tracks in an interdune and several gravel plain areas in the Namib respectively. By monitoring these tracks at intervals of several years, we are learning about the recovery process, including that of lichen fields. ORV tracks enable students to become sensitised to critical soil processes. At a stage when the use of ORVs appears to be increasing in western Namibia, this study provides a good basis for planning and for advice on driver behaviour.

Soil - Soil processes play a major role in sustaining productivity of agricultural production areas, including crop-lands as well as range systems (GREENLAND and SZABOLCS, 1994). This is important particularly in view of the concerns about desertification and land degradation in Namibia, but also for biodiversity research because many soil processes are mediated by the soil biota (ANDERSON, 1994). Soil processes are being monitored at NAPCOD sites in southern Kunene region. This concerns the biological integrity of farms under different land tenure and different land-use intensities, with particular emphasis on the communal farm Olifantputs. At six different monitoring plots of 1 ha each, soil parameters such as nutrient status and biophysical parameters are measured twice a year since 1997. This is done to gauge the effect of various kinds of communal and commercial farming on rangeland and habitat condition in western Namibia (ZEIDLER *et al.*, 1998; ZEIDLER, 1999). These data are correlated with the biodiversity of soil organisms. Other long-term studies can be initiated by revisiting sites that were part of other applied studies associated with Gobabeb (e.g., WARD *et al.*, 1998).

Ephemeral Rivers - For many years, the Kuiseb river ecosystem has been studied from the biophysical and ecological perspectives, while it also serves as an excellent outdoor classroom for students who can learn how natural and anthropogenic processes in this linear oasis contrast with and influence the surrounding desert habitats. Since 1963 floods and groundwater level have been monitored, and on several occasions since 1976, aerial photographs were taken (e.g. HUNTLEY, 1985; JACOBSON *et al.*, 1995). The effect of changes, natural as well as anthropogenic, on vegetation and ultimately wild and domestic animals are recorded (e.g., HAMILTON *et al.*, 1977; SEELY, BUSKIRK and HAMILTON, 1980; THERON and VAN ROOYEN, 1980; TILSON and HENSCHER, 1986; GABRIEL, 1992; DAUSAB *et al.*, 1993; ZEIDLER, 1995; AMOOMO *et al.*, 2000). Such studies were the departure point for the ephemeral rivers project that established similar monitoring points in other ephemeral rivers throughout north-western Namibia during 1994 (JACOBSON *et al.*, 1995), and these are periodically revisited. This, in turn, has given rise to a more detailed study of the Hoanib River Catchment (LEGGETT, 1998, 2000). This scientific information has been transferred into public documents, including videos and maps (e.g., JACOBSON *et al.*, 1995; HEYNS *et al.*, 1998). These materials address issues related to the use of natural resources, explaining biodiversity and other important ecological concepts to laymen.

Plants

Welwitschia - Long-lived plants are affected by long-term processes and long-term study is required to understand these. *Welwitschia mirabilis* is a widespread endemic of the central and northern Namib (BORNMAN, 1978). In some areas it is the dominant perennial plant present and may serve as shelter or food source for various animals. As the only member of the family Welwitschiaceae, and a most unusual desert plant, the *Welwitschia* is a botanical curiosity and is protected by national legislation. MARSH (1982, 1987, 1990) and BRINCKMAN and VON WILLERT (1987) emphasised that it is important to know more about the phenology of this plant and its satellite fauna in order to improve the management of *Welwitschia* fields. Accordingly, the DRFN initiated a monitoring programme in 1985 in which the degree of leaf growth of 11 male and 10 female plants is measured at monthly intervals. Seasonal patterns are correlated with air humidity, while annual differences are affected by rainfall, but fog does not influence growth patterns and germination (HENSCHHEL and SEELY, in press). The continuing study elucidates the reproductive output, seed dispersal, recruitment, water availability, age structure, and ecological differences between the sexes, and long-term life history strategies. Because of its stark differences to other plants and clear responses to environmental conditions, *Welwitschia* has high demonstration value for students.

!Nara - Another long-lived plant is being monitored, namely, *!nara Acanthosicyos horridus* (Cucurbitaceae), a potential keystone species of the Namib dunes (KLOPATEK and STOCK, 1994). *!Nara* appears to be affected by various environmental factors, such as climate, ground-water availability, herbivory and fruit harvesting, while plant condition may affect numerous invertebrates and vertebrates that feed on or shelter in it. *!Nara* fruits are used by the local Topnaar people as food, fodder, for medical purposes, and as a cash crop under a unique form of tenure (DENTLINGER, 1977). The Topnaar community requested advice from the DRFN on sustainable resource management of the *!nara*. Subsequent investigations have involved both participatory and academic research, with some of the latter being conducted by students and in-service trainees. Unpublished recent studies by DRFN and associates concern the socio-economic importance and marketing of products (BÜTTENDORE, 1999), the plant's water sources, requirements and transpiration, photosynthesis, pollination, seed dispersal, germination and seedling establishment, herbivory, and the importance of *!nara* for animal ecology. The *!nara* is an excellent training tool as it enables students to conduct a wide variety of valuable exercises, ranging from biophysical factors, to ecology and resource management.

Grass - Grass is a major source of primary productivity and detritus and forms the basis of a major food web. For example, SEELY (1990b) described the long-term development of hummocks of the perennial dune grass *Stipagrostis sabulicola*, which are being monitored at intervals of several years since 1976. Ephemeral grasses germinate and can complete their life cycle after a minimum rainfall event of 10-12 mm (SEELY, 1978a, b; JACOBSON, 1992; GÜNSTER, 1993) and the biomass increases with increasing rainfall. In most years since 1988, the annual distribution of ephemeral grasses have been monitored in the central Namib (GÜNSTER, 1995) and was found to be highly variable over space due to patchy rainfall (BURKE, 1997). This provides an excellent opportunity to elucidate the connection between spatial heterogeneity and biodiversity. Establishing this relationship would

increase our endeavour to understand biodiversity in variable environments throughout Namibia (BARNARD, 1998a, b).

Plant communities - Another important LTER project entails the re-examination of all plant communities on the gravel plains of the southern Central Namib (HACHFELD, pers.comm.) over 20 years after ROBINSON's (1976) study. By comparing the spatial distribution of various plant species and of the species composition of various communities, long-term changes in community development can be determined. Furthermore, this study is providing an invaluable baseline for understanding plant biogeography (and associated fauna; e.g., WHARTON, 1980) in the Namib (HACHFELD, 1996). Similarly, re-examining mycorrhiza and lichens at existing study sites (DANEEL, 1992a; SCHIEFERSTEIN and LORIS, 1992; JACOBSON, 1997a), provide important insights into the dynamics of biotic components of the soil surface, particularly in the dunes and on the gravel plains.

On long-term monitoring plots established on three farms in southern Kunene region along gradients of different land-use intensities and different types of land tenure (see above) various vegetation parameters are monitored. These include plant species composition and annual net primary production. These are commonly used indicators of range condition (BEHNKE *et al.*, 1994) and are included in an index of biological integrity developed for Namibian rangelands by DRFN (ZEIDLER *et al.*, 1998; BARNARD *et al.*, 1999; ZEIDLER, 1999). Plants are the key to people's perceptions of their own land and affect their decisions on land use. Perceptions are therefore directly incorporated into the monitoring (ZEIDLER *et al.*, 2000).

Photography - Fixed point photography is a convenient and objective method of monitoring changes of vegetation over seasons, years and decades. This technique lends itself to qualitative and quantitative analyses (e.g. identity, number and size of plants and their relative cover). The DRFN has established fifty fixed points in the Namib and along a transect between Gobabeb and Windhoek and another similar set in southern Kunene region at all NAPCOD sites. These photographs lend themselves to very broad application, and form historical records. For example, ROHDE (1997a, b) used repeat photography, retaking of historical photographs, to identify changes in vegetation over decades to a century.

Animals

Large mammals - Knowledge of the long-term population dynamics and movements of large mammals is very important for their effective management. The Ministry of Environment and Tourism, for example, found that the distribution of gemsbok was primarily determined by the availability of water, whereas grazing affected abundance (BERRY and SIEGFRIED, 1991; KILIAN, 1995; KOK and NEL, 1996), as established through longer-term data records (NEL, 1983). It is important to continue this monitoring and it is suggested to include mountain zebra into the monitoring, as the Namib-Naukluft Park is the distribution centre of this endemic species (JOUBERT, 1974). Troops of baboons living along the Kuiseb River are being monitored at regular intervals since 1970 (HAMILTON *et al.*, 1976; HAMILTON, 1985, 1986; BRAIN, 1992) and in the Swakop River since 1990 (COWLISHAW,

1999). For baboons water is a principal limiting resource and its availability affects reproduction, infant mortality, disease, and inter-troop relationships in troops living at the most arid extreme of the species. Domestic animals, particularly donkeys and goats, are common in and near the lower Kuiseb riverbed. Their changing populations may affect indigenous species, including plants, game, and invertebrates (GABRIEL, 1992; ZEIDLER, 1995). Outside the Kuiseb catchment, there are numerous other studies of Namib mammals that are continued or repeated on a long-term basis, including in communal areas. An example involving DRFN is LEGGETT's (1998, 2000) and Save the Rhino Trust's extension of elephant monitoring previously conducted by VILJOEN (1988).

Invertebrates and small vertebrates - A major contribution to the long-term account of Namib Desert animal populations is the intensive pit-trapping programme of DRFN. Although it focuses primarily on the monitoring of Tenebrionid beetles as "key organisms", it also provides a model for other desert organisms. The concepts that are derived in this study underpin many biodiversity studies undertaken in hyper-arid and arid environments, specifically Namibia. Because of its significance, the tenebrionid long-term research will be dealt with in some detail below. However, the long-term pit-trapping programme is also monitoring many other animals besides tenebrionids in the Namib, e.g. other beetles, solifugids, and lizards. The repeated monitoring technique has been used for the lizard *Meroles anchietae* (ROBINSON, 1990; MUTH, pers.comm.), golden mole *Eremitalpa granti* (FIELDEN, 1989; SEYMOUR and SEELY, 1996) and gerbils (DICKMAN, pers.comm.). These small vertebrates appear to track long-term trends in rainfall. Ants (MARSH, 1986) and soil microarthropods (ANDRÉ *et al.*, 1997) are also good candidates for repeated monitoring due to their importance. Annual monitoring of three spider species since 1987 is revealing how their populations relate to different aspects of the environment, namely food (mainly tenebrionids and ants), and substrate instability, e.g., due to sand storms or trampling by ungulates. Long-term monitoring and modelling demonstrate how these factors affect clusters of spiders and how clusters develop, shift, and decline over generations (HENSCHEL, 1990, 1995; HENSCHEL and LUBIN, 1992; EISINGER *et al.*, 1998; HENSCHEL, 1997). Such work also serves as a model for other animal populations surviving in extreme and variable environments. These invertebrates and small vertebrates provide substance to several kinds of field ecology courses at Gobabeb.

At the DRFN research sites in the farming areas in southern Kunene region pit-trapping studies of tenebrionid beetles (see below) as well as monitoring of biodiversity of abundant representatives of the soil fauna, namely termites (Isoptera), were recently established. Working over most of Namibia, ZEIDLER (1997) analysed termite communities throughout Namibia 30 years after COATON and SHEASBY (1972) had done so. Further current studies are directed at elucidating some underlying reasons for differences in termite communities, such as changes in rainfall and land use patterns (e.g. ZEIDLER, 1999). The biodiversity measures of the termites are related to the soil processes and resources affected by them.


Tenebrionid Beetles - The long-term pit-trapping project that focuses on tenebrionid beetles (Coleoptera) in the Namib serves as an example of the DRFN marrying findings from Namib-LTER with case

studies in farming-LTERs. Tenebrionids are good indicators of environmental conditions because their populations integrate several environmental factors, namely, detritus, leaves and dung on which they feed, vegetation cover under which they shelter, the hardness, moisture and stability of the soil, and the availability of water from rain, fog, and runoff. Furthermore, tenebrionids are abundant, conspicuous, diverse, flightless, and easy to capture and identify. Taxonomic, ecological, behavioural and ecophysiological studies have devoted much attention to Namib Tenebrionidae (more than 180 published papers). These factors make these beetles excellent subjects for further in-depth research and for environmental education.

In an ongoing study initiated in 1968, we are investigating the long-term population dynamics and the species composition of tenebrionid beetles in six habitats near Gobabeb, namely the Kuiseb riverbed, gravel plains, the interdune plains, dune hummocks, dune slope and dune slipface (HENSCHEL, 1994; HENSCHEL, SEELY and POLIS, 1998). Pit traps are operated at weekly to bimonthly intervals. Various diversity indices are derived from the pit trap data and population trends are revealed by comparing the time sequences of abundance data for each species. Different population trends (increasing / decreasing, rapid / slow, abundant / rare / absent) characterise changes in species composition and diversity. Possible causal factors, such as climate, community interactions, habitat characteristics, and periodicity are examined by correlation and autocorrelation.

The extraordinary high diversity of Namib tenebrionids (total >200) has attracted much attention (e.g., KOCH, 1950, 1962; ENDRÖDY-YOUNGA, 1982; PENRITH, 1986; IRISH, 1990; HENSCHEL, SEELY and POLIS, 1998). Most of the 82 species found near Gobabeb are endemic to the Namib. To date, we have live-trapped 460 000 tenebrionids in the 24-year study period since the studies of LOUW and SEELY (1980), WHARTON and SEELY (1982) and SEELY (unpubl.) initiated the project. We found that abundance is highest in those habitats where detritus is richest (riverbed and slipface), while diversity is relatively high in the habitat where food resources fluctuated most strongly (gravel plains). The time series for various species at first appear to be chaotic (e.g., Fig.2) and difficult to interpret (ZHOU *et al.*, 1997), but some patterns emerge upon closer examination (WOLDA, 1978; THOMAS, 1996). Many of these patterns seem to be related to water availability. Rainfall, fog and river floods appear to be important for population growth of most (but not all) Namib tenebrionid species, but the species differed in the type of water source to which they responded as well as in response rate ("response" being defined as an annual increase in population) (e.g., Fig.2). Population decline after growth also differed between species. These differences may explain coexistence of so many species that use a common food source.

Figure 2:



Long-term data series at an interdune LTER site in the Namib dunes of (top) log abundance of the ten most common tenebrionid beetles and (bottom) effective rainfall (>10mm per week) during summer or winter. Compare date from the shaded areas as examples of short-term studies within long-term trends.

We found that during the years between effective rainfalls at Gobabeb (enough to enable plants to germinate and to produce seeds), Namib tenebrionid populations underwent substantial decline (e.g., Fig. 1). Only few small areas appeared to remain relatively rich in species and biomass. By placing pit traps in and next to these rich patches, we are gaining knowledge of their dynamics (e.g., what factors affect their richness) and how these areas relate to other rich areas (metapopulations) and to poor areas between the rich patches. By examining long-term changes we endeavour to improve our understanding of biodiversity and other ecological processes and to improve the management of source areas. Consequently, the Namib tenebrionid LTER includes areas that can be compared as possible sources and sinks, such as dune slopes (near Kahani and at Khommabes located 10 km SW and 10 km W of Gobabeb respectively) with different quantities of perennial vegetation such as hummocks of the grass, *Stipagrostis sabulicola*, and !nara, *Acanthosicyos horridus*.

The long-term data series on tenebrionid populations demonstrate how LTER can form a valuable backdrop to the interpretation of short-term studies (Fig. 2). Beetle community composition appeared to differ in relation to food, soil, vegetation cover, water availability and climatic variability. Some of these factors are susceptible to local change by people, and this forms part of the study at Olifantputs. There we are examining how potential source areas of tenebrionids and termites (e.g., areas where land use by livestock is light) can serve as potential areas to recolonise disturbed areas (e.g., areas heavily grazed and trampled by livestock). This demonstrates one way by which the knowledge gained in the Namib LTER can be applied in communal farming areas such as Olifantputs (PARENZEE, 2000). Here, pit trapping and other monitoring research activities of tenebrionid and termite biodiversity have been initiated to investigate these organisms' potential as indicators of biological integrity at sites with different levels of land use intensity (BARNARD *et al.*, 1999; PARENZEE *et al.*, 2000; ZEIDLER *et al.*, 2000).

People and Livestock

Livestock - At the farming LTER sites the monitoring of domestic animals plays a particularly important role. In order to establish land use histories and long-term records of land use intensity, livestock and domestic animal demographics need to be constructed (ZEIDLER, 1999). In the past the agricultural extension officers have been collecting data which are integrated into the DRFN databases that include our current data and information on historic trends obtained from the rural community members, e.g. by Participatory or Rapid Rural Appraisal (PRA and RRA) methods (ZEIDLER, 1999). The data set should also incorporate data on fodder subsidies during periods of drought ("drought relief") and their effects on the numbers of livestock that are being kept beyond the numbers that can be sustained by the natural grazing available (HENSCHEL, 1996).

People - Long-term data on rural populations assist the interpretation of the interactions of people with the natural environment and the management of natural resources. At the NAPCOD study sites in southern Kunene region baseline data on human demography have been recorded using PRA and RRA methods (ZEIDLER, 1999). Long-term records include human demography and the use of natural resources, e.g. borehole water and grazing. Information on village populations and migration patterns to emergency grazing areas are being incorporated into this monitoring programme. This information is being compared to our concurrent data on environmental indicators, such as vegetation, termites, tenebrionids and soil characteristics. The latter, in turn, is compared to the Namib LTER data. A similar approach is being used to monitor natural resource management by the Topnaar community along the Kuiseb River, particularly concerning !Nara, water and livestock (BOTELLE and KOLWALSKI, 1995; HENSCHEL, MTULENI *et al.*, 1998; SHILOMOLENI, 1998; BÜTTENDORF and HENSCHEL, 1999; AMOOMO *et al.*, 2000).

Using and Applying LTER Data

Various examples of how DRFN is using, combining and applying LTER were given throughout the paper. This section describes the general research process of the DRFN as well as the outputs that are geared towards increasing the awareness and understanding of arid lands, including its biodiversity (see also SEELY *et al.*, submitted).

First, many short term research results, particularly those that feature aspects of biodiversity in arid environments, can be interpreted against the background of long-term trends. This is of great value for studies conducted throughout arid Namibia, but also when looking at arid land ecology worldwide.

Second, it is a policy of the DRFN to make its LTER data available under given conditions (DRFN, 2000). LTER data are housed in a central place, advertised and made accessible for further use. Sharing of data is of fundamental importance for involvement with collaborators, including the Na-LTER and ILTER networks (Na-LTER, 2000).

Third, data and tested methods of the Gobabeb LTER serves as baseline or comparative information

for many other studies and applied programmes in arid and semi-arid parts of Namibia. Some examples were given above in relation to NAPCOD. Further examples relating to the goal of reducing people's vulnerability to loss of sustainable livelihoods in variable environments relate to characterising the state of the environment and identifying vulnerability (e.g., QUAN *et al.*, 1994; SEELY and JACOBSON, 1994; BEHNKE and SCOONES, 1992; JACOBSON, 1997b; WARD and NGAIRORUE, in press). The characterisation through LTER of natural variability is crucial for distinguishing apparent from real environmental changes (e.g., SULLIVAN and KONSTANT, 1997; ROHDE, 1997a; SULLIVAN, 1998, 1999; WARD *et al.*, 1998). Furthermore, LTER provides details of mechanisms underlying long-term changes identified, for example, in archaeological studies (e.g., SANDELOWSKY, 1976; KINAHAN, 1991; SCOTT, 1995) and palaeoclimatology (e.g., HEINE, 1998; LANCASTER, 1998). In this way, LTER effectively complements other research fields concerned with characterising environmental changes.

Fourth, information available is incorporated into text books on desert ecology, resource materials for school teachers, natural history overviews, nature films, coffee table books, and information packages for resource managers and policy makers (described by SEELY *et al.*, this volume, submitted).

Fifth, national resource management schemes and policy-making projects require long-term data. Prominent among these in Namibia the Framework Convention on Climate Change (FCCC), the Convention of Biological Diversity (CBD) and the Convention to Combat Desertification (CCD), and the National Drought Policy. State of the Environment Reports and formal Workshops serve to provide feedback from Namibian and international forums (SEELY *et al.*, submitted).

Sixth, the most important point, all DRFN research is connected to an explicit training component involving students as well as natural resource users and managers in the collection and interpretation of the data. Such a participatory research process contributes to the capacity building of natural resource users and managers. Rural resource users are directly involved in research on natural resources so that they themselves can provide effective feedback towards conclusions. This facilitates devising nation-wide mechanisms for sustainable resource management and environmental monitoring. The NAPCOD programme serves as leading example of this connection.

General Discussion

Biodiversity reflects a combination of environmental and historical events at a site and changes in biodiversity can provide a sensitive measure of ecologically relevant changes in the environment. LTER can provide historical data, while allied studies focus on the proximate relationship between organisms (including people) and the environment (RISSER, 1995; BARNARD, 1998a).

The Gobabeb and other LTER sites described here are part of a larger network of LTER sites concerning arid lands. Prominent among these are further NAPCOD sites that include old SARDEP sites, the Kalahari Transect covering parts of South Africa, Namibia, Botswana and Zambia (SCHOLES and PARSONS, 1997), the planned BIOTA African Transcontinental Transect (JÜRGENS, pers.comm.)

and transects covering other deserts of the world (KOCH *et al.*, 1995). These and other initiatives are interconnected via the Na-LTER network (Na-LTER, 2000) and through collaboration along several other avenues. It is anticipated that a regional SADC network of LTER sites may arise within the ILTER network (ILTER, 1999) and that the Na-LTER sites will contribute to these.

Continuity and consistency in methods are required for LTER. However, LTER should be more than generating and interpreting data and it ultimately needs to involve human communities. The Gobabeb LTER projects demonstrate that baseline information and understanding derived from an undisturbed, arid environment is invaluable to the interpretation of shorter-term data collected in other arid areas. This should ultimately enable brief assessments of biodiversity to be used to describe habitat quality. Likewise, it allows an interpretation of the effects of people on biodiversity and facilitates the planning of projects in populated areas that may change these effects.

Biophysical and socio-economic LTER projects are extremely important for the Namibian environment. This has long been recognised by several other organisations besides the DRFN, e.g. the Ministry of Agriculture, Water and Rural Development, and the Ministry of Environment and Tourism, as well as other NGO's like the Save the Rhino Trust (over 20 institutions have thus far declared interest in Na-LTER). However, today it becomes more and more difficult to maintain long-term sites. There appear to be several reasons. First, the maintenance of long-term environmental research is costly and demands much dedication. Second, the value of long-term data collection is often not fully recognised. This may partly be because LTER data and conclusions are sometimes not published appropriately and timely nor made accessible. The current paper demonstrates the value of published LTER data for many social and environmental fields in Namibia.

Conclusions

In this paper we show how LTER has developed at Gobabeb since Independence, including generating, sharing, understanding, using and applying research results for training and management purposes. LTER is important for arid land studies. Its ultimate value depends on the alliance between basic research and applied research that underlines the importance of both. In the long run both types of research cannot do without each other. Basic research is vital because it provides first-hand knowledge that can be translated and applied into relevant information to relevant target groups. Basic research often comes up with novel insights that were not initially predicted. Knowledge can in this way advance in quantum leaps. However, without its application, basic research can develop into an ivory-tower condition that a developing country can ill afford. It is the fruitful interaction between basic and applied research that increase the value of both. Well-tested conclusions and honed skills can be transferred from basic to applied research, while applied questions form a guiding framework for basic research. LTER embraces both.

Acknowledgements

The Ministry of Environment and Tourism granted permission for LTER at Gobabeb in the Namib-Naukluft Park. The National Museum of Namibia played an important supporting role over the years.

Technical assistants at Gobabeb were ANGELA SUTTLE, LINDA MALAN, CHRISTINE HÄNEL and VILHO MTULENI and these were assisted by numerous other staff, research associates, and volunteers. The participatory research approach at Olifantputs involved numerous enthusiastic members of the community, and, in particular, we would like to thank DUDU MURORUA, LESLEY PARENZEE and DEON SHARURU for assistance. BARBARA PATERSON kindly assisted with the development of a computerised management information system. KATHY JACOBSON, MARK ROBERTSON and DAVID WARD commented on the manuscript.

References

- AMOOMO, H., GASEB, N., HOVEKA, V., MBANGULA, E., MUHARUKUA, E. and ZAARUKA, B. 2000. *Determining the water reserve of the Kuiseb River - sustainable use of Namibia's ephemeral rivers, life-lines for development*. Windhoek: Desert Research Foundation of Namibia, Summer Desertification Programme 8.
- ANDERSON, J.M. 1994. Functional attributes of biodiversity in land use systems. In: GREENLAND, D.J. and SZABOLCS, I. (Eds.), *Soil resilience and sustainable land use*, pp. 267-290. Oxon, UK: CAB International.
- ANDRÉ, H.M., NOTI, M.-I. and JACOBSON, K.M. 1997. The soil microarthropods of the Namib Desert: a patchy mosaic. *Journal of African Zoology* 111: 499-517.
- BARNARD, P. 1998a. *Biological diversity in Namibia, a country study*. Windhoek: Namibian National Biodiversity Task Force, Directorate of Environmental Affairs.
- BARNARD, P. (Ed.) 1998b. Special issue: The biological diversity of Namibia (ten papers by several authors). *Biodiversity & Conservation* 7: 415-559.
- BARNARD, P., ROBERTSON, M., ZEIDLER, J. 1999. Abstract. Developing an early warning system for environmental degradation in Namibia. *Vlth International Rangeland Congress Proceedings, Townsville, Australia* 2: 662-663.
- BEHNKE, R. H., and SCOONES, I. 1992. Rethinking range ecology: implications for rangeland management in Africa. *International Institute for Environment and Development, Overseas Development Institute, London* 33: 1-43.
- BEHNKE, R.H., SCOONES, I., and KERVEN, C. (Eds.). 1994. *Range ecology at disequilibrium*. London: Overseas Development Institute. 248 pp.
- BENDER, K. (Ed.) 1999. *Coastal profile of the Erongo Region*. Swakopmund, Namibia: Integrated Coastal Zone Management Project, Erongo Regional Council. 214 pp.
- BERRY, H.H. and SIEGFRIED, W.R. 1991. Mosaic-like events in arid and semi-arid Namibia. In: REMMERT, H. (Ed.), *The mosaic-cycle concept of ecosystems*, pp. 147-160. Berlin: Springer.
- BESLER, H. 1970. Geomorphologie der Wüste. *Namib und Meer* 1: 59-68.
- BESLER, H. 1972a. Klimaverhältnisse und klimageomorphologische Zonierung der zentralen Namib (Südwestafrika). *Stuttgarter Geographische Studien* 83: 1-209.
- BESLER, H. 1972b. Geomorphologie der Dünen. *Namib und Meer* 3: 25-35.
- BORNMAN, C.H. 1978. *Welwitschia: paradox of a parched paradise*. Struik Publishers, Cape Town, 71 pp.

- BOSS, G. 1941. Niederschlagsmenge und Salzgehalt des Nebelwassers an der Küste Deutsch-Südwestafrikas. *Bioklimatische Beiblätter der Meteorologischen Zeitschrift* 8(1): 1-14.
- BOTELLE, A. and KOWALSKI, K. 1995. *Changing resource use in Namibia's lower Kuiseb river valley: perceptions from the Topnaar community*. Maseru & Windhoek: Institute of Southern African Studies at the University of Lesotho and the Social Sciences Division at the University of Namibia. 90 pp.
- BOYER, D.C. 1989. Some characteristics of the plant communities of three dunes situated across a climatic gradient in the Namib Desert. *Madoqua* 16: 141-148.
- BRAIN, C. 1992. Deaths in a desert baboon troop. *International Journal of Primatology* 13: 593-599.
- BRINCKMANN, E., and VON WILLERT, D.J. 1987. Injury and recovery of *Welwitschia mirabilis*. *Dinteria* 19: 69-76.
- BURKE, A. 1997. Is the grass always greener at certain sites? A contribution to patch dynamics in the central Namib desert. *Tropical Ecology* 38: 125-127.
- BÜTTENDORF, U. 1999. *Former and current !Nara utilisation and its potential for development, an interim report. Gobabeb, Namibia*: Desert Research Foundation of Namibia.
- COATON, W.G.H., and SHEASBY, J.L. 1972. Preliminary report on a survey of the termites (Isoptera) of South West Africa. *Cimbebasia Memoir* 2: 7-129.
- COWLISHAW, G. 1999. Ecological and social determinants of spacing behaviour in desert baboon groups. *Behavioural Ecology and Sociobiology* 45: 67-77.
- CRAWFORD, C.S., and GOSZ, J.R. 1982. Desert ecosystems: their resources in space and time. *Environmental Conservation* 9: 181-195.
- DANEEL, J.L. 1992a. *The impact of off-road vehicle traffic on the gravel plains of the Central Namib Desert, Namibia*. M.Sc. thesis, University of Natal, Pietermaritzburg, South Africa. 117 pp.
- DANEEL, J.L. 1992b. The visual effects of off-road vehicles on the gravel plains of the Central Namib. Gobabeb, Namibia: Desert Ecological Research Unit. 14 pp.
- DAUSAB, F., FRANCIS, G., JÖHR, G., KAMBATUKU, J., MOLAPO, M., SHANYENGANA, E. and SWARTZ, S. 1994. *Water usage patterns in the Kuiseb catchment area - with emphasis on sustainable use*. Windhoek: Desert Research Foundation of Namibia, Summer Desertification Programme 2, Occasional Paper 1. 243 pp.
- DEAN, W.R.J. and MILTON, S.J. (Eds.) 1999. *The Karoo, ecological patterns and processes*. Cambridge: Cambridge University Press.
- DENTLINGER, U. 1977. An ethnobotanical study of the !Nara plant among the Topnaar hottentots of Namibia. *Munger Africana Library Notes* 38: 3-39.
- DIAS, P.C. 1996. Sources and sinks in population biology. *Trends in Ecology and Evolution* 11: 326-330.
- DRFN, 2000. *Data sharing policy of the DRFN*. Windhoek: Desert Research Foundation of Namibia. 2 pp.
- ECKARDT, F. 1996. *The distribution and origin of gypsum in the Central Namib Desert, Namibia*. Ph.D. thesis, University of Oxford, Hertford College, Oxford, U.K. 320 pp.

- EISINGER, D., JELTSCH, F., HENSCHER, J., ULBRICH, K., LUBIN, Y. and WISSEL, C. 1998. Das lokale Ausbreitungsmuster der Wüstenspinne *Seothyra henscheli* - ein räumlich-explizites gitterbasiertes Simulationsmodell. (The local dispersal pattern of the desert spider *Seothyra henscheli* — a spatially-explicit grid-based simulation model.) *Verhandlungen der Gesellschaft für Ökologie* 28: 141-150.
- ENDRÖDY-YOUNGA, S. 1982. Dispersion and translocation of dune specialist tenebrionids in the Namib area. *Cimbebasia* (A) 5: 257-271.
- FIELDEN, L. J. 1989. *Selected aspects of the adaptive biology and ecology of the Namib Desert golden mole (Eremitalpa granti namibensis)*. Ph.D. thesis, University of Natal, Pietermaritzburg, South Africa. 247 pp.
- FRANKLIN, J.F., BLEDSOE, C.S. and CALLAHAN, J.T. 1990. Contributions of the long-term ecological research program. *BioScience* 40: 509-523.
- GABRIEL, M. 1992. *The impact of stock along a defined stretch of the Lower Kuiseb River, Central Namib Desert: a first attempt at quantification. Gobabeb, Namibia: Desert Research Foundation of Namibia.*
- GREENLAND, D.J. and SZABOLCS, I. (Eds.). 1994. *Soil resilience and sustainable land use*. Oxon, U.K.: CAB International.
- GOSZ, J.R. 1996. International long-term ecological research: priorities and opportunities. *Trends in Ecology and Evolution* 11: 444.
- GOUDIE, A. 1972. Climate, weathering, crust formation, dunes, and fluvial features of the central Namib Desert, near Gobabeb, South West Africa. *Madoqua* 1: 15-31.
- GOUDIE, A.S. and PARKER, A.G. 1998. Experimental simulation of rapid rock block disintegration by sodium chloride in a foggy coastal desert. *Journal of Arid Environments* 40: 347-355.
- GOUDIE, A.S., VILES, H.A. and PARKER, A.G. 1997. Monitoring of rapid salt weathering in the central Namib Desert using limestone blocks. *Journal of Arid Environments* 37: 581-598.
- GTOS. 2000. *Global Terrestrial Observation System*. Website: <http://www.fao.org/gtos>.
- GÜNSTER, A. 1993. Seed bank dynamics of serotinous plants in the central Namib. Ph.D. thesis. University of Münster, Münster, Germany. 103 pp.
- GÜNSTER, A. 1995. Grass cover distribution in the central Namib - a rapid method to assess regional and local rainfall patterns of arid regions? *Journal of Arid Environments* 29: 107-114.
- HACHFELD, B. 1996. *Vegetationsökologische Transektanalyse in der nördlichen Zentralen Namib*. M.Sc. thesis, Institut für Allgemeine Botanik, Universität Hamburg, Hamburg, Germany. 137 pp.
- HAMILTON, W.J. 1985. Demographic consequences of a food and water shortage to desert Chacma baboons, *Papio ursinus*. *International Journal of Primatology* 6: 451-462.
- HAMILTON, W.J. 1986. Namib Desert Chacma baboon (*Papio ursinus*) use of food and water resources during a food shortage. *Madoqua* 14: 397-408.
- HAMILTON, W.J., BUSKIRK, R.E. and BUSKIRK, W. H. 1976. Defense of space and resources by Chacma (*Papio ursinus*) baboon troops in an African desert and swamp. *Ecology* 57: 1264-1272.

- HAMILTON, W.J., BUSKIRK, R.E. and BUSKIRK, W.H. 1977. Intersexual dominance and differential mortality of Gemsbok *Oryx gazella* at Namib Desert waterholes. *Madoqua* 10: 5-19.
- HEINE, K. 1998. Klimawandel und Desertifikation im südlichen Afrika - ein Blick in die Zukunft. *Geographische Rundschau* 50: 245-250.
- HENSCHEL, J.R. 1990. The biology of *Leucorchestris arenicola* (Araneae: Heteropodidae), a burrowing spider of the Namib dunes. In: SEELY, M.K. (Ed.), *Namib Ecology - 25 years of Namib research*, pp. 115-127. Pretoria: Transvaal Museum Monograph 7.
- HENSCHEL, J.R. 1994. Pithy pits: population dynamics of Namib tenebrionids. *Namib Bulletin* 11: 4-5.
- HENSCHEL, J.R. 1995. Tool use by spiders: stone selection and placement by corolla spiders *Ariadna* (Segestriidae) of the Namib Desert. *Ethology* 101: 187-199.
- HENSCHEL, J.R. 1996. Subsidisation empoverishes local resources and increases dependency of local consumers: a lesson from spiders and aquatic insects. *Namib Bulletin* 13: 13.
- HENSCHEL, J.R. 1997. Psammophily in Namib Desert spiders. *Journal of Arid Environments* 37: 695-707.
- HENSCHEL, J.R. 1999. Namibia moving towards networking LTER. *Abstract, Annual General Meeting of the International Long Term Ecological Research Network, August 1999, Skukuza, South Africa.*
- HENSCHEL, J.R., BÖHME, E., DENTLINGER, L., HENSCHEL, I.A., MÜLLER, M. and SEELY, M.K. 2000. *DRFN Bibliography 2000*. Windhoek: Desert Research Foundation of Namibia. 86 pp.
- HENSCHEL, J.R. and LUBIN, Y.D. 1992. Environmental factors affecting the web and activity of a psammophilous spider in the Namib Desert. *Journal of Arid Environments* 22: 173-189.
- HENSCHEL, J., MTULENI, V., GRUNKOWSKI, N., SEELY, M. and SHANYENGANA E. 1998. *Namfog, Namibian application of fog-collecting systems, I: Evaluation of fog water harvesting*. Desert Research Foundation of Namibia, Occasional Paper 8: 1-75.
- HENSCHEL, J.R. and SEELY, M.K. in press. Long-term growth patterns of *Welwitschia mirabilis*, a long-lived plant of the Namib Desert (including a bibliography). *Plant Ecology*: in press.
- HENSCHEL, J.R., SEELY, M.K., POLIS, G.A. 1998. Long term population dynamics of tenebrionid beetles in the Namib Desert. *Abstract: INTECOL: VII International Congress of Ecology, July 1998, Florence, Italy.*
- HEYNS, P., MONTGOMERY, S., PALLETT, J. and SEELY, M. (Eds.) 1998. *Namibia's water: a decision makers' guide*. Windhoek, Namibia: The Department of Water Affairs, Ministry of Agriculture, Water and Rural Development, and The Desert Research Foundation of Namibia. 173 pp.
- HOLM, E. 1970. *The influence of climate on the activity patterns and abundance of xerophilous Namib desert dune insects*. M.Sc. thesis, University of Pretoria, Pretoria. 44 pp.
- HUNTLEY, B.J. (ed.). 1985. The Kuiseb environment: the development of a monitoring baseline. *South African National Scientific Programmes Report* 108: 1-138.
- ILTER. 1999. Abstracts: Annual General Meeting of ILTER. *International Long Term Ecological Research Network, August 1999, Skukuza, South Africa.*

- ILTER. 2000. *International Long-Term Ecological Research Network*.
Website: <http://www.ilternet.edu>.
- IRISH, J. 1990. Namib biogeography, as exemplified mainly by the Lepismatidae (Thysanura: Insecta). In: SEELY, M.K. (Ed.), *Namib Ecology - 25 years of Namib research*, pp. 61-66. Pretoria: Transvaal Museum Monograph 7.
- JACOBSON, K.M. 1992. *Factors affecting VA-mycorrhizal community structure in the Namib Dune Field; and the population biology of an ectomycorrhizal Basidiomycete: Suillus granulatus*. Ph.D. thesis, USA: Virginia Polytechnic Institute, Virginia State University, Blacksburg, Virginia.
- JACOBSON, K.M. 1996. *Macrofungal ecology in the Namib desert: a fruitful or futile study?* *McIlvainea* 12: 21-32.
- JACOBSON, K.M. 1997a. Moisture and substrate stability determine VA-mycorrhizal fungal community distribution and structure in an arid grassland. *Journal of Arid Environments* 35: 59-75.
- JACOBSON, K.M. 1997b. An overview of research issues relating to sustainable natural resource management in drylands. Windhoek: Desert Research Foundation of Namibia. 149 pp.
- JACOBSON, K.M. and JACOBSON, P.J. 1998. Rainfall regulates decomposition of buried cellulose in the Namib Desert. *Journal of Arid Environments* 38: 571-584.
- JACOBSON, P.J., JACOBSON, K.M. and SEELY, M.K. 1995. Ephemeral rivers and their catchments: sustaining people and development in western Namibia. Windhoek: Desert Research Foundation of Namibia.
- JELTSCH, F., MILTON, S.J., DEAN, W.R.J., VAN ROOYEN, N. and MOLONEY, K. 1998. Modeling the impact of small-scale heterogeneities on tree-grass co-existence in semi-arid savannas. *Journal of Ecology* 86: 780-793.
- JOUBERT, E. 1974. Notes on the reproduction in Hartmann zebra *Equus zebra hartmannae* in South West Africa. *Madoqua Series I* 8: 31-35.
- KAMBATUKU, J.R. 1996. *Historical profiles of farms in former Damaraland: notes from the archival files*. Windhoek: Desert Research Foundation of Namibia, Occasional Report 4. 58 pp.
- KILIAN, J.W. 1995. *The ecology of gemsbok (Oryx gazella gazella) in the southern Namib*. Windhoek: Directorate of Specialist Support Services, Ministry of Environment and Tourism.
- KINAHAN, J. 1991. *Pastoral Nomads of the central Namib Desert: the people history forgot*. Windhoek: Namibia Archaeological Trust. 167 pp.
- KLOPATEK, J.M. and STOCK, W.D. 1994. Partitioning of nutrients in *Acanthosicyos horridus*, a keystone endemic species in the Namib Desert. *Journal of Arid Environments* 26: 233-240.
- KOCH, C. 1950. The Tenebrionidae of Southern Africa 1: first account of the Tenebrionidae collected on the University of California-Transvaal Museum expedition, 1948. *Annals of the Transvaal Museum* 21: 273-367.
- KOCH, C. 1962. The Tenebrionidae of southern Africa 31: Comprehensive notes on the tenebrionid fauna of the Namib Desert. *Annals of the Transvaal Museum* 24: 61-106.

- KOCH, G.W, SCHOLES, R.J., STEFFEN, W.L., VITOUSEK, P.M. and WALKER, B.H. 1995. The IGBP terrestrial transects: Science plan. *International Geosphere-Biosphere Programme, Stockholm, Report 36*: 1-53.
- KOK, O.B., and NEL, J.A.J. 1996. The Kuiseb river as a linear oasis in the Namib desert. *African Journal of Ecology* 34: 39-47.
- LANCASTER, N. 1989. *The Namib sand sea: dune forms, processes and sediments*. Rotterdam: A.A.Balkema.
- LANCASTER, N. 1998. Dune morphology, chronology, and Quaternary climatic change. In: ALSHARHAN, A.S., GLENNIE, K.W., WHITTLE, G.L., KENDALL, C.G.S., (Eds.), *Quaternary Deserts and Climatic Change*, pp. 339-349. Rotterdam: A.A. Balkema.
- LANCASTER, J., LANCASTER, N. and SEELY, M.K. 1984. Climate of the central Namib Desert. *Madoqua* 14: 5-61.
- LANDRES, P.B., MORGAN, P. and SWANSON, F.J. 1999. Overview of the use of natural variability concepts in managing ecological systems. *Ecological Applications* 9: 1179-1188.
- LEGGETT, K. 1998. *The Hoanib River catchment study: a summary*. Windhoek: Desert Research Foundation of Namibia.
- LEGGETT, K. 2000. The Hoanib River Catchment study. *Namib Bulletin* 15: 22.
- MARSH, A.C. 1986. Checklist, biological notes and distribution of ants in the central Namib Desert. *Madoqua* 14: 333-344.
- MARSH, B.A. 1982. An ecological study of *Welwitschia mirabilis* and its satellite fauna. *Namib Bulletin* 4: 3-4.
- MARSH, B.A. 1987. Micro-arthropods associated with *Welwitschia mirabilis* in the Namib Desert. *South African Journal of Zoology* 22: 89-96.
- MARSH, B.A. 1990. The microenvironment associated with *Welwitschia mirabilis* in the Namib desert. In: SEELY, M.K. (Ed.), *Namib Ecology - 25 years of Namib research*, pp. 149-153. Pretoria: Transvaal Museum Monograph 7.
- MILTON, S.J. 1995. Spatial and temporal patterns in the emergence and survival of seedlings in arid Karoo shrubland. *Journal of Applied Ecology* 32: 145-156.
- Na-LTER. 2000. *Namibian Long-Term Ecological Research Network*.
Website: <http://www.netwise.drfn.org.na/Na-LTER.html>.
- NEL, P.S. 1983. *Monitoring van die beskikbaarheid, gehalte en benutting van voer op die gruisflaktes van die Kuiseb-studiegebied*. M.Sc. thesis, University of the Orange Free State, Bloemfontein, South Africa. 233 pp.
- NICHOLS, N. and WONG, K.K. 1990. Dependence of rainfall variability on mean rainfall, latitude and the Southern Oscillation. *Journal of Climate* 3: 162-170.
- NOY-MEIR, I. 1981. Understanding arid ecosystems: the challenge. In: GOODALL, D.W. and PERRY, R.A. (Eds), *Arid land ecosystems: structure, functioning and management*, pp. 447-449. Cambridge: Cambridge University Press.
- PARENZEE, L. 2000. *Use of tenebrionid beetles as indicators of habitat quality*. M.Sc. thesis, University of Witwatersrand, Johannesburg. submitted.

- PARENZEE, L., ZEIDLER, J. and SEELY, M. 2000. Testing biodiversity indicators for community use - a case study from Namibia. *Abstract, 13th Congress of the German Society for Tropical Ecology, March 2000, Würzburg, Germany.* p. 37.
- PENRITH, M.-L. 1986. Relationships of the tribe Adesmiini (Coleoptera: Tenebrionidae) and a revision of the genus *Stenodesia* Reitter. *Annals of the Transvaal Museum* 34: 275-302.
- PULLIAM, H.R. 1988. Sources, sinks and population regulation. *American Naturalist* 132: 652-661.
- QUAN, J., BARTON, D., CONROY, C. and ASHLEY, C. (1994). A preliminary assessment of the economic impact of desertification in Namibia. *Department of Environmental Affairs Research Discussion Paper* 3: 150 pp.
- REYNOLDS, J.F., VIRGINIA, R.A., KEMP, P.R., DE SOYZA, A.G. and TREMMEL, D.C. 1999. Impact of drought on desert shrubs: effects of seasonality and degree of resource island development. *Ecological Monographs* 69: 69-106.
- RISSER, P.G. 1995. *Long-term ecological research: An international perspective, SCOPE 47.* Chichester, U.K.: John Wiley.
- ROBINSON, E.R. 1976. *Phytosociology of the Namib Desert Park, South West Africa.* M.Sc. thesis, University of Natal, Pietermaritzburg, South Africa. 220 pp.
- ROBINSON, E.R. 1978. Phytogeography of the Namib Desert of South West Africa (Namibia) and its significance to discussions of the age and uniqueness of this desert. In: VAN ZINDEREN BAKKER, E.M. and COETZEE, J.A. (Eds.), *Palaeoecology of Africa and the surrounding islands, Vol. 10*, pp. 67-74. Rotterdam: A.A. Balkema.
- ROBINSON, M. D. 1990. Comments on the reproductive biology of the Namib desert dune lizard, *Aporosaura anchietae*, during two years of very different rainfall. In: SEELY, M.K. (Ed.), *Namib Ecology - 25 years of Namib research*, pp. 163-168. Pretoria: Transvaal Museum Monograph 7.
- ROBINSON, M.D. and SEELY, M.K. 1980. Physical and biotic environments of the southern Namib dune ecosystem. *Journal of Arid Environments* 3: 183-203.
- ROHDE, R.F. 1997a. *Nature, cattle thieves and various other midnight robbers: images of people, place and landscape in Damaraland, Namibia.* Ph.D. thesis, University of Edinburgh, Edinburgh, U.K.
- ROHDE, R.F. 1997b. Looking into the past: interpretations of vegetation change in Western Namibia based on matched photography. *Dinteria* 25: 121-149.
- SANDELOWSKY, B.H. 1976. The beginning of archaeo-ethno-botany in the Namib. In: VAN ZINDEREN BAKKER, E.M. (Ed.), *Palaeoecology of Africa* 9, pp. 136-143. Rotterdam: A.A. Balkema.
- SCHIEFERSTEIN, B. and LORIS, K. 1992. Ecological investigations on lichen fields of the Central Namib. *Vegetatio* 98: 113-128.
- SCHOLES, R.J. and PARSONS, D.A.B. 1998. The Kalahari Transect: research on global change and sustainable development in southern Africa. *International Geosphere-Biosphere Programme, Stockholm, Report* 42: 1-63.

- SCOTT, L. 1995. Pollen evidence for vegetational and climatic change in Southern Africa during the Neogene and Quaternary. In: VRBA, E.S., DENTON, G.H., PARTRIDGE, T.C. and BURCKLE, L.H. (Eds.), *Paleoclimate and Evolution with Emphasis on Human Origins*, pp. 65-76. Yale: Yale University Press.
- SEELY, M.K. 1973. Factors controlling reproduction of certain Namib Desert tenebrionids. *Madoqua*, Series II 2(63-68): 63-65.
- SEELY, M.K. 1978a. Standing crop as an index of precipitation in the central Namib grassland. *Madoqua* 11: 61-68.
- SEELY, M.K. 1978b. Grassland productivity: the desert end of the curve. *South African Journal of Science* 74: 295-297.
- SEELY, M.K. (Ed.) 1990a. *Namib Ecology - 25 years of Namib research*. Pretoria: Transvaal Museum Monograph 7.
- SEELY, M.K. 1990b. Patterns of plant establishment on a linear desert dune. *Israel Journal of Botany* 39: 443-451.
- SEELY, M.K. 1998. Can science and community action connect to combat desertification? *Journal of Arid Environments* 39: 267-277.
- SEELY, M.K., BUSKIRK, W.H. and HAMILTON III, W.J. 1980. Lower Kuiseb river perennial vegetation survey. *Journal of the South West Africa Scientific Society* 35: 57-86.
- SEELY, M.K. and HAMILTON, W.J. 1978. Durability of vehicle tracks on three Namib Desert substrates. *South African Journal of Wildlife Research* 8: 107-111.
- SEELY, M.K., HENSCHHEL, J.R., ZEIDLER, J. and SHANYENGANA, E.S.C. 2000. Namib research: its development at Gobabeb and implications for Namibia. *Journal Namibia Scientific Society* Volume 48.
- SEELY, M.K. and JACOBSON, K.M. 1994. Desertification and Namibia: a perspective. *Journal of African Zoology* 108: 21-36.
- SEELY, M.K. and LOUW, G.N. 1980. First approximation of the effects of rainfall on the ecology and energetics of a Namib Desert dune ecosystem. *Journal of Arid Environments* 3: 25-54.
- SEELY, M.K., ZEIDLER, J., HENSCHHEL, J.R., BARNARD, P. submitted. Creative problem solving in support of biodiversity conservation. *Journal of Arid Environments*.
- SEYMOUR, R.S. and SEELY, M.K. 1996. The respiratory environment of the Namib Desert golden mole. *Journal of Arid Environments* 32: 453-461.
- SHANYENGANA, E.S. 1997a. Baseline study: Gobabeb Training and Research Centre programme: appropriate technology for arid lands. Windhoek: Desert Research Foundation of Namibia. 56 pp.
- SHANYENGANA, E.S. 1997b. Appropriate technology at Gobabeb: a look into the future. *Namib Bulletin* 14: 18.
- SHILOMBOLENI, A. 1998. The !nara and factors that lead to its decline in productivity. Third year Diploma in-service training thesis, Department of Natural Resource Management of the Polytechnic of Namibia and Desert Research Foundation of Namibia, Windhoek. 17 pp.
- SOUTHGATE, R.I., MASTERS, P., SEELY, M.K. 1996. Precipitation and biomass changes in the Namib Desert dune ecosystem. *Journal of Arid Environments* 33: 267-280.

- SULLIVAN, S. 1998. People, plants and practice in drylands: socio-political and ecological dimensions of resource-use by Damara farmers in north-west Namibia. Ph.D. thesis, University College London, London. 448 pp.
- SULLIVAN, S. 1999. The impacts of people and livestock on topographically diverse open wood- and shrub-lands in arid north-west Namibia. *Global Ecology and Biogeography Letters* 8: 257-277.
- SULLIVAN, S. and KONSTANT, T.L. 1997. Human impacts on woody vegetation, and multivariate analysis: a case study based on data from Khowarib settlement, Kunene Region. *Dinteria* 25: 87-120.
- THERON, G.K. and VAN ROOYEN, M.W. 1980. Vegetation of the lower Kuiseb River. *Madoqua* 11: 327-345.
- THOMAS, L. 1996. Monitoring long-term population change: why are there so many analysis methods. *Ecology* 77: 49-58.
- TILSON, R.L. and HENSCHER, J.R. 1986. Spatial arrangement of spotted hyaena groups in a desert environment, Namibia. *African Journal of Ecology* 24: 173-180.
- TYSON, P.D. 1986. Climatic change and variability in southern Africa. Oxford, U.K.: Oxford University Press.
- TYSON, P.D. and DYER, T.J. 1975. Mean annual fluctuations of precipitation in the summer rainfall region of South Africa. *South African Geographical Journal* 57: 104-110.
- UNEP, 1995. Convention to Combat Desertification. United Nations Environmental Programme, Geneva, Switzerland.
- VILJOEN, P.J. 1988. The ecology of the desert-dwelling elephants *Loxodonta Africana* (Blumenbach, 1797) of western Damaraland and Kaokoland. Ph.D. thesis, University of Pretoria, Pretoria. 335 pp.
- WALTER, H. 1937. Die ökologischen Verhältnisse in der Namib-Nebelwüste (Südwestafrika). *Jahrbucher für wissenschaftliche Botanik* 84: 58-222.
- WALTER, H. 1971. Ecology of tropical and subtropical vegetation. Edinburgh, U.K.: Oliver and Boyd.
- WARD, D. and ELLIS, J. 1999. Abstract: The effect of grazing on biodiversity in arid lands. In: GOSZ, J.R. and SHACHAK, M. (Eds.), Workshop on Biodiversity in drylands: towards a unified framework and identification of research needs, June 1999. Mitrani Department of Desert Ecology, Ben-Gurion University of the Negeve, Beer Sheva and Sde Boqer, Israel.
- WARD, D. and NGAIRORUE, B.T. in press. Are Namibia's grasslands desertifying? *Journal of Range Management*.
- WARD, D., NGAIRORUE, B.T., KATHENA, J., SAMUELS, R. and OFRAN, Y. 1998. Land degradation is not a necessary outcome of communal pastoralism in arid Namibia. *Journal of Arid Environments* 40: 357-371.
- WATKINSON, A.R. and SUTHERLAND, W.J., 1995. Sources, sinks and pseudo-sinks. *Journal of Animal Ecology* 64: 126-130.
- WEATHERHEAD, P. J. 1986. How unusual are unusual events? *American Naturalist* 128: 150-154.

- WHARTON, R.A. 1980. Insects and arachnids associated with *Zygophyllum simplex* (Zygophyllaceae) in the central Namib Desert. *Madoqua* 12: 131-139.
- WHARTON, R.A. and SEELY, M.K. 1982. Species composition of and biological notes on Tenebrionidae of the lower Kuiseb River and adjacent gravel plain. *Madoqua* 13: 5-25.
- WOLDA, H. 1978. Fluctuations in abundance of tropical insects. *American Naturalist* 112: 1017-1045.
- YEATON, R.I. 1988. Structure and function of the Namib dune grasslands: Characteristics of the environmental gradients and species distributions. *Journal of Ecology* 76: 744-758.
- ZEIDLER, J. 1995. Spatial and temporal abundance of *Rhipicephalus gertrudae* (Acari, Ixodidae) along a degradation gradient along the Lower Kuiseb river, Namibia. M.Sc. thesis, Johann-Wolfgang Goethe University, Frankfurt-am-Main, Germany. 55 pp.
- ZEIDLER, J. 1997. Distribution of termites (Isoptera) throughout Namibia - environmental connections. M.Sc. thesis, University of Witwatersrand, Johannesburg. 91 pp.
- ZEIDLER, J., 1999. Establishing indicators of biological integrity for western Namibian rangelands. Ph.D. thesis, University of the Witwatersrand, Johannesburg. submitted.
- ZEIDLER, J. 2000. Linking the environment to rural development - Napcod III. *Namib Bulletin* 15: 9.
- ZEIDLER, J., SEELY, M., HANRAHAN, S. and SCHOLE, M. 1998. Abstract: Establishing indicators of biological integrity in western Namibian rangelands: combining an ecological with a participatory community-based research approach. *Proceedings of the Kalahari Transect Meeting, Gaborone, Botswana, 10-13 June 1998, START report No. 2.*
- ZEIDLER, J., SEELY, M. and PARENZEE, L. 2000. Environmental indicators for community management. Abstracts of the 13th Congress of the German Society for Tropical Ecology, March 2000, Würzburg, Germany. p. 53.
- ZHOU, X., PERRY, J.N., WOIWOD, I.P., HARRINGTON, R., BALE, J.S., and CLARK, S.J. 1997. Detecting chaotic dynamics of insect populations from long-term survey data. *Ecological Entomology* 22: 231-241.

Personal Communication

- COETZEE, Martin, and MULDER, Karl-Heinz - Fog Project, Swakopmund, Namibia.
- DICKMAN, Chris - School of Biological Sciences, University of Sydney, Sydney, Australia.
- HACHFELD, Berit - Botanical Institute, University of Cologne, Cologne, Germany.
- JÜRGENS, Norbert - Botanical Institute, University of Cologne, Cologne, Germany.
- MUTH, Allan - Boyd Deep Canyon Desert Research Center, Palm Desert, California, U.S.A.



Biography

Dr. MARY SEELY received her Ph.D. at the University of California, Davis, and a D.Sc. Honoris causa from the University of Natal, Pietermaritzburg. She is honorary professor at the University of Namibia and the University of Witwatersrand, and adjunct professor at the University of New Mexico and U.C.Davis. In 1970, she became director of the Desert Ecological Research Unit and in 1990 Executive Director of the Desert Research Foundation of Namibia. Apart from publishing 160 papers and 9 books, Dr. Seely has played a leading role in connecting science to development in Namibia.



Dr. JOH HENSCHER received a M.Sc. in Marine Biology at the University of Cape Town and a Ph.D. in Mammal Ecology at the University of Pretoria. After completing a research fellowship of the Alexander von Humboldt Foundation in Germany, he became coordinator of research and information services at the DRFN. His association with this institution dates back to 1977. Dr. Henschel has published 60 scientific papers and 35 popular articles. His current research fields include population dynamics, food webs, climatology, natural resource management, and long-term ecological research, particularly in the Namib Desert.



Ms. JULIANE ZEIDLER submitted her Ph.D. thesis on Indicators of Biological Integrity during 1999 at the University of Witwatersrand. After traveling for one year by car from Germany through Africa, she became associated with the DRFN in 1991, and has since completed two M.Sc. degrees, one on Arthropod Ecology, and the second in Conservation Biology. Ms. Zeidler is currently the DRFN project manager of the Namibian Programme to Combat Desertification in which she is linking ecological research with community based research in the Namib and other areas of Namibia.

JOURNAL 48
Namibia Wissenschaftliche Gesellschaft / Namibia Scientific Society
Windhoek, Namibia 2000
ISSN: 1018 - 7677

THE SPECIES DIVERSITY, DISTRIBUTION AND CONSERVATION OF NAMIBIAN REPTILES: A REVIEW

MICHAEL GRIFFIN

Ministry of Environment and Tourism, Biodiversity Inventory, Private Bag 13306, Windhoek,
Namibia

e-mail: ssarurus@iafrica.com.na

ABSTRACT

A brief historical chronology of Namibian reptile research is given. Namibia's extant fauna of ~ 258 species represents approximately 53% of the Southern African Subregion's species richness, 79% of generic richness, and 100% of the region's familial richness. Sixty-six species are presently recognized as endemic (75% or more of total population range). These endemics occur predominantly in the Namib desert, pro-Namib and adjoining inland escarpment, and are primarily rupicolous. The Namibian endemic reptile fauna is characterised by the speciose genera *Pachydactylus*, *Rhoptropus* and *Ptenopus* (Gekkonidae), *Meroles* and *Pedioplanis* (Lacertidae) and the Scincidae as well as the Cordylidae have also speciated extensively.

Approximately 67% of all Namibian species are provisionally regarded to be "Secure". However, due to a paucity of data, 60 species (25%) are assigned to possible or probable threat categories. Seven species are considered to be "Vulnerable".

Approximately 13% of the Namibian landscape is set aside for preservation purposes. Approximately 90% of the reptile fauna is represented within the formal protection network, ~65% of all species occur in three or more reserves and ~23% occur in 10 or more reserves.

Over-collecting (and gathering) and the alteration of habitat (wetlands as well as rupicolous substrates) are identified as conservation issues.

INTRODUCTION

History remembers Johan Wahlberg as an early Namibian explorer, who, nearly two years after landing in Walvis Bay in 1854, was killed by an elephant bull near Lake Ngami. Although not the first naturalist/explorer to travel through Namibia (he was preceded by, amongst others, Le Vaillant, Alexander, and Andersson), nor among the first to have an unfortunate encounter with local wildlife; George Bonfield, for instance, was knocked into the Kavango River by a hippo, and then immediately