

TRANSBOUNDARY SPECIES PROJECT

ROAN, SABLE AND TSESSEBE

ROWAN B. MARTIN

SPECIES REPORT FOR ROAN, SABLE AND TSESSEBE
IN SUPPORT OF
THE TRANSBOUNDARY MAMMAL PROJECT
OF THE
MINISTRY OF ENVIRONMENT AND TOURISM, NAMIBIA
FACILITATED BY
THE NAMIBIA NATURE FOUNDATION
AND
WORLD WILDLIFE FUND
LIVING IN A FINITE ENVIRONMENT (LIFE) PROGRAMME



COVER PICTURE ADAPTED FROM THE ILLUSTRATIONS BY CLARE ABBOTT IN
THE MAMMALS OF THE SOUTHERN AFRICAN SUBREGION

BY REAY H.N. SMITHERS

PUBLISHED BY THE UNIVERSITY OF PRETORIA

REPUBLIC OF SOUTH AFRICA

1983

CONTENTS

1. BIOLOGICAL INFORMATION	1
a. Taxonomy	1
b. Physical description	3
c. Habitat	6
d. Reproduction and Population Dynamics	12
e. Distribution	14
f. Numbers	24
g. Behaviour	38
h. Limiting Factors	40
2. SIGNIFICANCE OF THE THREE SPECIES	43
a. Conservation Significance	43
b. Economic Significance	44
3. STAKEHOLDING	48
a. Stakeholders	48
b. Stakeholder Institutions – Present and Future	51
c. Towards Trans-Boundary Institutions	53
4. MANAGEMENT	56
a. Present and Future Management of Roan, Sable and Tsessebe	56
b. Transboundary Issues	64

List of Appendices

1. A Generic Population Model for Roan, Sable and Tsessebe	66
2. Air Survey Estimates for Roan, Sable and Tsessebe in Namibia	70
3. Rainfall Data for Selected Sites	74
4. Financial Analysis of Sport Hunting Potential for Roan, Sable and Tsessebe	79
5. Protected Area Requirements in Southern Africa	88

BIBLIOGRAPHY	92
---------------------	----

List of Figures

1. Taxonomy of the Antelope Family (<i>Bovidae</i>) based on Smithers (1983)	2
2. The Distribution of Roan Antelope in Africa	3
3. The Distribution of Sable Antelope in Africa	4
4. The Distribution of Tsessebe Antelope in Africa	5
5. Habitat Niches for Roan, Sable and Tsessebe	8
6. Distribution of Roan Antelope in Southern Africa in Relation to Annual Rainfall	17
7. Distribution of Sable Antelope in Southern Africa in Relation to Annual Rainfall	18
8. Distribution of Tsessebe in Southern Africa in Relation to Annual Rainfall	19
9. Distribution of Roan Antelope in Namibia	20
10. Distribution of Sable Antelope in Namibia	21
11. Distribution of Tsessebe in Namibia	22
12. The Potential Range for Roan, Sable and Tsessebe in Namibia	23
13. Estimates of the Roan Population in Etosha National Park	28
14. Estimates of the Roan Population in Waterberg Plateau Park	29
15. Estimates of the Roan Population in Bushmanland including Khaudum	30
16. Estimates of Roan Populations in East and West Caprivi	31
17. Estimates of Sable Populations in Etosha and Waterberg	32
18. Estimates of Sable Populations in East and West Caprivi	33
19. Estimates of Tsessebe Populations in Etosha, Khaudum and Waterberg	34
20. Estimates of Tsessebe Populations in East and West Caprivi	35
21. Land Tenure in Namibia above the 400mm Rainfall Isohyet	49
22. Institution for Botswana-Namibia management of shared wildlife species populations	55
23. Effect of hunting quotas on age structure of adult males in the population	60

List of Tables

1. Habitat niches occupied by roan, sable and tsessebe	7
2. Habitat requirements of roan, sable and tsessebe	9
3. Grass species eaten by roan, sable and tsessebe	10
4. Woody plant species in the diet of roan and sable	11
5. Reproductive parameters for roan, sable and tsessebe	12
6. Sensitivity of the population to changes in adult female and juvenile survival	13
7. The potential impact of roan, sable and tsessebe on sport hunting income	44
8. Areas above the 400mm rainfall isohyet with roan, sable or tsessebe populations	50
9. Required Budgets for State Protected Conservation Areas	57
10. Effect of hunting quotas on age structure of adult males in the population	60

PREFACE

I have not given a list of acronyms at the start of this report because I have tried to avoid using them in the text and, where one is used, the meaning is given together with the acronym. This draft has not benefitted by having another person review it and is therefore likely to contain numerous typing errors, omissions and spelling mistakes. I seem to be deficient in noticing my own errors. Hopefully, any such mistakes can be corrected in a second draft.

I had hoped that this report would be shorter than the background study on buffalo which I completed in November last year, but it is much the same length. It is long partly because a large amount of background information is available on roan, sable and tsessebe and much of it is relevant to the subject in hand. There are more diagrams and less text in this report and a larger amount of detail has been consigned to the appendices. It is long also because some extremely interesting concepts and hypotheses have arisen in the course of the work and these have needed to be examined thoroughly.

I would like to thank all those people who gave so kindly of their time and valuable experience to this project. In particular, I thank Chris Brown of the Namibia Nature Foundation, who provided me with a large number of references in the literature on roan, sable and tsessebe, John Mendelsohn who assembled rainfall data for on a number of sites in northern Namibia and Peter Erb who gave me all of his working notes and recent data on the three species. I thank also Chris Weaver of the WWF LIFE programme, and Ben Beytell and Pauline Lindeque of the Ministry of Environment and Tourism who spared considerable time to discuss the subject matter. I am unused to receiving such a high level of assistance on consulting work and thank Barbara Paterson as the project co-ordinator for her support. I thank Kevin Dunham, Fay Robertson and David Cumming all of whom assisted me with key documents. Finally, I would like to thank Debbie Gibson and Colin Craig for accommodating me in Windhoek, making a large amount of aerial survey data available and spending much time in discussion of the study.

1. BIOLOGICAL INFORMATION

a. Taxonomy

The current state of the taxonomy of the entire order Artiodactyla (even-toed ungulates) is in some disarray. New suborders and superfamilies have been created, and there seems to be some uncertainty whether Tsessebe still belong in the genus *Damaliscus*. For the purposes of this study, the classification of the three species used by the Antelope Specialist Group of the IUCN Species Survival Commission (ASG 1998) has been adopted –

Roan antelope – *Hippotragus equinus* (Desmarest 1804)

Sable antelope – *Hippotragus niger* (Harris 1838), subspecies *H.n. niger*

Tsessebe – *Damaliscus lunatus* (Burchell 1824), subspecies *D.l. lunatus*

The taxonomic relationship of the three species within the antelope family **Bovidae** as given by Smithers (1983) is shown in **Fig.1** on the following page. The species shown in red font are those for which Background Studies have been carried out under the Transboundary Species Project of the Ministry of Environment and Tourism.

Smithers (1983) notes that Ansell (1972) lists six subspecies of roan for the African continent, including a subspecies *H. e. cottoni* occurring in northern Botswana, the Caprivi and extending into Angola and southern Zaire. However, in the latest classification of the Antelope Specialist Group there is no mention of this subspecies and it seems there is no longer a valid basis for recognising any subspecies of roan antelope.

Apart from the Giant sable in Angola (*H. niger variiani*), it seems that the taxonomic status of the other subspecies of sable (*H. n. roosevelti* which occurs in coastal Kenya and northern Tanzania, and *H. n. kirkii* found in the Selous Game Reserve) are under question at the moment.

Ansell (1972) lists 7 subspecies of *Damaliscus lunatus* but the Antelope Specialist Group has reduced this number to 5 subspecies: *D. l. lunatus* – Tsessebe in southern Africa; *D. l. korrigum* – Korrigum, in northern Benin, Nigeria and Cameroun; *D. l. tiang* – Tiang, occurring eastwards from Chad across southern Sudan to Ethiopia and northern Kenya; *D. l. topi* – Coastal Topi from East Africa and *D. l. jimela* – the Topi in western Tanzania extending into south-western Kenya and Uganda.

The maps showing the continental distribution of the three species (**Figs. 2, 3 & 4**) are shown on the three pages following the taxonomic diagram.




Figure 1: Taxonomy of the Antelope Family (*Bovidae*) based on Smithers (1983)




Figure 2: The Distribution of Roan Antelope in Africa. *Source: ASG (1998)*

b. Physical description

Roan are a large antelope, second only in size to eland. Adult males have a shoulder height of about 1.4 metres and weigh slightly under 300kg. Females are slightly smaller and lighter. Both sexes carry horns which rise from the head in a uniform backward curve and are ridged. The longest pair of horns on record (99 cm) are from a trophy taken in the Tokwe Valley, Zimbabwe (Best and Best, 1977). The females' horns are smaller than the males.

The body coat is greyish brown with touches of rufous colouring which varies amongst individuals. The legs are darker brown than the remainder of the body and the animal possesses a noticeable mane. The most distinctive features are the facial markings which resemble a black balaclava mask with elongated white 'cut-outs' around the eyes. The ears are unusually long with dark brown tufts on the tips.




Figure 3. The Distribution of Sable Antelope in Africa. *Source: ASG (1998)*

Sable are slightly smaller than roan, with adult males standing under 1.4 metres at the shoulder and weighing about 230kg. There is a distinct dimorphism: older males have shiny black coats whilst females are dark brown. Subadults and juveniles of both sexes tend to be a lighter shade of brown. All have white bellies with the colour extending inside the rear legs and up to the base of the tail. The face is marked longitudinally with white stripes extending from each eye towards the muzzle.

Both sexes carry horns but the fully developed male horns are far larger than those of the females. The horns sweep back from the head in a characteristic curve, with the largest trophies from the southern African region being just over 50 inches.




Figure 4: Distribution of tsessebe and other subspecies of the species *Damaliscus lunatus* in Africa.

Source: ASG(1998a)

Tsessebe are smaller than sable, with adult males standing about 1.2 metres at the shoulder and weighing about 140kg. The average female body weight is slightly less than 130kg. The general colour of the body is dark reddish-brown with a purplish sheen. The upper part of the head and muzzle is almost black. The colouration of the flanks and withers are darker than the remainder of the body giving the appearance of a ‘watermark’ from mid-body downwards. There is some yellowish colouring on the base of the tail, the back of the ears, the insides of the hind legs and on the abdomen. Tsessebe have a characteristic body shape which slopes downwards towards to the rump.

Both sexes possess horns which rise close together in the centre of the head and bend outwards and upwards in a uniform curve. A record trophy of 47cm (Best and Best, 1977) was taken in northern Botswana.

c. Habitat

Pienaar (1974) examined the range of physiognomic habitat types occupied by the different large mammal species in Africa. Coetsee (1980) re-ordered Pienaar's original table of vegetation types and clustered the various mammal species to show similarities of habitat occupation. At the broadest level, Coetsee categorised habitats into three main types: dense woody vegetation, non-desert open vegetation, and subdesert and desert vegetation. Roan, sable and tsessebe are found only in the second category. Coetsee noted specifically the close similarity between roan and tsessebe in their range of selected habitats.

I have revisited Pienaar's original classification and present the tables on the following page (**Table 1**) which show the 'niches' occupied by roan, sable and tsessebe. The niche for each species is essentially three-dimensional: the first axis (the columns in the matrix below) shows the occurrence of the species along a moisture gradient, the second axis (the rows in the matrix below) shows the degree of openness of the habitat, ranging from deserts to closed forests, and the third axis is an altitudinal gradient from lowland tropical to temperate montane conditions (the dual columns under the broad heading "Mesic" in each table). The content of the tables is represented pictorially in **Fig. 5** on the next page.

All three species occur only in the middle range of moisture conditions (mesic) – they are not found in very dry or in high rainfall areas. This is significant for attempts to conserve them in Namibia – the majority of the country falls below the lower limit of acceptable precipitation. Both roan and tsessebe are found across a wide altitudinal range – they occur from tropical lowlands to temperate montane conditions (e.g. the Nyika plateau in Malawi). Sable, however, are not found in the in the cool high altitude areas. All three species have a strong preference for open and fairly well treed savannas, although roan and tsessebe tend to avoid closed canopy woodlands. Roan is the only one of three species found in rocky outcrops and sable does not occur in swamp/grassland habitats.

I have summarised and compared the key findings on the habitat requirements of roan, sable and tsessebe in **Table 2** (page 9) which has been compiled from studies carried out by Dörgeloh (1998), Dunham and Robertson (2001), Erb (1993), Grobler (1974), Harrington (*et al* 1999), Joubert (1976a), Joubert and Bronkhorst (1977), Pienaar (1963), Smithers (1983), Wilson (1975) and Wilson and Hirst (1977).

A list of grass species eaten by roan, sable and tsessebe (**Table 3**, page 10) has been compiled from detailed feeding studies undertaken by Erb (1993), Grobler (1974, 1981b), Huntly (1972), Joubert (1976), Perrin and Taolo (1999), Wilson (1975) and Wilson and Hirst (1977). Because most of the studies have been done in restricted localities, the list is unlikely to be comprehensive. It nevertheless shows that there are a number of grass species common to the diet of all three antelope species. At the genus level, which is probably a more useful indicator of shared preferences on a regional scale, seven grass genres are common to all three antelope species.

Both roan and sable browse to a limited extent in the dry season – sable less so than roan. Wilson (1975) noted that roan in northern Transvaal remained in better body condition than sable because of being able to switch to browsing in critical periods. A list of woody species eaten by roan and sable (**Table 4**, page 11) has been compiled from the studies of Erb (1993), Grobler (1974), Joubert (1976), and Wilson (1969).

Table 1. Habitat Niches occupied by Roan, Sable and Tsessebe

ROAN	DRY	MESIC		WET
		TEMPERATE	TROPICAL	
ROCKY OUTCROP			1	
GRASSLAND/SWAMPLAND		2	2	
SHRUB SAVANNA		1	2	
OPEN TREE SAVANNA		2	2	
TREE SAVANNA			2	
WOODLAND			1	
THICKET				
FOREST				

SABLE	DRY	MESIC		WET
		TEMPERATE	TROPICAL	
ROCKY OUTCROP				
GRASSLAND/SWAMPLAND				
SHRUB SAVANNA			1	
OPEN TREE SAVANNA			2	
TREE SAVANNA			2	
WOODLAND			2	
THICKET				
FOREST				

TSESSEBE	DRY	MESIC		WET
		TEMPERATE	TROPICAL	
ROCKY OUTCROP				
GRASSLAND/SWAMPLAND		2	2	
SHRUB SAVANNA		2	2	
OPEN TREE SAVANNA		1	2	
TREE SAVANNA			2	
WOODLAND			1	
THICKET				
FOREST				

In the cells of each matrix, the number '1' indicates that the species has been recorded in the particular habitat and the number '2' indicates a strong preference for the habitat type.




Figure 5. Habitat Niches of Roan, Sable and Tsessebe

Table 2. Habitat requirements of Roan, Sable and Tsessebe

Roan	Sable	Tsessebe
All are predominantly grazers		
Include a small amount of browse in their dry season diet		Not known to browse
All are water dependent		
Usually found within 5km of water.	Usually found within 3km of water.	Usually found within 1km of water.
All occur primarily in savanna woodlands		
Prefer lightly wooded savanna with open areas of medium to tall grasses. Tolerant of low bush growth up to 1.5m provided it is open and patchy. Well adapted to grass heights up to 1.5m.	Dependent on cover. Prefer open woodlands adjacent to vleis or grassland with medium to high stands of grass.	Optimum habitats are open stands of healthy grasslands with easily accessible shelter in form of trees or shrubs either as an ecotone or as scattered islands in the grassland.
Avoid closed canopy woodland and thick closed stands of bush 1.5-4m high. Avoid short grass areas.	Avoid dense woodland and short grasslands especially when overused by other species.	Avoid habitats where plant height exceeds 2 metres.
Spend little time on burns even when a green flush occurs – prefer taller grass stands.	Not particularly attracted to burns.	Prefer burnt to unburnt areas. Readily concentrate on burned areas when a green flush appears.
All are highly selective feeders		
– characterised by narrow muzzles with which they can select particular clusters of leaves from grass swards		
Select climax green grass species with a high nutrient content and those that have a high leaf to stem ratio. Delicate feeders, using the higher parts of grasses from 8 cm above ground – not close croppers like wildebeest or zebra. In any given locality, two or three grass species make up the bulk of diet.	Prefer medium height green grasses and have a narrow range of acceptable grass species.	Strong predilection for young green grass shoots up to 60cm tall. Appear to select for stage of grass growth rather than species <i>per se</i> .
All are sensitive to habitat changes and have critical habitat requirements		
Physiognomic changes to vegetation structure such as those brought about by elephants are capable of a major impact on all three species. Loss of canopy trees resulting in changes to the species composition and structure of grass swards and the trampling effects of large numbers of elephant are all potentially negative influences. Overutilisation, inter-specific competition and trampling of grass by cattle and other large mammal species also renders habitats less favourable for roan, sable and tsessebe.		
Sensitive to any increase in the density of woody plants or reduction in grass cover.	Unable to cope with superabundant grass in good rainfall years. Grass swards are underutilised, resulting in favourable conditions for tick irruptions.	Structural changes to habitats which obstruct movement, affect access to water, visibility and cover all have a major impact on tsessebe.
As a result of these factors, the distribution of all three species is patchy and discontinuous across their range.		

Table 3. Grass species eaten by roan, sable and tsessebe

Roan	Sable	Tsessebe
<i>Andropogon schirensis</i>		
<i>Aristida spp</i>		
	<i>Brachiaria brizantha</i>	
<i>Brachiaria nigropedata</i>	<i>Brachiaria nigropedata</i>	<i>Bracharia nigropedata</i>
	<i>Cynodon dactylon</i>	<i>Cynodon dactylon</i>
<i>Digitaria pentzii</i>	<i>Digitaria pentzii</i>	<i>Digitaria pentzii</i>
<i>Digitaria serriata</i>		
<i>Digitaria spp</i>		
<i>Eragrostis curvula</i>	<i>Eragrostis gummiflua</i>	<i>Eragrostis chloromelas</i>
<i>Eragrostis plana</i>		
<i>Eragrostic superba</i>		
<i>Eragrostis jeffreysii</i>		<i>Eragrostis jeffreysii</i>
<i>Heteropogon contortus</i>	<i>Heteropogon contortus</i>	<i>Heteropogon contortus</i>
<i>Hyparrhenia hirta</i>	<i>Hyparrhenia hirta</i>	<i>Hyparrhenia hirta</i>
<i>Hyparrhenia spp</i>		
	<i>Hyperthelia dissoluta</i>	<i>Hyperthelia dissoluta</i>
<i>Melinis spp</i>		
<i>Panicum colorata</i>		
<i>Panicum kalaharensis</i>		
<i>Panicum maximum</i>	<i>Panicum maximum</i>	
	<i>Pogonarthria squarrosa</i>	
<i>Rhynchelytrum repens</i>		<i>Rhynchelytrum repens</i>
	<i>Schyzachyrium jeffreysii</i>	
<i>Schizachyrium sanguineum</i>		
<i>Schmidtia pappophoroides</i>		
	<i>Setaria perennis</i>	<i>Setaria perennis</i>
<i>Setaria woodii</i>		
<i>Setaria spp</i>		
		<i>Sorghum verticilaster</i>
<i>Sporobolus fimbriatus</i>		
<i>Themeda triandra</i>	<i>Themeda triandra</i>	<i>Themeda triandra</i>
	<i>Trachypogon spicatus</i>	
<i>Trichoneura grandiglumis</i>		
<i>Triraphis schinzii</i>		
	<i>Urochloa bolbodes</i>	
Grass species which are common to roan, sable and tsessebe		
Grass species which are common to two of the three antelope species		

An examination of the table shows the following grass genera are common to the diet of roan (R), sable (S) and tsessebe (T)

Genus common to RST	<i>Brachiaria, Digitaria, Eragrostis, Heteropogon, Hyparrhenia, Setaria, Themeda</i>
Genus common to RS only	<i>Panicum, Schizachyrium</i>
Genus common to RT only	<i>Rhynchelytrum</i>
Genus common to ST only	<i>Cynodon, Hyperthelia</i>

Table 4. Woody plant species in the diet of roan and sable

Species	Locality
<i>Acacia ataxacantha</i>	Waterberg, Namibia
<i>Acacia karroo</i>	Matopos NP, Zimbabwe
<i>Acacia</i> spp.	Percy Fyfe Nature Reserve, South Africa
<i>Dichrostachys cinerea</i>	Matopos NP, Zimbabwe
<i>Dombeya rotundifolia</i>	Matopos NP, Zimbabwe
<i>Faurea saligna</i>	Kruger NP, South Africa
<i>Grewia flava</i>	Matopos NP, Zimbabwe
<i>Grewia monticola</i>	Matopos NP, Zimbabwe
<i>Grewia reticulata</i>	Waterberg, Namibia
<i>Lippia javanica</i>	Matopos NP, Zimbabwe
<i>Lippia oatzii</i>	Matopos NP, Zimbabwe
<i>Lonchocarpus nelsii</i>	Waterberg, Namibia
<i>Ochna pulchra</i>	Waterberg, Namibia
<i>Olea africana</i>	Kruger NP, South Africa
<i>Philenoptera violacea</i> *	Percy Fyfe Nature Reserve, South Africa
<i>Tarchonanthus camphoratus</i>	Matopos NP, Zimbabwe
<i>Rhus lancea</i>	Kruger NP, South Africa & Matopos NP, Zimbabwe
<i>Rhus pyroides</i>	Kruger NP, South Africa
<i>Ziziphus mucronata</i>	Matopos NP, Zimbabwe

* Previously *Lonchocarpus capassa*

d. Reproduction and Population Dynamics

There is sufficient similarity in the biological parameters which determine the population dynamics of roan, sable and tsessebe to permit a generic population model to be used to examine their expected breeding performance under average conditions. The key parameters have been derived from Erb (1993), Dunham, Robertson & Swanepoel (2003), Dunham and Robertson (2001), Grobler (1974, 1979, 1980, 1981a), Joubert and Bronkhorst (1977), Penzhorn and van der Merwe (1993) and Smithers (1983) and are summarised in **Table 5** below.

Table 5. Reproductive parameters for roan, sable and tsessebe

Seasonal breeding	Roan breed throughout the year; sable give birth to calves January-March; tsessebe give birth to calves October-November.
Gestation	All 8-9 months
Age at first conception	Almost all females conceive after about two years
Age at first parturition	Almost all females produce calves in third year of life
Fecundity (adults)	All adult females are capable of producing a calf every year – fecundity of roan may be slightly above unity in favourable conditions
Longevity	Few animals survive beyond 12 years of age in the wild
Breeding longevity	Females are probably capable of breeding throughout their adult life although fecundity may be reduced in last few years
Mortality (juveniles)	About 25% in average years
Mortality (yearlings)	Females about 5%, males generally higher
Mortality (adult males)	Around 10%, increasing in the last few years of life
Mortality (adult females)	Less than 5% except in last few years of life
Adult sex ratio ♀♀:♂♂	About 2:1 depending on hunting regimes and predation

A simple population model using these parameters is given in **Appendix 1**. Under average conditions, roan, sable and tsessebe populations could be expected to increase at a rate of slightly under 14% per annum when they are below ecological carrying capacity.¹

The term “average conditions” is loaded. In semi-arid ecosystems, variations in annual rainfall may have profound effects on the performance of these three species. Dunham and Robertson (2001) and Dunham, Robertson and Swanepoel (2003) demonstrated a strong relationship between the survival of tsessebe and the accumulated surplus or deficit in rainfall over a long period. They also found a strong correlation between adult and juvenile mortality and the rainfall in the late dry season. Rainfall in late dry season appears to be critical, affecting the animals’ condition, survival rate, late stage of pregnancy and early stage of lactation.

1. Sable and tsessebe are recorded as reaching densities of over 4/km² in favourable habitats where rainfall is 500-600mm per year. At this level of rainfall, the ceiling densities for roan seldom exceed 2/km² (Erb 1993).

The population models developed by Dunham *et al* based on these variables (which also take into account tsessebe numbers, i.e. density dependence) appear to explain very adequately the variations in tsessebe populations in both Kruger National Park and on a mixed cattle/game ranch in Zimbabwe. Erb (1993) modelled the roan population in the Waterberg Plateau Park in Namibia incorporating the late dry season rainfall and obtained a very close fit between the actual population estimates and the number predicted from the model. In a study of sable in Zimbabwe, Grobler (1981) found that any animals weakened by stress during the dry season were likely to die from parasites soon after onset of rains. This phenomenon appeared particularly to affect neonatal mortality and subsequent juvenile mortality when losses increased towards end of the dry season caused by a downward trend in nutrition.

It seems eminently feasible that the long term surplus or deficit in rainfall² could be the main determinant of the vegetation structure, particularly of grasslands, in any given locality. A prolonged drying out process would affect water tables and the catenas in vegetation from river banks to the upper reaches of catchments. Species such as tsessebe would find their preferred habitats shrinking to narrow bands close to rivers. Sable and roan would be affected by changes in species composition in grass swards and shifts between perennial and annual grasses.

I have examined the extent to which the given regime of adult female and juvenile mortality in **Appendix 1** (which was derived for ‘average conditions’) would have to change in order to throw the ‘generic’ population into a decline (**Table 6**). In the two tables, each age-specific female survival rate has been changed by the proportion indicated.

Table 6. Sensitivity of the population to changes in adult female and juvenile survival

Table 6a. Effects of changes in adult female survival on population growth rate

Change in adult female survival %	+2%	+1%	0	-5%	-10%	-15%	-20%
Rate of population growth %	15.5	14.5	13.6	8.9	4.2	0.0	Decline

Table 6b: Effects of changes in juvenile survival on population growth rate

Change in juvenile survival %	+30%	+20%	+10%	0	-10%	-20%	-30%	-40%	-50%	-55%	-60%
Rate of population growth %	19.7	17.4	15.6	13.6	11.5	9.3	6.8	4.1	1.0	0.0	-ve

In the population model of Appendix 1, juvenile survival was set at 75% (a mortality of 25%). It is apparent from Table 6b that the population can tolerate very large increases in juvenile mortality – if survival is halved, the population continues to increase – albeit at a low growth rate (1%). The same is not true for adult female survival (for the purposes of this test, all female age classes other than juveniles were treated as ‘adults’). An overall decrease of 15% in the existing schedule of age-specific survival rates of adult females brings population growth to zero. The linkage between rainfall and adult mortality would not have to be particularly pronounced to produce major swings in the population.

2. In mathematical terms, this is effectively the integral of the deviations above and below the mean rainfall. It is the area under the curve of surpluses and deficits.

e. Distribution

(1) Regional Distribution

The distribution of roan, sable and tsessebe in southern Africa in relation to average annual rainfall is shown in **Figures 6-8** (pages 17,18,19). The data are from ASG(1998a) and are somewhat crude at the scale of individual subpopulations. Nevertheless, it is notable that –

- **There are no ‘naturally’ occurring populations of roan, sable or tsessebe below the 400mm rainfall isohyet;**
- Apart from a very few minor introduced populations of sable and tsessebe elsewhere in the region,³ **Namibia is the only southern African country attempting to maintain populations of roan, sable and tsessebe in areas where the average rainfall is below 400mm.**⁴
- **Many of the observed population ‘crashes’ in roan, sable and tsessebe populations in the region have occurred in areas where the average annual rainfall is close to the lower limit which the species can tolerate.**

Mills, Biggs and Whyte (1995) concluded that rainfall was the principal determinant of numbers of common ungulates in southern African savannas. This fundamental constraint appears to receive too little emphasis in the plethora of proposed research studies and management measures designed to address the declines in roan, sable and tsessebe in parts of southern Africa (e.g. Grant and van der Walt 2000, Grant *et al* 2002, Harrington 1995, Harrington *et al* 1999, RARE 2002 and numerous other references). Dunham and Robertson (2001) demonstrate very clearly that rainfall accounts for the observed declines in tsessebe populations in Kruger National Park yet there appears to be a reluctance to accept this too-simple finding or apply it to other ‘rare species’ populations.

(2) Distribution of Roan, Sable and Tsessebe in Namibia

In the next three figures, the distribution of each species is shown in relation to rainfall.

Roan

The present and historical distribution of roan antelope in Namibia is shown in **Fig. 9** (page 20). The original range for the species has been taken from Shortridge (1934) and falls entirely above the 400mm rainfall isohyet. Hahn (1925) records roan as far east as 14°E along the Cunene River in Ovamboland although noting that these were temporary movements during the rains. Gaerdes (1963) has examined early records of roan and gives none outside the range mapped by Shortridge. Joubert and Mostert (1975) note that the range for roan had decreased considerably since Shortridge’s work.

-
3. Below the 400mm rainfall isohyet, introduced populations of sable and tsessebe occur on private land in the vicinity of the Limpopo River in the northern province of South Africa and in the Beit Bridge area of Zimbabwe, and there is an introduced population of tsessebe in the Vaalbos National Park near Kimberley in South Africa.
 4. The African Antelope Specialist Group (ASG 1998) remark that the populations of roan, sable and tsessebe in Namibia are outside of the species’ ‘natural range’.

The present ‘natural range’ for roan includes a variable number of animals in the East and West Caprivi, Khaudum Game Reserve and the Nyae Nyae Conservancy. These are not separate subpopulations: rather they are the extreme animals on the outskirts of the larger roan population in northern Botswana.

Roan were introduced to Etosha National Park in 1970 and to the Waterberg Plateau Park in 1975. The Kaross area in the west of Etosha where the main roan population is located falls below the 300mm rainfall isohyet and the population has not thrived (see next section on numbers). The Waterberg population is above the 400mm rainfall isohyet and has done better.

The various commercial farms onto which roan have been introduced have been accurately plotted in **Fig. 8**. Most of the farms lie below the 400mm rainfall isohyet⁵ and it is likely that the long-term survival of roan on these properties will be dependent on supplementary feeding.

Sable

The present and historical distribution of sable antelope in Namibia is shown in **Fig. 10** (page 21). Shortridge (1934) did not record the species outside of the Caprivi and it is not mentioned by Hahn (1925) as occurring in Ovamboland. Joubert and Mostert (1975, page 21) refer to sable in their list of “unrecorded mammals” and regard the odd sightings of the species in Caprivi as temporary excursions of Botswana animals. However, surveys of the Caprivi since 1980 suggest that there is a permanent population in several parts of the Caprivi.⁶

Sable were introduced to Etosha in 1978 and today there is a small population in the Khaobendes paddock in the extreme west of Etosha National Park which has remained fairly constant at around 50 animals for some years and there are about 30 sable in the park proper (Killian, pers. comm.). Except for the eastern end of the Park, the entire area falls below the 400mm rainfall contour and must be regarded as marginal range for sable. Sable were introduced to the Waterberg Plateau Park in 1980 and they have increased to around 100 animals over the past twenty years.

There are a large number of sable on commercial farms (two-thirds of the national population) but the majority of the properties shown in **Fig. 10** lie below the 400mm rainfall isohyet in marginal habitats for the species. As with roan, it is likely that they will require supplementary feeding in below average rainfall years to maintain their numbers.

Tsessebe

The present and historical distribution of tsessebe in Namibia is shown in **Fig. 11** (page 22). The early range for tsessebe has been taken from Shortridge (1934) and it is notable that the species did not occur below the 500mm rainfall isohyet. Shortridge notes that the species was occasionally seen in what is now the extreme northerly end of Khaudum Game Reserve and, as recently as the mid-1980s, they were recorded during aerial surveys of this area.

5. One farm, the Ohorongo Game Reserve, lies below the 300mm rainfall isohyet and is not far from the Etosha Kaross population.

6. Mahango National Park, the western Core Area of the Caprivi Strip and the Kwando Triangle.

In the Caprivi, tsessebe were relatively abundant in the early part of the twentieth century. Gaerdes (1969) does not mention the species outside of the Caprivi and it is not mentioned by Hahn (1925) as occurring in Ovamboland.

Tsessebe were recorded in significant numbers in the Caprivi⁷ in 1994 in two independent and comprehensive air surveys (ULG 1994 and Rodwell *et al* 1995). Since then there have not been any surveys covering all of the Caprivi in a single year – however, the limited surveys which have been done (Craig 1998 and Craig 2000) suggest that the tsessebe populations in the Caprivi have almost disappeared. Craig (1998) saw tsessebe in small numbers (fewer than 30 animals) in both Mahango and in the Kwando Triangle but, in the last air survey of Mahango (Craig 2000), no tsessebe were seen.

Tsessebe were unsuccessfully introduced to the Waterberg Plateau Park in 1981 with all of the founder population dying before being released from the holding paddock. In May 1984 a male was transferred from Khoabendes to Waterberg and in May 1985 14 adults were received from Percy Fyfe Nature Reserve, Transvaal (Erb 1992). The population has not thrived and only two animals are thought to be surviving now.

There are seven populations of tsessebe on commercial farms (**Fig.11**), four of which number more than 20 animals. Apart from one small group (12 animals) on Kamapu-Oos,⁸ all of these newly established populations lie below the 400mm rainfall isohyet in far from optimum locations.

Summarising this section, the most suitable habitats for roan, sable and tsessebe lie in the extreme north-west of Namibia where mean annual rainfall is above 500mm – primarily in the Caprivi and Kavango regions (**Fig.12**). Over the long term, the three species should survive above the 400mm isohyet but they will always be vulnerable to droughts in this zone.

Not unexpectedly, the highest human population densities also occur in the most favourable habitats for the three species. However, the parts of the former ‘natural range’ for roan, sable and tsessebe in which human densities exceed 10 persons/km² are less than 10% of the total potential range and, in theory at any rate, the potential still exists for the north-west of the country to carry large populations of all three species. However, competition with cattle (whose numbers are not indicated in maps of human population density) will undoubtedly reduce the available range.

Namibia’s ‘wild’ populations of roan, sable and tsessebe all lie on the fringe of the larger Botswana populations. The present international veterinary fences separating the two countries limit the movements of the animals and this could result in isolated subpopulations on either side of the fence.

The combined limiting effects of rainfall, human settlement and veterinary fences are considered again Section 1.g on page 40.

7. Tsessebe were recorded in Mahango, the Kwando Triangle and Mudumu.

8. Farm No. 24, Otjiwarongo District.




Figure 6. Distribution of roan antelope in southern Africa in relation to annual rainfall




Figure 7. Distribution of sable antelope in southern Africa in relation to annual rainfall






Figure 8. Distribution of tsessebe in southern Africa in relation to annual rainfall



SUMMARY TABLE (rounded numbers)

COMMERCIAL FARMS	400
STATE PROTECTED AREAS (excluding Caprivi)	250
CONSERVANCIES (Nyae Nyae)	100
WEST CAPRIVI (including Mahango and Kwando Triangle)	25
EAST CAPRIVI (including Mudumu, Forest Reserve and Conservancies)	25
TOTAL	800


Figure 9. Distribution of Roan Antelope in Namibia



SUMMARY TABLE (rounded numbers)

COMMERCIAL FARMS		800
STATE PROTECTED AREAS	Hoffnung Part II	1
	Western Core (1998)	21
	Kwando Triangle (1998)	70
	Etosha National Park (2003)	78
	Waterberg Plateau Park (2000)	119
	Mahango Game Reserve (2000)	130
	TOTAL	419
	TOTAL	1,200

Figure 10. Distribution of Sable Antelope in Namibia



SUMMARY TABLE (rounded numbers)

COMMERCIAL FARMS	175
STATE PROTECTED AREAS	175
TOTAL	350

Figure 11. Distribution of Tsessebe in Namibia




Figure 12. The Potential Range for Roan, Sable and Tsessebe in Namibia showing Rainfall, Human Population Densities and Veterinary Fences

f. Numbers

Most aerial surveys tend to focus on elephants and buffalo and, as a result, roan, sable and tsessebe are usually under-counted. Sable are more visible from the air than roan and tsessebe so that the estimates for this species may be better than the other two. In deriving estimates for these species a dual problem may occur. If the animals are not present in large groups or large numbers, they tend not to be seen and the resultant estimates suffer from large confidence intervals because too few animals make up the sample for the estimate. If all the animals are being seen but the population is sparse, the estimates will still have large confidence intervals for the same reason. Typical confidence limits for estimates of the species are seldom better than $\pm 50\%$ of the value of the estimate even when the numbers are fairly high, and caution needs to be exercised in pronouncing apparent upward or downward trends in populations.

The data which follow are not very valuable for comparative purposes or for detecting trends because the same areas have not been surveyed consistently from year to year (this effect is particularly prevalent in the Caprivi) and because of the variety of survey techniques which have been used on different surveys.⁹ At present there is no acceptable alternative to the standard transect survey method or the random block count method. Systems which rely on “total counts” or “actual observations” are statistically inferior because no accuracy or precision can be attached to the estimate. In the final part of this report under ‘best practices’ for management, the subject is pursued further.

(1) Namibia

The data for roan, sable and tsessebe from surveys carried out in Namibia since 1970 are presented in **Appendix 2** and shown in **Figs. 12-19** on the pages which follow.¹⁰ Also shown on each of these figures is the cumulative surplus/deficit in rainfall since the 1960s and the late dry season rainfall. The rainfall data are presented in **Appendix 3**.

Dunham and Robertson (2001) found that a relationship of the following form adequately described the rise and fall of tsessebe numbers in Kruger National Park –

$$\text{Adult survival} = f\left((\text{Dry season rainfall}), \int (\text{Annual rainfall}), (\text{Tsessebe density}) \right)$$

Adult tsessebe survival is a function of the amount of dry season rainfall, the integral of the annual rainfall (the accumulated deficit/surplus) and the numbers of tsessebe actually present at the time. The effects of these variables on a tsessebe population would be as follows –

- Rainfall in the late dry season appears to be critical, affecting the animals’ ability to maintain condition and, hence, survival. The effect on females during the late stage of pregnancy and the early stage of lactation is particularly pronounced.

9. The types of surveys which have been carried out include waterhole counts, total counts from fixed wing aircraft, total counts from helicopters and sample surveys based on line transects with calibrated strip widths.

10. Where estimates have been based on standard sample surveys this is indicated for the year concerned.

- The long term surplus and deficit in annual rainfall appears to act on habitats. A prolonged sequence of years where rainfall remains below the average may result in unfavourable physiognomic changes in habitats.
- The last variable, population density, would act negatively on population increase. The higher the number of animals, the greater would be the degree of intra-specific competition for food and the amount of energy expended by adult males on maintaining territories. With the low numbers of tsessebe (and roan and sable) in the various areas where they are present in Namibia, this effect is unlikely to be marked.

Erb (1993) found that the relationship between early season rainfall and calving success was important for roan antelope in the Waterberg. Roan start to calve as early as August when the veld is still dry. Good rains in September/October result in an early green flush which provides lactating cows with the necessary dietary protein to meet a demand which escalates during late pregnancy and early lactation.

I would like to have developed a model incorporating these rainfall parameters and observed the extent to which it explained the performance of roan, sable and tsessebe populations in Namibia. Unfortunately, the population estimates for all areas (with the exception of roan in the Waterberg from 1975-1990) are too erratic to justify any attempts to fit modelled data to observed numbers. The data of **Figs. 12-19** allow at best a broad overview of trends in rainfall and population numbers but, in examining the data, it is necessary to be conscious at all times that many of the observed swings in population numbers may be no more than artifices arising from irregular and incomplete surveys. The following broad observations can be made –

- In all of the main areas where roan, sable and tsessebe population numbers have been estimated since the 1970s,¹¹ it is apparent that the accumulated rainfall surplus reached a peak in the five years between 1975 and 1980. It was also during this time that the populations of all three species appeared to be booming.
- After 1980 the surplus began to decrease although it did not change to a deficit in all areas simultaneously.¹² After 1994 all areas went into a rainfall deficit mode¹³ and this appears, on most of the figures, to coincide with population ‘slumps’ for all species populations.

-
11. Etosha National Park, Waterberg Plateau Park, Bushmanland (including Khaudum and the Nyae Nyae Conservancy) and East and West Caprivi.
 12. Year in which rainfall surplus changed to a deficit: Etosha – 1982; Waterberg – 1991; Bushmanland and West Caprivi – 1994; East Caprivi – 1991. It must be noted that this transition from surplus to deficit is dependent on the time span of data considered (1960-1997): with a longer time span the zero point could shift upwards or downwards.
 13. The most recent rainfall data (1997 onwards) is awaited from the Namibia Weather Bureau and is not shown on the figures.

- The impression given from the figures is that as long as the cumulative rainfall is in a surplus mode populations appear to be maintaining themselves or increasing. It is not necessary to hypothesize that there is any tight, direct relationship between the surplus rainfall and the breeding performance – in fact, other limiting factors may well be operating when the rainfall is in this mode. As soon as the cumulative rainfall changes into a deficit mode, this becomes the factor limiting the population.
- It might be expected that this relationship with the cumulative rainfall surpluses and deficits becomes increasingly critical as the lower rainfall limits of the species range are approached. In the Eastern Caprivi, where annual rainfall is normally well above 500mm, the effects appear less striking. In Kruger National Park, the range preferred by roan and tsessebe is in a marginal rainfall area and the rainfall deficit/surplus relationship is strong. The decline of roan and tsessebe since the 1970s in north-western Zimbabwe may be due to the same phenomenon.
- There is little to be gleaned from the late dry season rainfall data shown in the figures – mainly because the small ups and downs in population numbers from year to year which this factor would influence are obscured by the poor population estimates.

The following observations relate to the individual figures –

- Fig.12: The roan population in Etosha National Park crashed sharply in 1982-83 coinciding exactly with the transition from a cumulative rainfall surplus to a deficit. Since then its numbers have hovered around below 100 animals. This population (mainly in Kaross) is in an area below the 300mm rainfall isohyet and it can be expected that it will always suffer from any small deficit in rainfall.
- Fig.13: The introduction of roan to the Waterberg Plateau Park coincided fortuitously with an optimum part of the rainfall cycle. Since the curve has moved into a deficit mode, the population has decreased to well below 200 animals.
- Fig.14: The roan population in Bushmanland was at its zenith (over 500 animals) in 1985 when the cumulative rainfall surplus was also at a peak. Since this went into deficit mode in 1994, roan estimates have not exceeded 100 animals.
- Fig.15: At first sight, the roan in East and West Caprivi would appear to have reached a peak in 1992-94 (not that 150 animals necessarily represents much of a ‘peak’). In fact the real peak was probably that shown for 1980 when the rainfall surplus was high – thereafter the survey data is erratic and the so-called peak in 1992-94 relates more to a time when a number of comprehensive surveys were carried out. The most recent survey data from 1998 onwards suggests very low roan numbers and the cumulative rainfall deficit may well be the cause of this.

- Fig.16: The sable population in Etosha has hovered around 50 animals for a number years.¹⁴ The Khaobendes paddock in the extreme west of the park cannot be viewed as ideal sable habitat either from a rainfall or vegetation point of view. The introduced sable population in the Waterberg increased to about 150 animals while the cumulative rainfall curve was in a surplus mode but appears to have declined since then.
- Fig.17: The Caprivi sable population appeared to reach a peak of over 1,000 animals in 1994 (over 500 animals in both the West and the East) – however this coincided with two comprehensive surveys done in that year (Rodwell *et al* 1995 and ULG 1994). It is difficult to state whether sable numbers have decreased as drastically as indicated because there have been no comparable surveys since 1994. If they have declined, then it is to be noted that the cumulative rainfall data from both Andara and Katima Mulilo indicate a deficit.
- Fig.18: Tsessebe were introduced to Etosha in 1978 and appear to have gone extinct. The numbers are too low to draw any conclusions about a relationship with the cumulative rainfall graph – it may be pure coincidence that they crashed exactly at the time when this went into a deficit mode. In any case, it is unlikely that the rather special habitat requirements of tsessebe would be satisfied in Etosha.
- The Waterberg tsessebe population appears never to have exceeded 20 animals and Erb (1992) notes that a larger founder population is an urgent necessity. However, it is possible that the required ecotonal habitats for tsessebe do not occur in the park.
- Tsessebe have not been seen in Khaudum since 1987. However, there is always a possibility that re-colonisation will occur from Botswana when the cumulative rainfall regime moves once again into a surplus mode.
- Fig.19: Tsessebe numbers were modest (>100) in East Caprivi in 1980 when the cumulative rainfall surplus was at its peak. As with the sable, the population estimates thereafter are erratic and it is far from certain that the decline which appeared to take place between 1980 and 1994 was real. In 1994 more than 100 tsessebe were estimated in Mudumu and, given that the cumulative rainfall surplus persisted throughout this period, it is most likely that this population had been in place all the time. After 1991 the cumulative rainfall changed to a deficit and the current status of the species must be regarded as uncertain. Being linked to the larger Botswana population, the possibility exists that animals will re-appear when conditions are favourable.

The figures provide plausible evidence for the linkage between roan, sable and tsessebe numbers and cumulative rainfall surpluses and deficits. However, it is not intended to suggest that this is the only mechanism affecting the species populations. Other factors such as human population increases, new areas opened up for settlement, competition with cattle, veterinary fences and illegal hunting are all likely to exert a negative influence. However, the analysis does suggest that when the cumulative rainfall enters a deficit mode, this factor is likely to override all other limiting factors.

14. Werner Killian (pers. comm. Feb, 2003) estimates there are about 50 sable in the Khaobendes paddock and a further 28 at large in the greater park.

ROAN – Etosha National Park




Figure 13. Estimates of the roan population in Etosha National Park




Figure 14. Estimates of the Roan Population in Waterberg Plateau Park




Figure 15. Estimates of the Roan Population in Bushmanland including Khaudum

ROAN – East and West Caprivi




Figure 16. Estimates of Roan Populations in East and West Caprivi





Figure 17. Estimates of Sable Populations in Etosha and Waterberg

SABLE – Caprivi




Figure 18. Estimates of Sable Populations in East and West Caprivi

TSESSEBE – Etosha, Khaudum, Waterberg




Figure 19. Estimates of Tsessebe Populations in Etosha, Khaudum and Waterberg




Figure 20. Estimates of Tsessebe Populations in East and West Caprivi

(2) Neighbouring countries

The countries neighbouring Namibia with populations of roan, sable and tsessebe are Angola, Botswana, Zambia and Zimbabwe. No data is to hand for Zambia and Angola.

The wildlife authorities in Botswana have systematically carried out nation-wide aerial surveys of large mammals in both dry and wet seasons for most years since 1987.¹⁵ ULG (1995) produced pooled estimates of the northern Botswana roan, sable and tsessebe populations and, in the table below, these have been compared with the estimates for the Caprivi in 1994 when two comprehensive surveys were carried out.

	Roan	Sable	Tsessebe	Area
Estimate	1,357	3,138	10,015	145,605 km ²
Confidence intervals	± 49 %	± 35 %	± 22 %	
Density (km ² /animal)	107	46	15	
Caprivi population (1994)	197	1,200	206	20,000 km ²
Percentage of Botswana figures	15 %	38 %	2 %	14 %

The overall densities of the three species are very low in both Botswana and Namibia. In relation to the areas involved, in 1994 the Namibian sable population was at a higher density than the Botswana population, the roan population was more or less at the same density and the tsessebe population was at a much lower density. It is perhaps to be expected that tsessebe numbers in northern Botswana would be high because of the optimum habitats in the vicinity of the Okavango Swamps. Movements of animals between Botswana and the Caprivi are likely to be at their lowest during the dry season and, therefore, the dry season estimates for the Caprivi are probably indicative of the size of the “permanently resident” populations.

The linkages between the roan, sable and tsessebe populations in north-west Zimbabwe and the Caprivi are more tenuous than those of Botswana-Namibia ‘axis’. The survey results are only presented in the context of a long term vision for a trans-frontier conservation area where these species populations are able to move freely between Botswana, Namibia and Zimbabwe. The realities of the present situation are that although veterinary control fences do not theoretically prevent movements of Zimbabwe wildlife westwards into northern Botswana,¹⁶ only minor movements have been recorded.¹⁷ This may be because the physical gap between the international boundary and the nearest permanent water supplies in Chobe National Park is sufficiently large to deter most movement except in the wet season. Beyond that, the access for Zimbabwean animals to the eastern end of the Caprivi is barred by relatively dense human settlement.

15. A request has been made to the Botswana wildlife authorities for recent survey results of roan, sable and tsessebe and the data is expected in due course.

16. Except in the extreme south-western corner of Hwange National Park where the international boundary fence has caused the deaths of a large number of animals of many species.

17. An exception may be the roan population around Kazuma Pan on the Botswana-Zimbabwe border about 100km south of Kazungula.

Sample count aerial surveys have been consistently and regularly carried out in north-western Matabeleland area of Zimbabwe since 1980 and some of the estimates¹⁸ are presented below.

The roan population in Matabeleland North was estimated at over 1,000 animals in the early 1980s. Since then it has declined drastically. The confidence intervals on the surveys in 1998 and 1999 are such that there is no significant difference between the two estimates. The present population is thought to be about 300 animals.

Sable numbers have remained fairly constant at about 5-6,000 animals. The areas of basalt soils in Matabeleland North provide optimum habitats for sable and the Matetsi Safari Area is renowned for its consistently high hunting trophy quality.

Prior to 1995, aerial surveys in Matabeleland North focussed almost entirely on elephant, buffalo and sable¹⁹ – which explains the absence of any meaningful data for tsessebe which have never been abundant in this region.

Year	Roan	Sable	Tsessebe
1989	213		
1990			
1991		7,484	
1992	31	12,713	–
1993	32	6,598	7
1994	218	5,356	–
1995			
1996			
1997	113	5,613	
1998	806	5,424	–
1999	315	5,636	316
2000	–	<i>no survey</i>	–
2001	292	5,854	80

The estimates for the three species permit little speculation about relationships with long term cumulative surpluses and deficits in rainfall – perhaps because the region enjoys a relatively high rainfall like Katima Mulilo (usually above 500mm).

With the low densities of the populations of all three species in the Caprivi and Bushmanland linkages with the Botswana population are highly desirable.

- (i) The subpopulations in the west of the Caprivi (Mahango and the western “Core Area”) are effectively isolated from the remainder of the Caprivi by the arid terrain in the central part of the Caprivi Game Reserve and also effectively isolated from Botswana by the veterinary fence along the international boundary.
- (ii) If settlement and subsistence agriculture continue to develop in the vicinity of the Kwando River in Namibia, the subpopulations in Mudumu and the western “Core Area” of Caprivi Game Reserve will become isolated and will be linked only through Botswana.

18. The blank cells are those for which data was not available at the time of writing this report.

19. It was considered that reliable aerial survey estimates could only be obtained for these three species and hence other species were excluded (this probably resulted in better survey results than many of the present day surveys which attempt to record all species!). Species such as roan and black rhino were sufficiently rare that they were always recorded when seen.

g. Behaviour

Roan, sable and tsessebe have been the subject of a number of PhD and MSc theses in the southern African region (e.g. Erb 1993, Garstang 1982, Grobler 1978, Joubert 1976, Wilson 1975) and it would be possible to include a wealth of interesting observations on the behaviour of the three species. However, the behavioural attributes which should be included in this report are those which have implications for management and therefore the discussion will be kept brief.

The question of territoriality has obvious implications for management. Species which are strongly tied to territories are likely to be vulnerable in semi-arid environments when drought or habitat changes alter their situations. Species which have the flexibility to be able to undertake local migrations when water or food is limiting may have a survival advantage. It may only be necessary for the females of the species to possess this attribute to ensure the persistence of populations.

Intra-specific aggression and competition are established behavioural characteristics of all three species and are an adaptation to ensure sufficient good quality nutrition. However, they are unlikely to play a significant rôle at the low densities at which most of the ‘wild’ populations in Namibia occur. They may be factors to be taken into account in managing enclosed populations on commercial farms or in small protected areas such as the Waterberg Plateau Park. Wilson and Hirst (1977) note that when roan and sable are constrained on small properties or areas of low quality habitat, intra-specific aggression and competition with other species intensifies.

A brief description of the relevant points relating to home ranges and territory for the three species follows.

Roan

The social structure of a roan population consists of nursery herds, bachelor groups and dominant bulls (Smithers 1983). They are gregarious animals with breeding herds usually numbering 10-20 individuals – although as many 80 have been recorded in a single group. Breeding herd home ranges vary from 60 – 100 km² (Erb 1993). Social hierarchies are well developed in both sexes. Males are forced out of the herd at 2-3 years of age and either become solitary or form bachelor groups. Breeding groups are usually led by the dominant female.

Depending on the situation, roan may or may not exhibit territorial behaviour. In some areas dominant bulls have an ‘activity zone’ rather than a territory and defend a harem of females, e.g. in Kruger National Park a specific male is usually associated with a particular female group throughout the year (Joubert 1976). On the other hand, there are instances of males establishing territories through which a group of females with their offspring may move, e.g. in the Waterberg (Erb 1993). The determining factor for which of these two modes of behaviour is adopted may be the degree of aridity. Roan may be forced to move over a large range in low rainfall areas prone to sporadic droughts – in which case it would seem pointless to defend a territory. Where the rainfall is higher and less erratic or in areas which are relatively small and constrained (e.g. in the Waterberg) the situation may be more favourable for the males to invest in territory.

Sable

As with roan, sable are gregarious, form similar-sized herds and their social structure consists of nursery herds, bachelor groups and dominant bulls (Grobler 1978). Unlike roan and tsessebe, sable bulls appear to be strictly territorial and are never attached to a mobile group of females. The home ranges of territorial males lie between 25-40ha whereas breeding herds use ranges of 2-5 km². Juvenile males are evicted from herds at about 3 years old by dominant males and, at 5-6 years of age, they are ready to establish their own territories, either by challenging dominant males or by dispersing to set up a new territories.

Tsessebe

Although tsessebe share similar social structures with both roan and sable, subtle differences also exist. Average herd sizes are generally smaller, e.g. (excluding juveniles) less than 4 in northern Botswana (Child *et al* 1972); less than 6 in Zimbabwe and Kruger National Park (Grobler 1973, Joubert and Bronkhorst 1977) and about 8 animals in the P.W. Willis Nature Reserve, South Africa (Garstang 1982).

Dominant tsessebe males establish fixed territories but, unlike sable, a specific harem herd is usually associated with each bull. This harem herd also has a fixed home range but it may not necessarily coincide exactly with that of the bull because of the partiality of females for particular habitat types. Garstang (1982) found that these harem home ranges overlapped in the hot/wet season but not in the cool/dry season. There is a high degree of fidelity to breeding herds (i.e. no new animals from other groups are permitted to join the herds). Tsessebe in Kruger National Park are fairly sedentary under such a system (Joubert and Bronkhorst 1977). However, other social structures have been observed where harem herds move between male territories at will – as in the case of sable. Child *et al* (1972) observed seasonal changes in home ranges at Kwhai in Botswana.

This ability of both roan and tsessebe to alter their system of range use according to circumstances ought to enhance their survival. However, of the three species, sable would appear to be slightly more numerous than both roan and tsessebe in the ‘natural range’ in north-east Namibia.

h. Limiting Factors

Coe, Cumming and Philipson (1976) noted that it was a distinct characteristic of large mammal communities in African savannas that, although many species might contribute to the biological diversity in any area, the major biomass contribution would be made by a limited number of species. Usually, these dominant species are bulk feeders such as elephant, buffalo and hippo. There seem to be very few cases where specialist feeders such as roan, sable or tsessebe have ever been a significant component of the large mammal biomass.²⁰ Thus, in developing a management plan for roan, sable and tsessebe, **there should be no expectations that any of these three species might ever come to dominate a savanna landscape in Namibia.**

Far from becoming dominants in southern African savannas, roan, sable and tsessebe populations have declined in many areas in recent years and given conservationists cause for grave concern. A welter of plausible hypotheses have been put forward for these declines²¹ but few unifying theories have emerged.

One such theory has been put forward by Dunham and Robertson (2001) where they demonstrate the correspondence of the observed numbers of tsessebe in Kruger National Park from 1977-2000 with the long term cumulative deficits and surpluses in annual rainfall and the late dry season rainfall (see page 24). In a study of tsessebe on a mixed cattle and game ranch in south-western Zimbabwe, the same variables accounted for an observed decline in tsessebe (Dunham *et al* 2003).²²

Wilson and Hirst (1977) saw the whole subject of factors limiting roan and sable as a complex one involving disease, malnutrition and habitat quality and thought that nutritional status might be affecting populations. Grant *et al* (2002) review the various hypotheses which have been in place at various times to explain the decline in roan, sable and tsessebe in Kruger National Park and the effects of the management actions which have resulted from these hypotheses. Dunham and Robertson (2001) and (Dunham *et al* 2003) reviewed hypotheses put forward to account for the tsessebe decline in both Kruger and Shangani and were able to dismiss all of the following – based either on the fact that they could not account for the situations over the full time period or that the data supporting them were inadequate –

- (1) Competition with other wild herbivores;
- (2) Predation;
- (3) Fire;
- (4) Effects of artificial waterpoints;

20. In southern Africa, the exceptions might be the roan population on the Nyika plateau in Malawi, the Matetsi sable population and the tsessebe populations on Shangani Ranch and De Beers Block in Zimbabwe.

21. Wilson and Hirst (1975), Harrington (1995), Harrington *et al* (1999), Grant and van der Walt (2000), Grant *et al* (2002) and RARE (2002).

22. An additional variable, that of competition with cattle, had to be incorporated into the equation in order to fit a population model closely to the observed data.

- (5) Illegal hunting;
- (6) Emigration;
- (7) Excessive sport hunting and capture offtakes; and
- (8) Direct food shortages caused by poor rainfall in any given year.

Animals were suffering from undernutrition – and might even have been killed by predators because of this – but these were proximate rather than ultimate causes. The only satisfactory explanations for the overall population performance under a range of rainfall conditions lay in the relationship with the long term cumulative rainfall deficit/surplus and the late dry season rainfall. Rainfall in late dry season appears to be critical, affecting animals' condition, survival rate, late stages of pregnancy and early stages of lactation. On Shangani Ranch, an increase in cattle stocking rates midway through the time span under consideration caused direct competition for food which resulted in a negative correlation of cattle and tsessebe biomass. The more serious indirect effects of competition with cattle in the longer term were the structural changes in habitats caused by bush encroachment following a high biomass of cattle.

In my assessment of the data from this study, the limiting factors for roan, sable and tsessebe can probably be arranged in a hierarchy –

- (1) Many of the areas in which it is being attempted to conserve roan, sable and tsessebe lie below the lower rainfall limit which the subcontinental data indicates are acceptable for the species.
- (2) In the areas which are marginal for the species from a rainfall perspective (300-400mm of annual rainfall), the performance of all three species appears linked to the long term cumulative rainfall surpluses and deficits.
- (3) A surplus in the accumulated rainfall need not necessarily produce a linear increase in population growth rates – it should rather be seen as the removal of a primary limiting factor.
- (4) When it occurs, **a deficit in the accumulated rainfall is likely to be the primary limiting factor** for roan, sable and tsessebe populations. All management efforts directed at secondary factors are unlikely to surmount this fundamentally negative effect.
- (5) Given that the rainfall regime is favourable (i.e. in a period of accumulated surplus), management efforts directed at a number of other potentially limiting factors may enhance population growth – e.g. illegal hunting, fire, provision of artificial water in specific locations.
- (6) Management interventions aimed at reducing competition with other species do not appear compatible with the general aim of increasing biological diversity. This simply results in an ongoing need for such interventions which, when they are withdrawn, result in the situation reverting to the *status quo*. However, the specific case of elephants may be an exception (see next paragraph).

- (7) Roan, sable and tsessebe are specialist feeders with habitat requirements which, despite much research, may not yet be fully understood. In the Sebungwe region in Zimbabwe all three species have been in decline for a number of years (roan and tsessebe are almost extinct). This coincides with a period where the elephant population has continued to increase and has wrought major structural changes in habitats – changes which have not been favourable to roan, sable and tsessebe. If sable and roan prefer parkland savannas with dappled sunlight shining through tree canopies to favour specific grass communities, then those conditions have gone. **The negative influence of large numbers of elephants on the habitats required by the three species is likely to be the most severe limiting factor after rainfall.**
- (8) Roan, sable and tsessebe may be susceptible to various **diseases** of which anthrax is likely to be the most serious (Pienaar 1961). De Vos and Imes (1976) document a rare skin disease contracted by sable in a holding facility in Kruger National Park. However, there is no evidence in the literature that disease has ever been a significant limiting factor for these species and, if it were, there is little in the way of management measures available to mitigate the effects (de Vos *et al* 1973). Together, predation and disease tend to be secondary factors acting on undernourished animals. Disease may differentially affect juveniles but the resultant mortality is likely to cause population fluctuations rather than any long term alterations to basic population growth rates (Sinclair 1974b).
- (9) **Veterinary fences are an important limiting factor** in their influence on movements of roan, sable and tsessebe between Botswana and Namibia. Many populations are becoming isolated as a result of the placement of fences.²³
- (10) Within the Caprivi and to the west of Khaudum Game Reserve **the *ad hoc* location and spread of human communities and their cattle** is resulting in loss of wildlife range and direct competition for grazing resources. In the background study on buffalo which preceded this work (Martin 2002), unplanned human settlement was seen as more than a ‘limiting factor’ – it was a direct threat to the long term survival of the species.

23. e.g. Mahango, Khaudum and Nyae Nyae.

2. SIGNIFICANCE OF THE THREE SPECIES

a. Conservation Significance

In the *Background Study on Southern Savanna Buffalo* immediately prior to this, the Taxon Data Sheet of IUCN's (1997) Conservation Assessment and Management Plan (CAMP) was completed as part of the work. This was not a particularly worthwhile exercise as the data sheet is intended to capture the conservation status of the global population of the species and is not applicable to subpopulations within that global population. For this reason, the CAMP form has been omitted from this study.

Under the IUCN Red Data Book system, roan sable and tsessebe are all classified as “Lower Risk (conservation dependent)” by the Antelope Specialist Group (ASG 1998) and it is evident from the distributional data of Figures 2-4 that these species cannot be regarded as threatened in any global or regional context. All three species are of conservation concern at the national level in Namibia because their numbers are low and the various subpopulations making up the national metapopulation are isolated from one another. However, it has been observed earlier in this study that most of the main body of the country is outside the limit of the rainfall range in which the three species are found ‘naturally’.²⁴ Although substantial populations of roan and sable (and, to a lesser extent, tsessebe) have been built up on private land in the main body of the country, it would be a mistake to regard these as secure because of their permanent vulnerability to rainfall regimes.²⁵ Because the areas in which roan, sable and tsessebe are found ‘naturally’ in north-eastern Namibia are spatially linked to larger populations in Botswana, they would not qualify independently for any category of threat based on population numbers.

The greatest danger to the Namibian populations of these species is the potential fragmentation which could arise if links were severed with the Botswana population due to injudicious application of veterinary control fencing or the spread of settlement and subsistence agriculture in the north east of the country.

It was difficult to argue on conservation grounds that more buffalo were needed in Namibia (Martin 2002c). However, the case for roan, sable and tsessebe is stronger. If they were to disappear from their former range in north-eastern Namibia where the rainfall conditions are favourable, this would be a loss of biological diversity and a failure of wildlife management. **Their persistence in viable numbers could be seen as an indicator of ecosystem health.** Several factors threaten ecosystems in north-eastern Namibia including the uncontrolled spread of human settlement, an overabundance of cattle (with the attendant veterinary control measures) and, perhaps greatest of the threats for roan, sable and tsessebe, is the burgeoning elephant population which is likely to modify their habitats unacceptably.

24. The Antelope Specialist Group describes the Namibian populations as ‘extra-limital’.

25. They may also become vulnerable to political factors such as a demand for land re-distribution. In Zimbabwe, thousands of animals worth millions of dollars in recently introduced wildlife populations on private land have been killed in less than two years in the course of land invasions.

b. Economic Significance

In examining the potential contribution of buffalo to non-hunting tourism income, it was stated that the presence of buffalo would be unlikely to cause any marginal increase in tourism income if they were introduced into the main body of the country or if they were more abundant in the Caprivi (Martin 2002c). The same may not be true for roan, sable and tsessebe. Substantial numbers of animals of these species are likely to impress game-viewing tourists and might result in a marginal increase in wildlife tourism. The effect is likely to be more pronounced in the Caprivi or Khaudum than in (say) Etosha, which is already a guaranteed tourist destination. However, it would be difficult to attach quantitative values to this proposition.

A corollary to the possible enhancement which substantial populations of roan, sable and tsessebe might bring to tourism in the north-east of the country is the negative impression which their disappearance from their former range would give. For knowledgeable tourists visiting the Caprivi or Khaudum, the absence of roan, sable and tsessebe is likely to be noted unfavourably.

At the outset of this study, my main preoccupation was with the precarious conservation status of these species and I gave little thought to their possible value in the sport hunting industry. However, as they are all highly prized hunting trophies (sable and roan more so than tsessebe), it would be derelict not to attempt to assess the potential economic (or financial) contribution that their inclusion in international sport hunting quotas might make.

To explore the financial and potential land use rôle of the three species, two scenarios are examined in **Appendix 4** and the results are summarised in **Table 7** below. It is a relatively simple financial exercise with no pretensions to being a full economic analysis on the lines of the various studies carried out by Barnes (see Bibliography).

Table 7. The potential impact of roan, sable and tsessebe on sport hunting income

	WITHOUT ROAN, SABLE AND TSESSEBE	WITH ROAN, SABLE AND TSESSEBE
Area	4,000 km ²	4,000 km ²
International hunting client days	664	2,952
Gross income US\$/hectare	5.90	10.06
Operating costs US\$/hectare	1.09	4.37
Net income US\$/hectare	4.81	5.69
Potential net earnings from 4,000km ²	1,923,020	2,275,280

The assumptions and methods used to perform the analysis are given in detail in Appendix 4 and summarised briefly on the next page together with a discussion of the results.

- (i) The financial modelling for sport hunting is limited to consideration of a ‘core wildlife range’ of about 4,000km² in the Caprivi.
- (ii) Two hypothetical communities of species populations which might be expected in this core area at ‘carrying capacity’ are developed – one without roan, sable and tsessebe and the other with numbers of these species at modest densities typical of southern African savannas with rainfall above 500mm. In both scenarios, the densities of species have been adjusted so that the total metabolic biomass of the wildlife community is the same (1LSU/10ha).
- (iii) Hunting quotas are set for each scenario and the hunts are ‘packaged’ into different types of hunts aimed at maximising the income possible from the available animals.²⁶
- (iv) The presence of significant numbers of roan, sable and tsessebe results in a very large increase in the number of hunter-days possible with international clients (2,952 v.664)
- (v) A safari operator needs about 180 hunter days/year to be viable and, therefore, the hunting quota when roan, sable and tsessebe are included would allow for as many as 16 operators, each using 250km².
- (vi) The annual operating costs for a safari operator are calculated for a single camp hunting operation in 250km². The capital costs of setting up the operation are included in the operating costs by depreciating the capital over 5 years and recovering one-fifth of the cost each year. The operating costs are about US\$4.37/ha.
- (vii) In the scenario without roan, sable and tsessebe, the number of international client hunter-days allows only for about four safari operators each hunting some 1,000km². It has been assumed that the operating costs calculated as outlined in the previous paragraph would not change significantly because the hunting area is increased – which reduces the unit cost to about one-quarter (i.e. US\$1.09/ha).
- (viii) In this same scenario, there is a large quota of minor plains game animals available which simply cannot be marketed on international hunts because there are insufficient premier trophy animals to which to attach them. It is assumed that these would be sold by the operator on ‘biltong hunts’ at a lower daily rate and at half the trophy fee which international clients would pay. This type of hunting would involve the safari operator in little cost because the ‘biltong hunters’ would largely fend for themselves.

The net outcome of this analysis is interesting. The gross income from the sport hunting almost doubles as a result of altering the species composition of the wildlife community to include substantial numbers of roan, sable and tsessebe. However, the costs of realising this income also increase – so that the net income produced is only some 18% higher.

26. The packaging process is entirely automated within the table using Boolean logic and simulates the normal procedure which safari operators use to market their hunts.

This analysis could have been performed in many different ways. Alternative configurations of safari hunting operations could be developed which improve the profit margin. The amounts set for trophy fees and for daily rates are not independent and are very much up to the individual operator. The hunting client will take into account the combination of both in choosing a safari (i.e. the ‘bottom line’). However, the prices which have been used are representative of the sport hunting industry in southern and central Africa.

All costs and income have been internalised within a safari operator’s budget. The apparent profit of US\$5.7/ha (a 130% profit margin for the operating cost of US\$4.4/ha) would not, of course, accrue to the operator. This is the sum from which all state revenue or community income would be derived and it is obvious that a very large surplus would be available. If the safari operator were left with a 50% profit on operations (US\$2.2/ha),²⁷ the balance available as concession rental or community income under any form of joint venture or would be US\$3.5/ha. This is almost double the projected cash income for most conservancies in the Caprivi.

This analysis shows that roan, sable and tsessebe have the potential, through their value in the sport hunting industry, to raise the overall value of net income from land significantly if their numbers can be increased to the modest levels used in the calculations. It was shown in the recent buffalo study that an increase in the numbers of this species alone could double the returns from land use in the Caprivi. Enhancing roan, sable and tsessebe populations within the species mix would further increase income. As existing wildlife uses are financially and economically more profitable than subsistence agriculture and cattle husbandry, the potential rôle of these species in a land use context is very significant.

It should also be expected that the development of substantial populations of roan, sable and tsessebe on private farms will increase their viability. Barnes (*et al* 2001) state that in the medium to long term the comparative advantages of land use based on domestic livestock can be expected to decline as international subsidies are phased out. They also point out that the comparative advantages of wildlife land uses can be expected to increase over time, due to continuing rapid expansion in international tourist markets, increasing scarcity of wildlife elsewhere, and the development of markets to capture international wildlife non-use values as income. Their results show that commercial livestock ranching has limited potential to compete economically with wildlife use because it is capital intensive and requires access to external markets.

27. Due to intense competition in the safari hunting industry in southern Africa, few safari operators are realising profits of 50% – far more common are margins below 20%.

I conclude this section by repeating some more general observations on the wildlife industry made in the previous study (Martin 2002c).

In the development of the wildlife sector, non-consumptive tourism on high quality wildlife land will give by far the greatest economic returns (Barnes 2001, Martin 1999). However, only a limited amount of land in any country is suitable for high quality game viewing tourism and, if wildlife is to compete with alternative land uses over larger tracts of land, then it is necessary to harness a range of sustainable uses to maximise the income from wildlife. Safari hunting is one such use. Martin (1995) found that whilst high quality eco-tourism could very easily realise net returns greater than US\$25/ha, the net income values for safari hunting reached a ceiling of about US\$7/ha. This may, in many situations, be the highest valued use for wildlife and the highest valued overall land use.

Safari hunting is capable of producing competitive returns from land with less capital investment than that required for non-hunting tourism and with a lower adverse ecological impact. It has other advantages. Whilst it may take several years for any non-hunting tourism venture to build up markets, the returns from sport hunting are almost instantaneous – provided a minimum population of wildlife is present. This feature may be very important in the development of local community wildlife programmes where benefits are needed from the outset in order to provide the incentives for wildlife conservation.

Barnes (*et al* 2002) observe that instability in markets for wildlife can affect sustainability and give examples to show that recent political events in southern Africa have severely affected growth in non-consumptive tourism in parts of Namibia including some of the conservancies examined in their study. Safari hunting has been demonstrated to be far less susceptible to these types of market perturbations. It may be that the political instability to which Barnes (*ibid*) are referring obliquely is the present traumatic situation in Zimbabwe. It is significant to note that whilst the Zimbabwe ecotourism market collapsed very shortly after the inception of the said ‘political events’, its safari hunting market has persisted throughout – albeit slightly reduced in volume in the 2002 hunting season. A similar situation existed during the ‘liberation war’ in the 1970s in Zimbabwe. Where there was no ecotourism activity to speak of, a viable and resilient safari hunting industry continued throughout the war. This consideration should affect decision-taking on land uses in the areas of this study.

3. STAKEHOLDING

Much of this main section on stakeholders remains the same as that written for the *Background Study on Southern Savanna Buffalo* (Martin 2000c). The new material includes a geographic identification of stakeholders based on rainfall, which follows logically from the first main section (“Biological Information”).

a. Stakeholders


The term ‘stakeholder’ is often loosely applied and may include a range of parties whose so-called ‘stakes’ differ considerably in scale. For this reason it is essential to distinguish between various degrees of stakeholders and to base decisions on the magnitude of the ‘stake’ which each party brings to the table. The primary stakeholders who are affected by the occurrence, abundance or absence of roan, sable and tsessebe in Namibia are landholders, including those with traditional landholdings. Secondary stakeholders are those who have a direct financial investment in the land and the wildlife industry. Tertiary stakeholders are those who have an interest in the conservation of buffalo but do not contribute financially to the process.

Roan, sable and tsessebe occur ‘naturally’ in areas with an average annual rainfall above 400mm. In Namibia, the three species have been introduced to a number of areas, both in national parks and on private land, where the rainfall is well below this level. A conclusion of this study is that the populations living outside the ‘natural’ range of the three species will always be vulnerable to the vicissitudes of rainfall and, in the long term, attempts to maintain the species in such areas will be frustrating and ineffective.²⁸ The policy focus of the Ministry of the Environment and Tourism should be to promote populations of roan, sable and tsessebe in areas where they are likely to succeed.

If such a policy were to be adopted, the first step would be the identification of suitable areas. Land tenure categories in the part of Namibia which lies above the 400mm rainfall isohyet are mapped in **Fig. 21** on the next page. This should define the potential primary stakeholders under any new policy. They would consist of the landholders (the State, communal lands and commercial farms) highlighted in bold colours on the map. The *de facto* situation that there are a number of roan, sable and tsessebe populations established between the 300mm and 400mm rainfall isohyets (and even a few commercial farms lying below the 300mm isohyet) is also shown on the map and it is not suggested that, under any new State policy, the stakeholders in these areas should immediately be abandoned or dispossessed of their animals. However, it should be conscious policy to avoid further investment in such areas and to promote populations in the areas where they are more likely to be successful.

Also shown on the map are those parts of communal lands above the 400mm rainfall isohyet in the extreme north of Namibia where human population densities exceed 10 persons/km² and where it would be unlikely that populations of roan, sable and tsessebe could be established.

28. It is possible that with a substantial amount of supplementary feeding, some commercial farms will be successful in maintaining populations of roan, sable and tsessebe in low rainfall areas. However, this is not seen as a sustainable long term strategy for developing the wildlife industry. If wildlife is to out-compete cattle as a form of land use it should not rely on ‘subsidies’ to species populations.



Land tenure categories are taken from Map 5.21 of the 'Atlas of Namibia' project, DEA (2002)

Figure 21. Land Tenure in Namibia above the 400mm Rainfall Isohyet

In **Table 8** below, the areas lying above the 400mm rainfall isohyet which have populations of roan, sable or tsessebe (or have had them in the recent past) are listed. In the future, these landholders should be seen as the primary stakeholders, together with any new protected areas, conservancies²⁹ or commercial farms which might become recipients of the three species.

Table 8. Areas above the 400mm rainfall isohyet with roan, sable or tsessebe populations

Land Category	Individual Areas (km ²)	Total Area (km ² rounded)
State Conservation Areas ³⁰		20,700
Etosha National Park ³¹	7,500	
Waterberg Plateau Park	403	
Khaudum National Park	3,841	
Mangetti Game Camp	480	
Mahango Game Park	200	
Popa Game Reserve	20	
Caprivi Game Park	5,500	
Mudumu National Park	1,000	
Mamili National Park	280	
Caprivi State Forest	1,496	
Conservancies ³²		10,700
Nyae Nyae	9,003	
Kwandu	190	
Mayuni	151	
Mashi	250	
Wuparo	190	
Salambala	930	
Private Land		700
La Rochelle (MU927)	100	
La Rochelle (MU8470)	100	
Kamapu -Oos	120	
Otjiwa	80	
Neu Okatjuru	100	
Okawaka	120	
Klein Waterberg	120	
	TOTAL AREA (km²)	32,100

29. In 1996, a legislative amendment provided for custodial rights over wildlife to be granted to communities on communal land subject to their forming and registering “*Conservancies*”. The provision grants partial rights for common property management and use of wildlife in defined areas. By the end of 2002, 15 conservancies had been registered, and some 35 more were in the process of being formed (Travel News Namibia 2002).

30. The areas given for the Caprivi State Protected Areas are approximate, pending gazetting notices

31. The total area of Etosha is 22,912 km² of which about one-third is above the 400mm rainfall isohyet.

32. Three additional conservancies are proposed in the Caprivi: Malengalenga, Lianshulu and Impalila

The suitable regions in Namibia for roan, sable and tsessebe are the Caprivi; Okavango, Ohangwena and Oshikoto (excluding the areas of high human density in the extreme north of these regions); the southern parts of Omusati and Oshana regions (north of Etosha Pan in the lower human density parts of these regions); Otzondjupa (excluding the lower rainfall area at extreme southerly end of the region); the northern part of Omaheke region (north of Gobabis); and a small part in the extreme east of the Cunene region south of Etosha.

Recognition should be given to the fact that the largest populations of roan, sable and tsessebe in Namibia are now on commercial farms (see Figs.9-11) – albeit that many of these populations are in less than optimum locations. It is notable that, in relative terms, the amount of land in this category is very small – in Table 8 on the previous page, the seven farms with roan, sable and tsessebe which lie above the 400mm rainfall isohyet add up to less than 1,000 km² and the total area of all the other farms which hold these species (about 18) does not amount to 2,000 km². There is considerable scope for increasing the number of primary stakeholders in the Tsumeb, Grootfontein, Outjo (extreme east), Otjiwarongo and northern parts of the Okahandja and Gobabis farming districts.

In the very large areas of communal land above the 400mm rainfall contour where human populations are at densities less than 10 persons/km² (see Figs.12 and 21) there is considerable potential to expand the number of conservancies. Areas which are strategically important for the future viability of roan, sable and tsessebe are the eastern and western Caprivi, the area of Bushmanland immediately west of Khaudum, the areas north of Etosha (at its eastern end), and the areas west and south of Nyae Nyae conservancy.

Secondary stakeholders are those who are not landholders but who are investing in the development of wildlife-based land use in these areas. This group includes hunting outfitters, professional hunters, hunting guides, tourist lodge operators, businesses involved in processing trophies and, in general, all support systems for the wildlife industry. The group would also include all those organisations assisting to develop conservancy programmes, which are listed in full in the background study on buffalo (Martin 2000c). The large international donors investing funds in community based wildlife management tend to direct their investments through local NGOs but should also be seen as secondary stakeholders.

Tertiary stakeholders would include various wildlife societies and individuals concerned in general for wildlife conservation and the tourists who enjoy the recreational opportunities of Namibia.

b. Stakeholder Institutions – Present and Future

The progress which Namibia has made in developing policies and legislation which empower landholders to manage their wildlife resources both on commercial farms and in communal lands was noted in Martin (2000c) – as was the impressive record of development of the wildlife industry and the positive spirit of co-operation amongst the State, NGOs and private sector towards the larger goals.

In the same study, some disquiet was expressed that the initial positive steps towards devolution of rights over wildlife, for both communal land conservancies and private land, were not being followed by further progress. Institutional development in the wildlife industry in Namibia appeared stuck in a state of ‘aborted devolution’ (Murphree 1998).

The legislation (RN 1975) classifies various wildlife species into categories of “game” – such as “Specially Protected Game” (Schedule 3) “Protected Game” (Schedule 4) and “Huntable Game” (Schedule 5). Roan, sable and tsessebe are “Protected Game” – which implies that permits are needed from the Minister for these species to be hunted (and, presumably, for the capture or sale of live animals of the species). This type of legislation acts negatively towards the conservation of species because it fails to align authority, responsibility and incentives (Murphree 2000).³³ Authority without responsibility is meaningless or obstructive, responsibility without authority cannot be effective and, without responsibility or authority, there are no incentives to invest, manage or control. The governing hypothesis is, that given full authority over wildlife resources, the incentives will be present for landholders to use them sustainably.

Despite the fact that cattle are of a lower economic value than wildlife on commercial farms in Namibia (Barnes and de Jager 1995), the present system of mixed cattle and wildlife farming is likely to persist. The fact that Namibian farmers do not enjoy the same rights over their wildlife as they do over their cattle could be the single most important factor which is slowing down the progress of farmers converting to ‘pure’ wildlife systems. In competitive land use situations it is imperative that the rights of a wildlife farmer are no different to those which he enjoys over his cattle if he is to make choices which value one resource above the other.

Other perverse incentives are present in the legislation. The rights over “Huntable Game” which are conferred on private landholders if their properties are fenced to certain standards³⁴ would seem to mitigate against the desirable amalgamation of adjacent wildlife properties. Where roan, sable and tsessebe are concerned, it is evident that many of the populations on commercial ranches are being held at densities which exceed the carrying capacity for low rainfall areas. The formation of large “conservancies” (of the order of 1,000km² or more) would greatly enhance the likelihood of survival of the three species in times of drought and improve their breeding performance when rainfall conditions are favourable. Rather than pursue a “custodianship policy” under which commercial farmers are allowed to keep certain valuable wildlife species which are seen as the property of the State, greater incentives would be provided if the State used species such as roan, sable and tsessebe to promote the formation of larger amalgamated wildlife areas amongst commercial farmers – without attempting to retain controls on the use of the species.

The same principles apply to conservancies.

33. It is of interest that the only species which are a conservation failure in Zimbabwe are those which have been classified as “Specially Protected Species” (there are no categories equivalent to the Namibian “Protected Game” and “Huntable Game”). Roan antelope are Specially Protected Species in Zimbabwe and this categorisation has been responsible for their decline. Because the State retains the authority to control hunting and sale of live animals of the species, there is no investment in increasing their numbers.

34. In primary legislation, it is surprising to see detailed specifications for game fences. This type of technical detail belongs in regulations and subsidiary legislation.

To conclude this section on future stakeholder institutions, it would seem there are good reasons to examine the rôle of State Protected Conservation Areas in relation to neighbouring communities (including commercial farmers). In many places in the optimum range for roan, sable and tsessebe, there are configurations of parks and their neighbours which, as they stand, are not ideal. In the Caprivi, in particular, the mosaic of small parks and small conservancies would benefit from being amalgamated into larger, more viable blocks of land under wildlife management. Opportunities arise in respect of Khaudum and Nyae Nyae Conservancy to benefit from joint management on a large scale. Corbett and Jones (2002, p19) raise the option of partnerships in management and revenue-sharing arrangements and, inevitably, this would mean the development of new co-management institutions (Ruitenbeek and Carter 2001).

At the coming World Parks Congress,³⁵ the agenda will be dominated by the topic of relationships between protected areas and their neighbours. At a recent workshop³⁶ on *Local Communities, Equity and Protected Areas* aimed at preparing for the World Congress, the new Director-General of IUCN gave an inspirational address which indicated that the time had come to find new models for protected areas which were more resilient than those of the past. This is a challenge where Namibia could lead the way for other countries in the southern African region.

c. Towards Trans-Boundary Institutions

The complexity of the proposed 'Four-Corners Trans-Frontier Conservation Area' is daunting (Martin 2002a). To develop institutions involving not only the national governments of five countries (Angola, Botswana, Namibia, Zambia and Zimbabwe) but also the other primary stakeholders is a formidable task made more complicated by the different legal systems and institutional approaches which have already evolved in each country. This complexity was recognised in the round-table discussions at the inception of this study and a pragmatic approach was agreed upon whereby the larger vision of a massive trans-frontier area, whilst being recognised as an ultimate goal, should be preceded by the building of a number of incremental initiatives aimed at collaboration between Namibia and its immediate neighbour, Botswana.

Following the completion of the background study on buffalo (Martin 2002), the Ministry of the Environment, the Namibia Nature Foundation, the WWF LIFE Programme and the Botswana Department of Wildlife and National Parks initiated the first step towards collaboration by holding a workshop in Kasane, Botswana, at the end of November 2002 to discuss specific areas of co-operation with regard to buffalo and to consider further wildlife species which merited joint management.

A notional institution for collaboration between Botswana and Namibia was presented at the meeting for discussion purposes (**Fig.22**, page 55). The design of the institution was specifically aimed at the matching of ecological, functional and jurisdictional scales as outlined by Murphree (2000). The initial focus of the institution was on the Caprivi but, when roan, sable and tsessebe become involved, the institution should take into account stakeholders on either side of the north-south international border in the region of Khaudum and Nyae Nyae Conservancy.

35. Durban, South Africa, 8-17 September, 2003.

36. Workshop titled *Local Communities, Equity and Protected Areas* hosted by The Programme for Land and Agrarian Studies (PLAAS), Africa Resources Trust (ART) and IUCN, 26-28th October, Pretoria, S. Africa.

A key question is whether at the international level (i.e. between Botswana and Namibia) representation will be confined to government representatives or whether other primary stakeholders will participate (i.e. local community representatives from both countries).³⁷ The proposal in Fig.22 is that local communities should be represented but, ultimately, the decision on this issue lies between the two governments. The features of the organogram are as follows –


- (i) The organisation consists only of primary stakeholders and only involves two levels. A third level (not shown in the diagram) is the individual membership of each conservancy in Namibia and each community area organisation in Botswana.
- (ii) In Namibia, the individual conservancies would delegate limited powers upwards to an ‘Association’ which will represent them at the second level (CAs). This could comprise the existing Communal Lands Boards or be an association created especially for this purpose. The arguments in favour of using the Communal Lands Boards are that the interests of communities who have not formed conservancies would also be represented.
- (iii) A similar association would need to be identified on the Botswana side of the border to represent the various areas under community wildlife management.³⁸ This could be the relevant Land Board.
- (iv) The individual parks in northern Namibia and Botswana report to their Directorates which are represented on the second tier in each half of the structure.
- (v) In Namibia, the Directorate of Scientific Services is also represented in the second tier. In Botswana, the equivalent agency is contained with the Department of Wildlife and National Parks.
- (vi) The veterinary authorities from both countries are also represented at the second level.
- (vii) These two groups at the second level from Botswana and Namibia meet to constitute the international ‘institution’ which addresses joint management issues.

It could be argued that many more parties should participate in the final bilateral forum including more senior government representatives. In line with both governments’ efforts to decentralise, it seems more logical that this forum be treated as a technical and advisory panel which reports back to the relevant ministries on matters which may require high level decisions. If the principles of delegation upwards and accountability downwards are adhered to, there is no reason why all of the representatives at the national level cannot report back their particular constituencies rather than overload the international forum with unnecessary numbers. Finally, if it is agreed between the two delegations, there is no reason why any observers who may contribute to the discussion are not invited to the forum.

37. There was no such participation at this inception meeting in Kasane. For Botswana, representation was confined entirely to wildlife department staff and, on the Namibian side, only government staff and NGOs participated. It is possible in future meetings that this representation will be broadened.

38. The workshop identified a potential trans-boundary community collaborative opportunity in the Salambala Conservancy and the Chobe Enclave.

Figure 22. A Notional Institution for Botswana-Namibia Management of Shared Wildlife Species Populations



Key to Acronyms used in the diagram – see text for a fuller explanation of the structure

Namibia: CAs – Conservancies Association
 DSS – Directorate of Scientific Services
 DPW – Directorate of Parks and Wildlife
 DVS – Directorate of Veterinary Services

Botswana: DWNP -- Department of Wildlife and National Parks
 DAHP – Department of Animal Health and Production
 CAs – Community Areas Association

4. MANAGEMENT

The main finding of this study is that roan, sable and tsessebe are very much at the mercy of rainfall and that this effect is most pronounced near the limits of the ‘natural range’ for all three species, i.e. in areas with an annual rainfall lower than 400mm. With annual rainfall in most of Namibia falling below 400mm, the initial reaction to this finding is that there is little that can be done to secure the future survival of the three species.

There is, in fact, a great deal that can be done. Firstly, about one-quarter of the country does have an annual rainfall which is above 400mm and this zone must be the focus for conserving roan, sable and tsessebe. Within this zone, all of the conventional measures that enhance wildlife populations such as effective law enforcement, fire management and provision of water should be in place. But there are also opportunities to be pro-active within this zone. The establishment of new populations of roan, sable and tsessebe on any land where they are likely to be secure should proceed as rapidly as circumstances will permit. In this way, risk will be spread and the likelihood of ensuring the survival of the species is far greater.

Finally, continued collaboration with Botswana on the shared populations of these species is likely to produce benefits related to scale. Given that spatial linkages can be maintained and enhanced, the probability of the three species going extinct simultaneously throughout the region is minimal.

a. Present and Future Management of Roan, Sable and Tsessebe

(1) Protecting Existing Populations

Roan, sable and tsessebe will benefit from the general management measures aimed at conserving wildlife in State Protected Areas³⁹ and Conservancies. Within State Protected Areas a major effort is being made at present to contain illegal hunting, control fires and, in general, to implement park plans. The present MET staff numbers, equipment and infrastructure in many parts of Namibia are insufficient to meet the challenges (PW 1998, page iii) but improvements are taking place in all these aspects. Martin (1996) examined the minimum requirements of game guards and budgets for effective functioning of State Protected Areas and this analysis is summarised in **Appendix 5**.

Based on the formulae of Appendix 5, notional budgets have been developed for the protected areas which lie in the north of Namibia above the 400mm rainfall isohyet (**Table 9**, next page). The required operating budget for all of the State Areas is slightly under **US\$3 million** per annum.

It is of interest to note the effect of managing several small areas in the Caprivi rather than a single large area. The operating costs required for managing the Caprivi Parks (including the Forest Reserve and the Kwando Triangle) as individual areas is almost US\$2 million, i.e. almost two-thirds of the total budget. Were these areas to be managed as a single unit, the budget required for the Caprivi is halved and the total budget is reduced to less than US\$2 million.

39. Uncertainties surrounding the exact designation and final boundaries of many State Protected Conservation Areas in Namibia (Mendelsohn and Roberts 1997, p7; PW 1998) add to the law enforcement difficulties of the wildlife department. Clearly, a resolution of these issues will set the base line against which many other land use and conservation plans can be developed.

Table 9. Required Operating Costs for State Protected Areas in the Potential Range for Roan, Sable and Tsessebe

State Conservation Areas	Total Area km ²	Required Number of Guards	Required Annual Operating Budgets - US\$
Popa Falls Game Reserve	20	5	122,000
Mahango Game Park	200	15	177,000
Mamili National Park	280	17	193,000
Waterberg Plateau Park	403	21	216,000
Mangetti Game Camp	480	22	228,000
Mudumu National Park	1,000	32	300,000
Caprivi State Forest	1,496	39	359,000
Khaudum National Park	3,841	62	586,000
Caprivi Game Park	5,500	75	727,000
TOTALS . . .	13,220	288	2,908,000

Notes:

The eastern end of Etosha National Park, which is listed amongst the protected areas in Table 8, has been left out of this table because it is assumed that adequate operating budgets for the whole of Etosha are in place.

The operating costs for the Kwando Triangle are assumed to be included in the budget for the Caprivi Game Park

Costs of managing Caprivi Parks

State Conservation Areas	Total Area km ²	Required Number of Guards	Required Annual Operating Budgets - US\$
Popa Falls Game Reserve	20	5	122,000
Mahango Game Park	200	15	177,000
Mamili National Park	280	17	193,000
Mudumu National Park	1,000	32	300,000
Caprivi State Forest	1,496	39	359,000
Caprivi Game Park	5,500	75	727,000
TOTALS . . .	8,496	183	1,878,000
<i>Managed as a single unit . . .</i>	<i>8,496</i>	<i>93</i>	<i>963,000</i>

These budgets set a critical threshold. Where the State provides an annual operating budget equal to or greater than required, there is a high probability that the area will be adequately managed and conserved. Where budgets are lower than the amounts needed, it is unlikely that a wildlife agency will be able to handle any determined onslaught by illegal hunters. Shortfalls in the budgets provide a strong reason (but not the only reason) for seeking new institutions involving partnerships with neighbouring communities.

Progress with Conservancies in the Caprivi is impressive. Many of the apparently overwhelming conservation tasks expected of State wildlife agencies are likely to be reduced when protected areas are surrounded by functioning community land use systems based on wildlife and natural resources. The question which a State wildlife agency needs to ask itself is whether the combination of stand-alone parks (which may be seriously underfunded) and conservancies together provide the right land use planning framework for the future. In the section of this report on Stakeholders (page 53), the potential rôle of State Protected Areas in catalysing land use based on wildlife over a wider area was raised. Moves towards this will require continued innovation from enlightened Namibian bureaucrats and a re-definition of classic protected models.

(2) Land Use Planning

In the previous study on buffalo (Martin 2002c), it was argued that *ad hoc* settlement, clearing of land for agriculture and cattle grazing were resulting in a large loss of potential range for buffalo. Throughout the area of Namibia above the 400mm rainfall isohyet, this factor is likely to exert a strong negative influence on roan, sable and tsessebe populations. The process is not one which the Ministry of Environment and Tourism would normally be expected to influence greatly. However, unless the Ministry does involve itself in a pro-active manner in land use planning, it is likely that much of the high potential for land use under wildlife (which should be a national priority to realise) will be foregone.

(3) Veterinary Fences

The influence of veterinary fences on the well-being of roan, sable and tsessebe populations is less critical than for buffalo (Martin 2000c). Nevertheless, the fences along the international boundary between Namibia and Botswana undoubtedly impede movement of the three species and act against the maintenance of spatial linkages between the subpopulations. The effects are likely to be most severe in the vicinity of Khaudum, Nyae Nyae, Mahango and the Caprivi Game Park.

(4) Establishing New Populations

Translocations of roan, sable and tsessebe to commercial farms have proved a success,⁴⁰ although many of these farms are located in areas with lower rainfall than might be preferred. The establishment of roan and sable populations in Waterberg Plateau Park has also been successful and it is of interest that this park lies close to, but above, the 400mm rainfall isohyet. It is the main reservoir for the Ministry of Environment to draw from in establishing further populations of roan and it makes sense to remove animals continuously from the population to ensure that it remains below carrying capacity and able to breed at the maximum rate.

It is strongly recommended that the Ministry of Environment continue to pursue an aggressive policy of establishing new roan, sable and tsessebe populations. The features of such a policy should be –

40. The present numbers of these species on private land are estimated as: roan– 404, sable – 778, tsessebe – 175.

- (i) New populations will only be established in areas with an annual rainfall higher than 400mm per annum (see Fig.21, page 49);
- (ii) The chosen locality for any new introductions should be secure (see previous two subsections) so that factors such as illegal hunting, loss of range and competition with cattle are not likely to impede population growth;
- (iii) The habitats in the selected areas should be suitable for the introduced species – with roan and sable this unlikely to be as great an issue as with tsessebe, whose habitat preferences are fairly narrow and ecotonal.⁴¹
- (iv) The category of land on which the new population is being established (i.e. whether it is a State protected area, conservancy or commercial farm) should be of secondary importance and should not be subject to political influence;
- (v) In establishing new populations of roan, sable and tsessebe, the Ministry should use the opportunity to promote the conversion of larger areas of land to wildlife management. Accordingly, no special conditions (such as those pertaining to the “Custodianship Scheme”) should be attached to the introductions.

(5) Illegal hunting

Levels of illegal hunting could affect the survival of roan, sable and tsessebe on both State Land and Conservancies. The population model developed in **Appendix 1** has been used to explore the maximum illegal harvest which a population of roan, sable or tsessebe could sustain. It is assumed that mortality would affect both sexes and all ages equally. The calculation of the number of years a population would take to reach 2,500 animals is based on an assumed starting population of 250 animals (which is not far off the present estimates for the populations of all three species in their “natural range”), i.e. the time it would take for the population to increase ten-fold.

Illegal harvest %	0	1	2	3	4	5	6	7	8	9	10	11	12
Rate of population growth %	13.6	12.5	11.4	10.2	9.1	7.9	6.8	5.7	4.5	3.4	2.3	1.1	0.2
Years to reach 2,500 animals	18	20	22	24	27	31	34	42	53	69	102	211	∞

The assumptions in this analysis are that the generic population model is suitable for all three species and that the model applies only in a situation where rainfall is above 400mm and is not in any major long term rainfall deficit mode (see page 24). Without any illegal hunting the population grows at slightly under 14% per annum and it can sustain a maximum offtake of about 12%. The higher the proportional offtake, the lower is the growth rate of the population and the effects become severe above a 6% offtake.

41. Dunham and Robertson (2003) give a detailed description of the habitats for tsessebe on Shangani Ranch in south-western Zimbabwe. The population of tsessebe on this and neighbouring ranches was, until recently, the largest single subpopulation in Africa.

To examine rates of population decline when the illegal harvest exceeds 12%, it is not useful to examine percentage offtakes because these result in a lower and lower number of animals being killed as the population declines so that the population tends to stabilise at some low level. A more realistic examination of rates of decline for unsustainable harvests has been done with a fixed number being removed from the population each year which inevitably results in extinction. In the table below, the number of years to extinction is shown for various fixed offtakes from a starting population of 1,000 animals.

Illegal harvest (% of 1,000 animals)	14	15	16	17	20	25	30	40	50
Fixed annual offtake	140	150	160	170	200	250	300	400	500
Years to extinction	29	20	15	13	9	7	5	4	3

(6) Sport Hunting

I have used the population model of **Appendix 1** (page 66) to explore the effects of hunting quotas on roan, sable and tsessebe populations (**Fig.23** below). Again, the assumptions are that one model suits all. Hunting selectivity is centred on the 8-9 year old males, with 40% of trophies coming from this age group, but the rest spread fairly evenly over the range of age classes. The model works by attempting to take the available quota of trophy animals from the various age classes in the proportions set by the selectivity profile (i.e. if the quota was 100 animals it would take 5 animals from the 5 year old age-class, 10 animals from the 6 year-old age class, 15 animals from the 7 year-old age class . . . and so on). However, if there are insufficient animals in any age class to meet the quota demand, the animals are then sought in the age class immediately below it (as would happen in practice). The percentage quotas apply to the total population and it is assumed that males under 5 years old would not be hunted.

Fig.23. Effect of hunting quotas on age structure of adult males in the population



As the quota is increased from zero, the older age classes are ‘cleaned out’ very quickly. A quota of 1% results in all animals older than 12 years being removed from the population, 1.5% removes all animals 9 years and older and, at 2%, all 8 year-olds have gone, leaving only 7 year-olds and younger males to breed. This would suggest that **sport hunting quotas for roan, sable and tsessebe should never exceed 2%**. The annual recruitment to the part of the age pyramid from which males are hunted is very low – the proportion of males recruited annually to the 5 year age class is about 3% of the population and, to the 8 year-old age group, it is about 2% of the population. Thus a hunting quota of 2% will claim every male 8 years old and over.

The sex ratio in the population in the absence of hunting is 1♂:2♀ and, as hunting quotas increase, this shifts in favour of females. With the quota at 2%, the sex ratio becomes 1♂:3.3♀.

I conclude this section on sport hunting by discussing the methodology for setting quotas and monitoring sustainability. It is not necessary to know the numbers of animals in the population in order to set sustainable quotas – indeed a system based on population estimates is likely to be far less robust than an adaptive management system (Holling 1978, Bell 1986, Martin 1999) because, firstly, the confidence intervals on population estimates are very large and, secondly, the area of interest is not the total number of animals in the population but the number of adult males older than (say) 8 years – which is only about 3% of the population.

The key parameter to be monitored is the age of trophies taken from the population. If a criterion is set that there should always be a sufficient number of prime breeding males, then the requirement is that amongst the trophies (regardless of the selectivity of the hunting regime), there should be a representative number of males in the age classes above 7 years old. As soon as the cohort of hunted animals is missing all of the age classes older than 7 years of age, this is a robust indicator that the population is being overhunted and the quota should be reduced. An initial quota might be set by the crude method of applying 2% to the population estimate but thereafter that quota should be adjusted upwards or downwards by the ‘hard data’ (as opposed to the ‘soft data’ of population estimates) which comes from measuring trophies. Methods of ageing roan, sable and tsessebe from their dentition or horns are available (Grobler 1979,1980; Joubert 1976; Child *et al* 1972; Huntley 1973) – which could easily be applied by local community monitoring staff with some training. It is logical that conservancies should take on this monitoring rôle in all areas where the species are hunted in conservancies since it is effectively their resource being managed.⁴² In State protected areas where there is hunting this would be the responsibility of the parks staff.

The principle can be applied to the hunting of any species. Adaptive management is a better methodology than the blind application of percentage offtakes to populations since it tests the underlying hypotheses about population age structures and the sustainability of hunting quotas. In the case of roan, sable and tsessebe there is perhaps a more important reason for monitoring quotas by this method which is discussed in the next subsection.

42. The development of conservancy monitoring systems by the NACSO unit (Namibian Association of CBNRM Support Organisations) is impressive and may be well-suited to carrying out this specific monitoring.

(7) Monitoring

In general, most monitoring activities should be applied within an adaptive management framework. Bell (pers.comm. 2000) said “*In the early stages many of us saw adaptive management as a research process to reduce uncertainties. These days it should be stated as part of the definition of adaptive management that intensive levels of research or monitoring which will result in a system being too expensive, and hence unsustainable, cannot and should not be attempted.*” An example has been given of the use of adaptive management to set hunting quotas where there is a focussed objective and monitoring is aimed at realising that objective. A second situation arises, specifically within the Transboundary Species Project, with the need to assess whether the objective of increasing roan, sable and tsessebe populations is being achieved.

Present air survey techniques are not suited to precise or accurate estimates of roan, sable and tsessebe populations for the reasons given on page 24. However, this method is the best there is at present. The population estimates for all species in the Caprivi, for example, would be greatly improved if annual surveys using the standard methodology of Craig (2000) were done at a fairly high sampling intensity (10-20%) using the same strata on every survey and, where possible, the same observers. The budgets outlined in Table 9 (page 57) contain a provision for such a survey for each area every year.

There is an alternative (or additional) method that might be attempted for roan, sable and tsessebe which makes use of the monitoring data obtained from sport hunting. The data in Fig.23 show the shape of the age pyramid to be expected at different levels of hunting quotas. For example, if the hunting quota is 2% there are unlikely to be any males older than 8 years in the population. If the ages of the animals taken in any year are accurately determined and if it is found (say) that there are 7 year-old animals in the population but no 8 year-olds, one might be justified in assuming that the actual numbers of animals killed in the year concerned is **2% of the population**. This permits a reverse calculation to be carried out: if 10 animals were hunted and this, according to the age structure model should be 2% of the population, then total population would be 500 animals.⁴³

Obviously a large number of assumptions go into this method – including the assumption that the adult male survival rate and the selectivity for trophies are close to the values used in the model and that the age determinations can be carried out with some accuracy. However, over several hunting seasons through some iterations with the model and through deliberately pursuing a course of active, adaptive management⁴⁴, it might be possible to simultaneously refine the model, obtain better estimates of the populations from which the hunting trophies are derived and arrive at optimum hunting quotas.

43. A drawback of this type of calculation is that the quotient consists of a large number divided by a small number and the corresponding scope for error is very large.

44. i.e. the hunting quotas are deliberately varied, even in some cases to levels which are likely to be too high, in order to learn more about the population characteristics.

(8) Elephants

In the discussion of factors possibly limiting roan, sable and tsessebe populations (page 40), the large elephant population in the Caprivi (5,000-10,000 animals) was put forward as perhaps being responsible for fundamental structural changes to habitats. In areas where annual rainfall is 500-600mm, elephant densities greater than 1/km² result in marked changes to woody vegetation and it can be presumed that the grazing sward is also affected. There have been no population reductions of elephant in either northern Botswana or Caprivi as part of ecosystem management in recent times (if ever) and this management option could be considered. It is a topic which should be discussed jointly with the Botswana authorities.

(9) Artificial water

More habitats could be made available for roan, sable and tsessebe by the supply of artificial water – particularly in the Caprivi Strip and Bushmanland. All of these species are water-dependent⁴⁵ and seldom move further than a few kilometres from surface water (Table 2, page 9). In the Caprivi, this means they are tied to the large rivers for a large part of every year. This limits the ability of the populations in the eastern and western ends of the Caprivi Game Reserve to maintain contact and, in conjunction with the veterinary fences along the Botswana border and a hostile environment in Angola, could result in the total isolation of various subpopulations.

The development of game water supplies in the large Kalahari Sands area of the Caprivi Game Reserve would not be simple: Mendelsohn and Roberts (1997, page 39) show the average depth of water below the surface as varying from as much as 300 metres in the west of the Caprivi Strip to 35 metres in the east. A number of boreholes have been sunk in the area but most are non-functional or would only provide small quantities of water. However, this latter feature might prove valuable to roan, sable and tsessebe: if large amounts of water were available it is highly likely that the water points would be captured by elephants and large buffalo herds.

(10) Fire

Roan and sable are not particularly attracted to burns but tsessebe are – although they will seldom move more than short distances in order to feed on a burn (Joubert and Bronkhorst 1977). Fire can be a valuable tool to improve habitats for tsessebe: where bush encroachment has occurred through excessive cattle grazing, judicious burning may return areas to grassland.

In the Caprivi, the marginal gains which roan, sable and tsessebe might get from careful use of fire are more likely to be nullified by the loss of grazing caused by the existing fire regimes. Mendelsohn and Roberts (1997, pages 24-25) present a compelling picture of the gravity of the fire situation with burns commencing as early as April each year and continuing until December when over 60% of the vegetation has been burnt and the total count of individual fires may have exceeded 3,000. It seems that at one time there was an extensive network of firebreaks in Caprivi to control fires and it would obviously be beneficial if these could be resuscitated.

45. Garstang (1982) observes that tsessebe have ‘aesthetic preferences’ for certain types of surface water and will generally avoid drinking at concrete water troughs.

b. Transboundary Issues

A number of areas have been identified in this report where collaboration between Botswana and Namibia could enhance roan, sable and tsessebe populations. These are presented below in the form of a possible agenda for the next joint workshop with Botswana.

Botswana's roan population is about 1,500 animals, its sable population is about 3,000 and its tsessebe population is one of the largest in Africa – of the same order as Zambia's at about 10,000 animals. Together Botswana and Zambia hold more than two-thirds of Africa's tsessebe. Thus, Namibia stands to be the greater beneficiary from co-operation with Botswana on management issues than *vice-versa*. Namibia's primary conservation objectives are to increase numbers of roan, sable and tsessebe and to avoid fragmentation of the populations. Maintaining spatial linkages with Botswana will be important in achieving this.

The question of the scale at which roan, sable and tsessebe populations should be managed is an important one. In this course of this study it has not been possible to identify discrete subpopulations but it is known that movements of all three species are considerably less than is the case for buffalo. Therefore, the scale under consideration at the outset of this collaborative process need not necessarily embrace the full northern Botswana populations of the three species but could focus on the animals in Namibia and Botswana which are located within a certain distance (e.g. 50km) of the international boundary. Later, it may be possible to refine management to specific subpopulations.

Perhaps the key determinant of roan, sable and tsessebe numbers is rainfall and, if this is so, the remaining factors might seem to be of secondary importance. I do not personally believe this is so and identify a number of management issues which could make a substantial difference and where co-operation would be worthwhile. Most of the issues remain the same as those affecting buffalo but they differ individually in their priorities. They are listed below in order of importance.

(1) Elephants

The impact of the very large elephant population in the project area (more than 100,000 animals) on roan, sable and tsessebe habitats is viewed in this study as the most important factor after rainfall. Elephant management is a high-level issue where technical collaboration is essential.

(2) Introductions

Botswana might be able to provide animals from their large reservoir of tsessebe for introduction to identified project sites in Namibia in the areas which have been identified as having the greatest likelihood of success in re-establishing populations. Although roan and sable in Botswana are less abundant, there may be potential for introductions of these species, too – or the deliberate establishment of trans-border populations of all three species in specific localities.

(3) Illegal Hunting

Levels of illegal hunting in the Caprivi are higher than in northern Botswana and, if these cannot be contained, it will have a deleterious effect on Namibia's resident roan, sable and tsessebe populations. It is less likely to affect the larger populations of these species in Botswana, by virtue of the localised nature of the species' distribution (unlike the buffalo situation). Nevertheless, there may be collaborative measures that could assist in reducing illegal hunting in both countries.

(4) Veterinary Control Measures

Veterinary control measures are less of an issue with roan, sable and tsessebe than they are with buffalo. However, they are a factor acting to disrupt linkages between the Namibia and Botswana populations and should still be addressed with some priority. The trend in Namibia towards isolated subpopulations in Caprivi, Khaudum and Nyae Nyae is of concern. Scott-Wilson (2000) put forward four options to mitigate the effects of veterinary fences in northern Botswana and decisions are still awaited on these options – or an alternative solution.

(5) Population Estimates

The inadequacy of present air survey techniques for counting roan, sable and tsessebe is highlighted in this study and is reflected in the high confidence limits of both the Botswana and Namibian population estimates. In most years, Botswana conducts a national air survey of all of the northern wildlife areas which is done to the highest scientific standards. It would be cost-effective to extend the coverage of this survey into the roan, sable and tsessebe range in Namibia, with the extra costs being met by Namibia. This could be a major step forward in standardising air surveys between the two countries.

(6) Fire

The Caprivi suffers from an excessive burning regime every year. Whilst few of these fires originate from Botswana, this may be an area where co-operative effort would result in a reduction in the number and extent of fires.

(7) Liaison on Hunting Quotas

It is unlikely that excessive sport hunting quotas in either Namibia and Botswana would be likely to affect each other's safari hunting industry significantly because of the localised subpopulations which make up the roan, sable and tsessebe distribution. However, it is possible that in specific localities on either side of the international border there are good reasons to cooperate on joint hunting management. This is an area of liaison which would require little effort and could produce significant economic and conservation gains. In the areas on either side of the international border where hunting is taking place from what may be the same herds, there is good case for developing local institutions at the appropriate scale which would enable the proceeds from an overall quota to be shared proportionally amongst the participating community areas.

A Generic Population Model for Roan, Sable and Tsessebe

Using the broad reproductive parameters given in Table 5 of the main report (page 12), a simple population model has been constructed using a computer spreadsheet which enables analysis of population growth rates and age structures.

START	0												Nominal starting population:	1,000
YEAR	1													
AGE	1	2	3	4	5	6	7	8	9	10	11	12	TOTALS	
	Starting Cohort													
Males	114	76	53	40	32	25	20	16	12	8	3	0	399	
Females	114	96	80	68	58	50	43	36	29	18	8	1	601	
	Running Cohort (population from previous year)													
Males	114	76	53	40	32	25	20	16	12	8	3	0	399	
Females	114	96	80	68	58	50	43	36	29	18	8	1	601	
Fecundity	0.00	0.00	0.75	0.95	0.97	0.98	0.98	0.97	0.95	0.50	0.25	0.00		
Calves	0	0	60	64.6	56.3	49	42.1	34.9	27.5	9	2	0	346	
	Population after births													
Males	173	114	76	53	40	32	25	20	16	12	8	3	572	
Females	173	114	96	80	68	58	50	43	36	29	18	8	773	
	Population after annual mortality													
Survival ♂	0.75	0.75	0.80	0.85	0.90	0.90	0.90	0.90	0.90	0.70	0.50	0.10		
Survival ♀	0.75	0.95	0.95	0.97	0.97	0.98	0.98	0.95	0.90	0.70	0.50	0.10		
Males	130	86	61	45	36	29	23	18	14	8	4	0	454	
Females	130	109	92	78	66	57	49	41	33	21	9	1	686	
	TOTAL POPULATION												1140	
	RATE OF GROWTH %												14.00	
	ADULT SEX RATIO												2.01	

The model behaves in a manner similar to the Leslie matrix (Leslie 1984) but the calculations of births and deaths are separated into successive operations because it is designed to cycle within the row operations of a computer spreadsheet. The model operates as follows –

- (i) The starting year is set to zero, and a nominal starting population and average female fecundity are set in the indicated cells.
- (ii) In the two rows immediately following, the starting population is divided into equal numbers of males and females and further divided into numbers in each age class which approximate a stable age distribution.

- (iii) In the first year of the model, this cohort of males and females is transferred to the next two rows of the model (“Running cohort”). On each successive cycle of the model thereafter, the running cohort derives its population values (males and females) from the last two lines of the model.
- (iv) On each cycle of the model the individual number of females in each class is multiplied by the fecundity in the cell immediately below it to give the number of calves produced by each female age class.
- (v) The number of calves is summed at the end of the row, divided by two and inserted in the first two cells of the next two rows (“Population after births”). At the same time the number of males and females in each cell of the running cohort above is moved forward by one year and inserted in the cells following the one year old age class, i.e. at the same time as the births occur each animal in the population ages by one year.
- (vi) The individual male and female cells of the “population after births” are then multiplied by the survival values in the next row to give the “Population after annual mortality”.
- (vii) The population is then totalised and the growth rate is calculated using the increase in the population over the number at the start of that particular cycle (which is the total number of animals in the “Running cohort”).
- (viii) The cycle is then repeated as many times as desired (usually until the age structure becomes stable and the growth rate does not change from year to year).

The female survival values have been adjusted from their nominal starting values to give a sex ratio of 1 male : 2 females when the population has reached a stable age distribution.

The model has been amplified on the next page to examine the effects of illegal hunting and sport hunting. These two ‘treatments are inserted into the spreadsheet after the population has undergone its first step in the annual cycle. The model sequence is –

Births → Age by One Year → Illegal Hunting → Sport Hunting → Natural Mortality

For illegal hunting it is assumed that there is no selectivity – all animals in the population have an equal likelihood of being killed. When population numbers are low, the spreadsheet may give errors due to rounding and the actual number of animals killed may be more or less than the expected number. To avoid this effect, in the original spreadsheet compensation is performed using random numbers to round up or round down the numbers until the expected number matches the actual number of deaths. This is not shown in the table overleaf.

The modelling process for sport hunting is slightly more complicated. Firstly, it is assumed that the quotas will be set as a proportion of the total population after it has enjoyed its annual increment of births, i.e. it does not take into account the numbers of animals killed illegally and, in this respect it is likely to be representative present practices.

Expanded model to examine population response to illegal hunting and sport hunting

START	1	Nominal starting population:	1,000
YEAR	20		

AGE	1	2	3	4	5	6	7	8	9	10	11	12	TOTALS
-----	---	---	---	---	---	---	---	---	---	----	----	----	--------

Starting Cohort													
Males	114	76	53	40	32	25	20	16	12	8	3	0	399
Females	114	96	80	68	58	50	43	36	29	18	8	1	601

Running Cohort (population from previous year)													
Males	926	611	430	321	246	167	0	0	0	0	0	0	2,701
Females	927	774	647	553	473	407	351	293	234	144	64	6	4,873
Fecundity	0.00	0.00	0.75	0.95	0.97	0.98	0.98	0.97	0.95	0.50	0.25	0.00	
Calves	0	0	485.3	532.4	458.8	398.9	344.0	284.2	222.3	72.0	16.0	0.0	2,855

Population after births													
Males	1,403	926	611	430	321	246	167	0	0	0	0	0	4,104
Females	1,404	927	774	647	553	473	407	351	293	234	144	64	6,271

Illegal hunting													
Percent of population 2 %													
Expected offtake 208													
Male offtake	28	19	12	9	6	5	3	0	0	0	0	0	82
Female offtake	28	19	15	13	11	9	8	7	6	5	3	1	125
Males	1,375	907	599	421	315	241	164	0	0	0	0	0	4,022
Females	1,376	908	759	634	542	464	399	344	287	229	141	63	6,146

Sport hunting													
Quota 2 %													
Expected offtake 208													
Selectivity %	-	-	-	-	5	10	15	20	20	15	10	5	
Offtake 1					10	21	31	42	42	31	21	10	208
Deficits					0	0	0	-42	-42	-31	-21	-10	-146
Surpluses					305	220	133	0	0	0	0	0	658
Adjustments					512	207	-13	-146	-104	-62	-31	-10	
Corrected					10	34	164						208
Males	1,375	907	599	421	305	207	0	0	0	0	0	0	3,814

Population after annual mortality													
Survival ♂	0.75	0.75	0.80	0.85	0.90	0.90	0.90	0.90	0.90	0.70	0.50	0.10	
Survival ♀	0.75	0.95	0.95	0.97	0.97	0.98	0.98	0.95	0.90	0.70	0.50	0.10	
Males	1,032	681	479	358	275	186	0	0	0	0	0	0	3,011
Females	1,032	863	722	615	526	455	392	327	259	161	71	7	5,430

TOTAL POPULATION	8,441
RATE OF GROWTH %	11.45
ADULT SEX RATIO	3.44

The sport hunting simulation works as follows –

- (i) It is assumed that sport hunting is restricted to males only and no males under the age of 5 years will be taken as trophies;
- (ii) The hunting quota is calculated as a percentage of the total population after births and the number of animals expected to be taken is presented in a cell in the same row;
- (iii) In the next row ('Selectivity') a set of age-specific hunting selectivities are assumed – for any given hunting quota, 5% of the quota will be taken from the 5 year-olds, 10% from the 6 year-olds, 15% from the 7-year olds . . . and so on up to the twelve year-olds;
- (iv) In the next row ('Mortality') the theoretical offtake from each age class is calculated;
- (v) The male population may not be able to provide the trophies in the proportions demanded, so that the shortfalls in each age class are calculated in the next row ('Deficits');
- (vi) In the next row ('Surpluses') the male age classes which will have a surplus of animals after the age-specific hunting quotas have been deducted are identified and the surpluses calculated;
- (vii) In the next row ('Adjustments'), the deficit/surplus is cumulated, beginning with the oldest age class;
- (viii) In the next row ('Corrected'), the number of males to be deducted from each age class is calculated by scanning the + and - signs in the previous row. If there is a negative value, the number inserted in the cell is simply the maximum number of animals for that age class, i.e. all of the animals in the age class concerned will be killed. At the point where the cell entry changes from negative to positive the required deficits are made up from the available animals in the younger age classes.
- (ix) In the last row ('Males'), the remaining population of males is calculated by deducting the number in the row above;
- (x) The model then continues with the normal calculation of age-specific survival and completes its cycle.

In this way, the effect of hunting quotas which exceed the offtake possible from the older age classes can be compensated for by obtaining younger animals – as would the case in real life.

A full set of tests of the sustainable limits of illegal hunting and sport hunting have been done and these are presented in the main body of the report.

Air survey estimates for Roan, Sable and Tsessebe in Namibia

The tables on the next three pages summarise all air surveys (or, in some cases, best estimates when these are not based on air surveys) which have been carried in the areas where roan, sable and tsessebe occur (or have occurred) since 1970. Data for commercial farms (DSS 2003) is not included in the tables but is shown on Figures 9-11 of the main report.

The starting point for the tables is the report by DSS (2002) summarising most of the air surveys done in Namibia. Additional data has been added from –

- DSS (2003) National estimates for Roan, Sable and Tsessebe, 2003
- Erb (1992) Best estimates for roan in Waterberg Plateau Park
- Erb (1993) Improved estimates for sable and tsessebe in Waterberg Plateau Park
- Craig (1998) Additional data for Khaudum, Bushmanland and Caprivi
- Craig (2000) New data for north-eastern Namibia
- Killian (2003, pers. comm.) Recent information on sable and roan in Etosha
- LIFE (2002) Population estimates from the Nyae Nyae Conservancy
- Rodwell, Tagg and Grobler (1995) Survey of the Caprivi in 1994 and summary data of previous Caprivi surveys
- ULG (1994) Survey of the Caprivi in 1994

In cases where more than one estimate has been available for the same area in the same year, the higher of the two estimates has been put in the table. Italics have been used for some numbers in the subtotals for the east and west Caprivi where the survey results for individual areas within the east and west Caprivi add up to a subtotal which is lower than a separate survey result for the east or west Caprivi as a whole. In such a case the higher total has been used.

Year	ETOSHA						WATERBERG		BUSHMANLAND				WESTERN CAPRIVI					EASTERN CAPRIVI					Total	GRAND TOTAL	
	Main Park	West Etosha	Kaross	Khaobendes	Kaross + Khaobendes	Total	Waterberg	Waterberg captures	Bushmanland	Bushmanland + Nyae Nyae	Khaudum	Total	Mahango	Western Core	Central Caprivi	Eastern Core	Subtotal	Mudumu	Mamili	Forest Reserve	Eastern Floodplain	Subtotal			
1970	-	.	.	74	.	74
1971	-	.	.	92	89	92
1972	114	114
1973	-	.	.	.	159	159
1974	-	.	.	.	177	177
1975	-	.	.	.	136	136	70	70
1976	-	.	.	.	204	204	77	7
1977	-	.	.	.	160	160	80	.	3	.	.	3
1978	-	5	.	.	202	207	85	.	27	.	.	27	0	0	319
1979	0	.	.	.	243	243	88	.	16	.	.	16	0	0	347
1980	.	13	.	.	223	236	100	.	55	.	20	75	39	0	33	0	243	243	654	
1981	197	197	176	16	66	66	0	373
1982	19	.	.	.	11	30	220	32	0	9	0	71	71	321	
1983	.	0	.	.	12	12	275	14	0	1	0	15	39	326	
1984	2	0	.	.	21	23	275	.	108	410	518	26	0	1	0	27	27	843	
1985	210	-72	.	.	159	159	4	-	.	.	.	12	0	2	0	14	18	387	
1986	.	0	.	36	.	36	245	43	14	0	0	0	14	57	338	
1987	.	.	40	.	.	40	215	-37	.	.	237	237	61	16	22	99	591	
1988	.	.	40	.	.	40	156	-77	.	56	.	56	63	.	.	4	67	14	0	0	.	14	81	333	
1989	39	39	150	63	0	.	0	63	17	0	0	0	17	80	269	
1990	0	.	20	.	42	42	160	.	.	0	220	220	58	0	.	.	58	0	0	.	.	0	58	480	
1991	94	.	.	.	94	94	94
1992	323	323
1993	.	.	61	.	.	61	158	.	.	7	165	0	0	.	.	0	165	226	
1994	.	.	83	.	.	83	130	6	0	6	142	55	0	0	0	55	197	280	
1995	0	.	110	.	.	110	74	.	124	75	199	107	.	.	.	107	0	0	107	490	
1996	0	0	0
1997	0	0	162
1998	29	.	.	.	80	109	162	.	33	.	33	66	6	28	.	.	34	10	.	.	.	10	44	219	
1999	80	80	0	80
2000	25	25	137	.	.	0	99	99	22	.	.	.	22	22	283	
2001	89
2002	80

• No survey – Species not on list of animals surveyed 0 Species not seen on survey

Year	ETOSHA						WATERBERG		BUSHMANLAND				WESTERN CAPRIVI					EASTERN CAPRIVI					SABLE		
	Main Park	West Etosha	Kaross	Khaobendes	Kaross + Khaobendes	Total	Waterberg	Waterberg captures	Bushmanland	Bushmanland + Nyae Nyae	Khaudum	Total	Mahango	Western Core	Central Caprivi	Eastern Core	Subtotal	Mudumu	Mamili	Forest Reserve	Eastern Floodplain	Subtotal	Total	GRAND TOTAL	
1970	-	-	-
1971	-	-	-
1972	-	-	-
1973	-	-	-
1974	-	-	-
1975	-	.	.	.	0	0	0
1976	-	.	0	.	0	0	0
1977	-	.	.	.	0	0	0
1978	-	-	.	.	26	26	.	25	-	.	-	0	0	26	
1979	-	.	.	.	26	26	.	.	-	.	-	0	0	26	
1980	.	0	.	.	29	29	25	6	-	.	-	55	0	19	8	137	137	191	
1981	29	29	.	29	113	113	0	113	142	
1982	-	.	.	.	0	0	.	-2	47	0	26	0	158	158	158	
1983	.	0	.	.	0	0	37	-2	1	41	0	3	0	44	45	82	
1984	-	0	.	.	0	0	53	-1	.	-	-	31	0	1	0	32	32	85	
1985	61	-3	.	-	-	73	52	.	.	125	33	0	4	0	0	37	162	223	
1986	.	-	.	-	.	-	57	-2	.	.	.	68	.	.	.	68	14	0	0	0	0	14	82	139	
1987	-	.	-	.	.	-	57	-3	.	.	-	103	33	.	.	136	2	138	195	
1988	.	.	-	.	.	-	59	.	.	-	-	148	.	.	64	212	16	0	0	.	.	16	228	287	
1989	0	0	99	0	.	0	99	17	0	0	0	0	17	116	116	
1990	-	.	20	.	24	44	.	.	-	-	-	98	28	.	.	126	20	0	.	.	.	20	146	190	
1991	142	.	.	.	142	142	142	142
1992	989	989
1993	.	.	0	.	.	0	78	48	.	7	133	0	0	.	.	.	0	133	133	
1994	.	.	0	.	.	0	158	187	2	265	612	514	0	0	74	588	1200	1200		
1995	-	.	0	.	.	0	74	.	-	-	-	40	.	.	.	40	0	0	40	114	
1996
1997
1998	-	.	.	.	-	-	157	.	-	-	-	35	21	.	70	126	0	0	128	285	
1999	46	46	46
2000	-	.	.	.	50	50	119	.	-	-	-	130	.	.	.	130	130	299	
2001									-	-	-														
2002									-	-	-														

• No survey - Species not on list of animals surveyed 0 Species not seen on survey

Year	ETOSHA						WATERBERG		BUSHMANLAND				WESTERN CAPRIVI					EASTERN CAPRIVI					Total	GRAND TOTAL			
	Main Park	West Etosha	Kaross	Khaobendes	Kaross + Khaobendes	Total	Waterberg	Waterberg captures	Bushmanland	Bushmanland + Nyae Nyae	Khaudum	Total	Mahango	Western Core	Central Caprivi	Eastern Core	Subtotal	Mudumu	Mamili	Forest Reserve	Eastern Floodplain	Subtotal					
1970	-	-	
1971	-	.	.	.	-	-
1972	0	0
1973	-	.	.	.	1	1
1974	-	.	.	.	0	0
1975	-	.	.	.	0	0
1976	-	.	0	.	0	0
1977	-	.	.	.	0	0	.	.	-	.	-
1978	-	-	.	.	8	8	.	-	.	.	-	0	0	0	0	0	8
1979	-	.	.	.	10	10	.	-	.	.	-	0	0	0	0	0	10
1980	.	-	.	.	13	13	7	7	-	.	0	0	54	0	0	0	0	116	116	116	136	136	
1981	12	12	0	35	35	35	47	
1982	-	.	.	.	1	1	10	3	15	0	0	54	54	54	55	55	
1983	.	-	.	.	1	1	8	.	.	.	33	0	15	0	0	48	56	56	57	57	
1984	-	.	.	.	1	1	1	1	.	-	24	24	31	0	1	0	0	32	32	32	58	58	
1985	16	14	.	.	36	36	13	6	.	.	9	0	3	0	0	12	38	38	90	90	
1986	.	-	.	-	.	-	11	0	.	.	.	8	1	0	0	0	1	9	9	20	20	
1987	-	.	-	.	.	-	13	.	.	-	2	2	11	0	.	.	11	0	11	11	26	26	
1988	.	.	-	.	.	-	15	.	-	.	-	-	22	.	.	16	38	6	0	0	.	13	51	66	66	66	
1989	0	0	19	40	0	.	0	40	0	0	0	0	0	40	40	59	59	
1990	-	.	-	.	0	0	18	.	-	0	0	0	24	0	.	.	24	19	0	.	.	19	43	61	61	61	
1991	35	.	.	.	35	35	35	35	35	35
1992	0	0	0	0
1993	.	.	-	.	.	-	33	0	.	20	53	0	0	.	.	0	53	53	53	53	
1994	.	.	-	.	.	-	17	0	0	61	78	128	0	0	0	128	206	206	206	206	
1995	-	.	-	.	.	-	0	-	.	0	0	0	18	.	.	.	18	0	.	.	.	0	18	18	18	18	
1996
1997
1998	-	.	.	.	-	-	15	.	-	0	0	0	27	-	.	24	51	0	.	.	.	0	51	66	66	66	
1999	0	0	0	0
2000	-	.	.	.	-	-	2	.	-	0	0	0	0	.	.	.	0	0	2	2	2	2
2001
2002

• No survey - Species not on list of animals surveyed 0 Species not seen on survey

Rainfall Data for Selected Sites

In order to examine the effects of the accumulated surplus and deficit in rainfall and the late dry season rainfall on the roan, sable and tsessebe populations in northern Namibia, long term data were needed for several different areas – Etosha, Waterberg, Khaudum, East Caprivi and West Caprivi. These data were very kindly provided by John Mendelsohn of the Department of Environmental Affairs. The relevant sites for which rainfall data were available are –

Data used

Etosha	– Kamanjab, Ermo, Uries Ekango, Otjitambi and Ondjou	Kamanjab
Waterberg	– Otjiwarongo, Okakarara, Hohenfels, Okosongomingo, Etekero	Hohenfels
Khaudum and West Caprivi	– Andara	Andara
East Caprivi	– Katima Mulilo and Sesheke	Katima Mulilo

Where multiple sites were available, the data were plotted and examined. In all cases the differences between nearby rainfall stations in any particular locality were not great and so one station was selected for each area – mainly on the basis of the completeness and length of the rainfall record. The four rainfall sites selected had records going back to 1960-61⁴⁶ and these are shown on the pages which follow.

The seasonal rainfall for any given year was obtained by totalling the months July-December for the previous year and adding this to the rainfall from January to June for the given year. The mean annual rainfall was derived from the average of all the seasonal rainfall figures. The accumulated rainfall surplus/deficit for each year was calculated by subtracting the mean annual rainfall from the actual rainfall for each year and cumulating the resulting positive and negative values, beginning with the first year in the data sequence.

On examining the cumulative surplus/deficit data it was apparent that the data did not have a zero mean.⁴⁷ Dunham and Robertson (2001, 2003), in using the cumulative surplus/deficit in rainfall for modelling populations, simply began their data sequence with whatever the data value happened to be in the first year of their analysis. In this case I have forced the data sequence to have a zero mean by subtracting a fixed amount from each point on the curve such that the curve has equal areas above and below zero. In a sense, this gives an absolute value to the concept of deficit and surplus: the shape of the curve is unaltered but the total of the deficit area is equal to the total of the surplus area – lending credence to the idea of a constant long term mean. The calculations of the data used in **Figs. 12-19** are given in the tables which follow, together with the the dry season rainfall.

46. Unfortunately, the data for the most recent years (1997 onwards) were not available. Contact has been made with the Namibian Weather Bureau and the data has been promised. However it has not yet been received.

47. This is entirely to be expected. If one considers a sine wave where exactly one wave length is contained within the start and end of the data span (i.e. the first half cycle of the sine wave has positive values reaching a peak at a quarter wave length and the second half cycle has negative values reaching a trough at three-quarters of the wave length), then the integral of this curve is entirely positive. The area under the curve reaches a peak at the half-wavelength point and then decreases to zero as the negative values are subtracted in the second half of the cycle.

YEAR	Monthly rainfall (mm)												TOTAL	Seasonal Total	DEFICIT/SURPLUS		Dry Season Rainfall	
	Jan	Feb	Mar	Apr	May	June	July	Aug	Sept	Oct	Nov	Dec			1961 →	Zero Mean	Jul-Oct	Nov
1961	20.0	50.5	68.0	23.0	26.0	0.0	0.0	0.0	0.0	0.0	66.0	9.5	263.0					
1962	18.5	43.5	13.0	67.5	0.0	0.0	0.0	16.5	0.0	1.0	31.0	14.5	205.5	218	-70	-407		
1963	179.5	49.0	177.5	79.0	0.0	0.0	0.0	0.0	6.0	0.0	70.5	7.5	569.0	548	189	-148		
1964	8.5	101.4	13.7	33.8	0.0	0.0	0.0	0.0	0.0	9.0	5.5	8.2	180.1	241.4	142	-195		
1965	72.5	96.0	47.5	29.5	0.0	0.0	0.0	0.0	20.5	0.0	12.5	0.0	278.5	268.2	122	-215		
1966	157.0	17.0	304.7	24.5	0.0	0.0	0.0	0.0	5.0	16.0	0.0	19.0	543.2	536.2	370	33		
1967	60.0	86.5	122.0	0.0	0.0	0.0	0.0	0.0	0.0	14.0	87.0	98.5	468.0	308.5	390	53		
1968	21.0	40.5	80.2	7.5	0.0	0.0	0.0	0.0	0.0	0.0	39.5	24.5	213.2	348.7	450	113		
1969	47.0	112.5	88.5	5.0	0.0	0.0	0.0	0.0	0.0	4.0	10.5	15.0	282.5	317	479	142		
1970	89.5	49.5	48.5	1.5	0.0	0.0	0.0	0.0	0.0	11.0	5.5	0.0	205.5	218.5	409	72	11	6
1971	78.5	271.0	37.5	4.0	3.0	7.0	0.0	0.0	0.0	0.0	0.0	45.0	446.0	417.5	538	201	0	0
1972	112.0	9.5	124.0	0.0	0.0	0.0	0.0	0.0	0.0	11.0	0.0	0.0	256.5	290.5	540	203	11	0
1973	8.5	0.0	133.5	65.0	0.0	0.0	0.0	0.0	0.0	42.0	10.5	0.0	259.5	218	469	132	42	11
1974	170.5	189.5	40.0	0.0	0.0	0.0	0.0	0.0	0.0	94.0	21.0	0.0	515.0	452.5	634	296	94	21
1975	44.5	105.5	163.5	13.0	0.0	0.0	0.0	0.0	0.0	6.0	23.0	0.0	355.5	441.5	787	450	6	23
1976	148.5	86.5	108.0	34.5	0.0	1.0	0.0	0.0	0.0	0.0	0.0	8.0	386.5	407.5	906	569	0	0
1977	53.0	43.5	75.5	37.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	24.0	233.9	217.9	835	498	0	0
1978	74.7	51.8	55.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	21.6	0.0	203.8	206.2	753	416	0	22
1979	123.0	105.2	20.0	0.0	5.0	0.0	0.0	0.0	2.5	6.0	47.5	0.0	309.2	274.8	739	402	8.5	48
1980	0.0	44.0	94.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	21.0	7.0	166.4	194.4	645	308	0	21
1981	10.5	0.0	33.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	43.5	71.5	428	91	0	0
1982	25.5	93.0	80.5	0.0	0.0	0.0	0.0	0.0	0.0	6.5	5.0	62.5	273.0	199	339	2	6.5	5
1983	49.0	24.6	24.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	28.5	126.6	172.1	223	-114	0	0
1984	49.3	37.5	96.5	36.0	2.5	30.0	0.0	0.0	0.0	7.0	25.5	0.0	284.3	280.3	215	-123	7	26
1985	97.0	63.0	61.0	0.0	13.5	0.0	0.0	0.0	0.0	15.5	35.0	0.0	285.0	267	193	-144	15.5	35
1986	121.5	37.5	58.0	46.0	0.0	0.0	0.0	0.0	2.0	2.0	12.5	22.0	301.5	313.5	218	-119	4	13
1987	3.0	95.0	5.0	32.5	0.0	0.0	0.0	0.0	0.0	49.0	29.5	9.0	223.0	174	104	-233	49	30
1988	114.0	60.5	30.0	5.0	0.0	0.0	0.0	0.0	0.0	0.0	42.5	90.5	342.5	297	112	-225	0	43
1989	13.0	43.0	10.0	40.5	0.0	0.0	0.0	0.0	0.0	0.0	4.0	5.0	115.5	239.5	63	-274	0	4
1990	91.0	44.0	138.0	0.0	0.0	0.0	0.0	0.0	0.0	5.5	8.0	24.5	311.0	282	57	-280	5.5	8
1991	129.5	104.0	102.0	0.0	0.0	0.0	0.0	0.0	13.5	19.0	58.0	58.5	484.5	373.5	142	-195	32.5	58
1992	84.5	8.0	4.0	5.5	0.0	0.0	0.0	0.0	31.0	2.0	12.0	0.0	147.0	251	105	-232	33	12
1993	76.0	45.0	123.0	10.0	0.0	0.0	0.0	0.0	0.0	17.5	6.5	66.5	344.5	299	115	-222	17.5	7
1994	10.5	48.5	6.0	9.5	0.0	0.0	0.0	0.0	0.0	1.0	17.0	8.0	100.5	165	-8	-345	1	17
1995	0.0	298.5	109.0	28.0	0.0	0.0	0.0	0.0	0.0	0.0	25.5	1.0	462.0	461.5	165	-172	0	26
1996	73.0	24.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	19.0	116.0	123.5	-0	-337	0	0

KAMANJAB

STATISTICS	
Total	10094.70
Years	35
Average	288.42
DEFICIT /SURPLUS	
Offset for zero mean	
337.06	

YEAR	Monthly rainfall (mm)												TOTAL	Seasonal Total	DEFICIT/SURPLUS		Dry Season Rainfall		HOHENFELS
	Jan	Feb	Mar	Apr	May	June	July	Aug	Sept	Oct	Nov	Dec			1961 →	Zero Mean	Jul-Oct	Nov	
1961	82.3	61.4	112.0	69.3	15.5	0.3	0.0	0.0	0.0	1.2	32.5	11.0	385.5						
1962	41.5	60.4	14.0	31.8	0.0	0.0	0.0	27.0	0.0	43.0	13.5	4.0	235.2	192.4	-241	-655			
1963	284.0	0.0	170.0	40.0	0.0	0.0	0.0	0.0	0.0	0.0	174.0	49.0	717.0	581.5	-93	-507			
1964	40.0	64.0	34.0	0.0	2.5	0.0	0.0	0.0	0.0	0.0	30.0	31.5	202.0	363.5	-164	-577			
1965	90.5	65.0	97.5	111.0	0.0	0.0	0.0	0.0	0.0	0.0	48.0	12.5	424.5	425.5	-172	-585			
1966	316.5	56.8	45.5	46.0	0.0	0.0	0.0	0.0	30.8	0.0	0.0	57.0	552.6	525.3	-80	-494			
1967	159.0	198.0	74.0	16.0	57.0	0.0	0.0	0.0	0.0	0.0	166.7	102.5	773.2	591.8	78	-336			
1968	86.0	48.0	184.7	4.9	22.0	0.0	0.0	0.0	0.0	0.0	93.0	61.0	499.6	614.8	259	-154			
1969	75.0	141.0	101.5	53.0	0.0	0.0	0.0	0.0	0.0	0.0	23.0	41.5	435.0	524.5	350	-64			
1970	120.0	23.0	13.2	0.0	0.0	0.0	0.0	0.0	0.0	47.0	6.5	142.0	351.7	220.7	137	-277			
1971	79.5	265.0	27.0	45.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	4.0	420.5	612	315	-98			
1972	117.0	4.0	137.0	44.5	0.0	0.0	0.0	0.0	0.0	6.0	17.0	41.5	367.0	306.5	188	-226			
1973	14.0	0.0	214.0	46.5	0.0	0.0	0.0	0.0	0.0	24.0	56.5	63.5	418.5	339	93	-320			
1974	319.0	188.0	37.0	10.0	0.0	3.0	0.0	0.0	0.0	36.0	43.0	21.0	657.0	701	361	-53			
1975	107.0	64.0	137.0	64.0	0.0	0.0	0.0	0.0	0.0	12.0	59.0	23.0	466.0	472	399	-15	12	59	
1976	233.0	204.0	110.5	33.5	0.0	7.0	0.0	0.0	0.0	6.0	31.0	35.0	660.0	682	647	234	6	31	
1977	147.0	200.5	59.0	62.0	9.0	0.0	0.0	0.0	0.0	83.0	11.0	74.5	646.0	549.5	763	349	83	11	
1978	246.0	164.5	60.5	57.0	5.0	0.0	0.0	3.5	0.0	9.0	25.5	69.0	640.0	701.5	1031	617	13	26	
1979	103.5	322.0	3.0	15.0	3.0	0.0	0.0	0.0	2.0	18.0	86.5	33.5	586.5	553.5	1151	737	20	87	
1980	60.0	104.0	112.0	4.0	0.0	0.0	0.0	0.0	30.0	0.0	13.0	74.0	397.0	420	1137	723	30	13	
1981	53.0	68.0	18.0	4.0	0.0	0.0	0.0	6.0	0.0	0.0	16.5	0.0	165.5	260	963	550	6	17	
1982	66.5	140.5	62.0	32.0	0.0	0.0	0.0	0.0	2.0	15.0	71.0	53.0	442.0	323.5	853	440	17	71	
1983	74.5	21.5	32.0	9.0	7.5	34.0	0.0	0.0	0.0	31.0	49.0	162.5	421.0	319.5	739	325	31	49	
1984	48.0	60.0	77.5	137.5	0.0	0.0	0.0	0.0	0.0	9.5	7.0	7.0	346.5	565.5	871	457	10	7	
1985	160.0	109.2	25.0	18.7	0.0	0.0	0.0	0.0	0.0	3.0	16.0	24.5	356.4	336.4	773	360	3	16	
1986	84.5	226.0	53.5	0.0	0.0	0.0	0.0	0.0	0.0	8.5	4.0	24.0	400.5	407.5	747	334	9	4	
1987	14.0	125.0	38.0	38.0	0.0	0.0	0.0	0.0	0.0	38.0	52.0	14.0	319.0	251.5	565	151	38	52	
1988	98.0	57.0	0.0	183.0	0.0	0.0	0.0	0.0	0.0	9.0	66.0	82.0	495.0	442	573	160	9	66	
1989	101.0	167.0	6.0	41.0	3.0	0.0	0.0	0.0	0.0	3.0	1.0	5.0	327.0	475	615	201	3	1	
1990	114.0	49.0	112.0	37.0	0.0	0.0	0.0	0.0	0.0	4.0	26.0	70.0	412.0	321	502	88	4	26	
1991	113.0	211.0	0.0	0.0	0.0	0.0	0.0	0.0	12.0	23.0	55.0	106.0	520.0	424	492	79	35	55	
1992	34.0	3.0	64.0	3.0	0.0	0.0	0.0	0.0	1.0	0.0	0.0	14.0	119.0	300	358	-55	1	0	
1993	85.0	180.0	43.0	46.0	0.0	0.0	0.0	2.0	0.0	42.0	58.0	34.0	490.0	369	294	-120	44	58	
1994	276.0	44.0	16.0	4.0	0.0	0.0	0.0	0.0	0.0	0.0	9.0	5.0	354.0	476	336	-77	0	9	
1995	18.0	118.0	74.0	5.0	0.0	0.0	0.0	0.0	4.0	10.0	29.0	12.0	270.0	229	131	-282	14	29	
1996	98.0	35.5	12.5	16.5	0.0	0.0	0.0	0.0	0.0	4.0	0.0	63.0	229.5	217.5	-85	-498	4	0	
1997	229.0	94.0	108.5	20.0	0.0	0.0	0.0	0.0	17.0	58.0	15.0	122.0	663.5	518.5	0	-413	75	15	

STATISTICS	
Total	15612.9
Years	36
Average	433.7
DEFICIT /SURPLUS Offset for zero mean	
	413.45

YEAR	Monthly rainfall (mm)												TOTAL	Seasonal Total	DEFICIT/SURPLUS		Dry Season Rainfall		ANDARA
	Jan	Feb	Mar	Apr	May	June	July	Aug	Sept	Oct	Nov	Dec			1961 →	Zero Mean	Jul-Oct	Nov	
1961	71.7	97.4	171.6	47.5	9.5	0.0	0.0	0.0	0.0	0.0	43.9	66.5	508.1						
1962	154.2	161.7	11.3	37.2	0.0	0.0	0.0	3.0	0.0	0.0	77.3	130.1	574.8	475.0	-104.0	-109.0			
1963	137.6	66.0	67.3	3.5	0.0	0.0	0.0	0.0	1.1	40.0	137.6	72.0	525.1	485.0	-198.0	-203.0			
1964	24.0	23.5	31.5	0.0	0.0	0.0	0.0	0.0	0.0	6.2	20.0	108.7	213.9	330.0	-447.0	-453.0			
1965	57.5	67.0	15.0	37.5	0.0	0.0	0.0	0.0	0.0	9.5	34.3	46.5	267.3	312.0	-714.0	-720.0			
1966	160.5	257.6	175.5	25.6	0.0	6.0	0.0	0.2	16.5	0.0	29.5	63.0	734.4	716.0	-578.0	-583.0			
1967	193.5	89.0	99.5	61.3	0.0	0.0	0.0	0.0	0.0	19.0	223.0	69.0	754.3	553.0	-604.0	-609.0			
1968	171.5	78.5	79.0	11.5	0.0	0.0	0.0	0.0	1.5	0.0	38.0	53.5	433.5	652.0	-531.0	-537.0			
1969	108.5	173.5	78.5	6.1	0.0	0.0	0.0	0.0	4.0	24.0	63.0	72.0	529.6	460.0	-650.0	-656.0			
1970	58.5	125.5	21.0	0.0	0.0	0.0	0.0	0.0	0.0	2.0	58.0	115.5	380.5	368.0	-861.0	-867.0	2	58	
1971	231.7	152.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	12.0	98.0	78.6	572.8	560.0	-880.0	-886.0	12	98	
1972	294.5	54.6	365.0	14.5	0.0	0.0	0.0	0.0	0.1	10.0	17.5	19.9	776.1	917.0	-542.0	-547.0	10	18	
1973	88.0	213.5	84.5	7.6	0.0	0.0	0.0	0.0	0.0	39.8	49.5	296.0	778.9	441.0	-680.0	-685.0	40	50	
1974	273.9	274.8	19.5	132.5	5.0	0.0	0.0	0.0	10.0	1.1	51.6	27.7	796.1	1091.0	-168.0	-173.0	11	52	
1975	147.0	141.0	188.5	11.0	7.0	0.0	0.0	0.0	0.0	0.0	22.0	82.7	599.2	585.0	-162.0	-167.0	0	22	
1976	180.0	133.0	131.0	34.5	9.0	0.0	0.0	0.0	35.0	29.0	77.5	35.9	664.9	592.0	-148.0	-154.0	64	78	
1977	58.3	321.0	180.2	20.5	14.0	0.0	0.0	0.0	16.5	14.0	56.3	262.5	943.3	771.0	44.0	39.0	31	56	
1978	201.5	448.5	190.9	14.0	0.0	0.0	0.0	0.0	0.0	22.6	58.0	77.0	1012.5	1204.0	670.0	664.0	23	58	
1979	96.5	91.2	41.5	2.5	0.0	0.0	0.0	5.0	3.0	3.0	116.5	65.5	424.7	389.0	480.0	475.0	11	117	
1980	75.0	150.5	51.6	15.0	0.0	0.0	0.0	0.0	0.0	14.0	247.6	77.0	630.7	485.0	386.0	381.0	14	248	
1981	112.5	306.5	154.5	0.0	0.0	2.0	0.0	0.0	0.0	0.0	121.3	23.0	719.8	914.0	722.0	716.0	0	121	
1982	57.4	134.9	109.8	69.7	0.0	0.0	0.0	0.0	0.0	4.5	4.5	7.5	388.3	516.0	659.0	653.0	5	5	
1983	—	—	—	0.0	0.0	0.0	0.0	0.0	0.0	39.7	17.8	160.3	—	579.0	659.0	654.0	40	18	
1984	21.8	118.3	151.2	31.5	0.0	0.0	0.0	1.0	4.9	82.5	21.4	44.2	476.8	541.0	621.0	615.0	88	21	
1985	197.6	181.0	30.7	0.0	0.0	0.0	0.0	0.0	0.0	23.3	1.5	151.3	585.4	563.0	605.0	600.0	23	2	
1986	94.5	112.6	112.3	38.4	0.0	0.0	0.0	0.0	17.6	0.0	103.7	88.8	567.9	534.0	560.0	555.0	18	104	
1987	61.6	141.0	35.4	0.0	21.4	0.0	0.0	0.0	9.0	29.9	31.1	128.4	457.8	470.0	451.0	446.0	39	31	
1988	50.2	143.7	72.6	90.9	0.0	0.0	0.0	0.0	0.1	4.4	66.4	148.0	576.3	556.0	428.0	422.0	5	66	
1989	—	—	—	—	0.0	0.0	0.0	0.0	0.0	4.7	51.4	25.6	—	579.0	428.0	423.0	5	51	
1990	113.5	128.6	67.2	66.0	0.1	0.0	0.0	0.0	0.0	8.0	19.7	127.6	530.7	457.0	306.0	301.0	8	20	
1991	100.2	107.3	76.3	0.0	0.0	0.0	0.0	0.0	0.0	58.5	64.8	244.7	651.8	439.0	167.0	161.0	59	65	
1992	84.7	76.7	66.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	17.5	36.1	281.4	596.0	184.0	178.0	0	18	
1993	249.5	101.0	35.5	38.0	0.0	0.0	0.0	0.0	0.0	1.0	8.5	67.3	500.8	478.0	82.0	77.0	1	9	
1994	263.5	131.6	20.0	4.5	0.0	0.0	0.0	0.0	0.0	10.0	36.0	60.5	526.1	496.0	0.0	-6.0	10	36	
1995	3.5	54.0	41.0	41.5	0.0	0.0	0.0	0.0	0.0	—	—	—	—	579.0	0.0	-5.0	0		

Mean rainfall used where data is missing

STATISTICS	
Total	19680.7
Years	34
Average	578.8
DEFICIT /SURPLUS Offset for zero mean	
	5.41

YEAR	Monthly rainfall (mm)												TOTAL	Seasonal Total	DEFICIT/SURPLUS		Dry Season Rainfall		KATIMA MULILO
	Jan	Feb	Mar	Apr	May	June	July	Aug	Sept	Oct	Nov	Dec			1960 →	Zero Mean	Jul-Oct	Nov	
1960	89.4	178.3	121.3	24.8	10.0	0.0	0.0	0.0	2.0	4.0	125.5	41.8	597.1						
1961	154.3	58.9	235.5	41.8	2.5	0.0	0.0	0.0	0.0	8.8	72.4	68.8	643.0	666	39	-284			
1962	242.4	104.7	15.4	28.7	0.0	0.0	0.0	0.0	0.0	1.4	93.0	286.7	772.3	541	-48	-371			
1963	195.8	223.2	44.7	0.0	0.0	7.5	0.0	0.0	0.0	58.4	86.7	197.3	813.6	852	177	-146			
1964	87.2	54.4	1.5	0.0	0.0	0.0	0.0	0.0	7.5	18.7	42.5	51.2	263.0	486	35	-289			
1965	75.6	50.5	9.2	14.0	0.0	0.0	0.0	0.0	0.4	0.7	73.2	72.5	296.1	269	-324	-647			
1966	68.5	185.4	140.0	16.5	0.0	5.4	0.0	6.0	0.0	12.0	34.5	156.9	625.2	563	-389	-712			
1967	239.6	79.0	68.0	10.5	0.0	0.0	0.0	0.0	0.0	124.3	109.9	156.9	788.2	607	-410	-733			
1968	239.8	86.8	50.8	50.2	0.7	0.0	0.0	0.0	1.8	14.5	38.8	95.2	578.6	819	-219	-542			
1969	91.1	142.8	195.5	1.5	0.0	0.0	0.0	0.0	0.0	26.2	82.2	168.8	708.1	581	-265	-588			
1970	120.1	54.5	14.0	6.6	0.0	0.0	0.0	0.0	0.5	5.4	123.9	136.6	461.6	472	-420	-743	6	124	
1971	268.7	54.4	13.0	3.5	0.0	0.0	0.0	0.0	0.0	8.8	98.0	125.5	571.9	606	-442	-765	9	98	
1972	411.3	77.3	226.4	57.5	0.0	0.0	0.0	0.0	0.0	9.2	27.4	80.0	889.1	1005	-65	-388	9	27	
1973	64.9	166.2	40.1	7.9	0.0	0.0	0.0	0.0	0.0	55.0	41.7	236.3	612.1	396	-297	-620	55	42	
1974	206.1	465.6	101.7	13.2	0.0	0.0	0.0	0.0	7.5	28.0	120.0	319.0	1261.1	1120	195	-128	36	120	
1975	99.2	90.7	135.7	16.5	8.9	0.0	0.0	0.4	0.0	0.0	17.8	92.4	461.6	826	393	70	0	18	
1976	191.0	62.9	122.2	15.9	7.0	0.0	0.0	0.0	27.1	21.5	150.0	151.4	749.0	510	275	-48	49	150	
1977	97.7	379.0	236.4	38.8	0.6	0.0	0.0	0.4	6.1	0.0	68.5	302.2	1129.7	1103	749	426	7	69	
1978	231.7	418.3	111.7	59.7	0.0	0.0	0.0	0.0	0.0	68.5	72.2	75.6	1037.7	1199	1320	997	69	72	
1979	135.8	145.9	44.5	10.2	5.0	0.0	0.0	2.7	0.0	19.3	72.6	54.8	490.8	558	1250	927	22	73	
1980	99.6	159.7	164.0	12.4	0.0	0.0	0.0	0.8	9.1	14.0	84.8	92.4	636.8	585	1208	885	24	85	
1981	227.9	189.7	66.7	11.7	0.0	3.7	0.0	0.0	4.3	32.4	75.6	140.4	752.4	701	1281	958	37	76	
1982	94.9	131.8	5.2	34.6	1.1	0.0	0.0	0.0	0.0	39.4	47.2	43.5	397.7	520	1174	850	39	47	
1983	122.2	51.5	35.1	56.4	5.1	0.0	0.0	0.0	0.0	25.6	76.6	150.3	522.8	400	946	623	26	77	
1984	77.4	93.0	134.9	8.0	0.8	0.0	0.4	0.1	7.2	10.5	90.2	51.3	473.8	567	885	562	18	90	
1985	167.8	72.3	62.5	12.3	0.0	0.0	0.0	0.0	0.0	43.5	66.1	197.1	621.6	475	732	409	44	66	
1986	216.0	63.8	106.6	17.3	0.0	0.0	0.0	0.0	2.1	52.2	82.1	91.9	632.0	710	815	492	54	82	
1987	30.0	112.8	37.0	3.4	0.0	0.0	0.0	0.0	0.0	0.0	51.7	198.8	433.7	412	599	275	0	52	
1988	30.8	221.7	60.9	16.5	0.0	0.0	0.0	0.0	18.7	23.9	36.2	184.3	593.0	580	551	228	43	36	
1989	249.6	151.9	43.2	2.6	0.0	0.0	0.0	0.0	0.0	56.0	21.5	98.6	623.4	710	634	311	56	22	
1990	193.2	43.9	46.0	52.8	0.0	0.0	0.0	0.0	0.0	13.0	17.5	72.7	439.1	512	518	195	13	18	
1991	72.9	157.4	141.6	0.0	0.0	0.0	0.0	0.0	0.0	6.0	86.6	281.1	745.6	475	366	43	6	87	
1992	36.6	3.3	108.2	2.6	0.0	0.0	0.0	0.0	0.0	44.9	49.8	135.4	380.8	524	262	-61	45	50	
1993	158.7	165.6	31.2	36.0	0.0	0.0	0.0	0.0	1.7	0.5	148.6	70.1	612.4	622	256	-67	2	149	
1994	101.8	125.2	25.6	0.0	0.0	0.0	0.0	0.0	0.0	5.4	34.5	154.5	447.0	474	102	-221	5	35	
1995	146.0	175.4	61.7	7.8	12.1	0.0	0.0	0.0	0.0	9.4	60.5	187.9	660.8	597	72	-251	9	61	
1996	139.4	117.4	41.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	109.1	95.2	503.0	557	1	-323	0	109	
1997	173.5	147.6	99.9	1.8	0.0	0.0	0.0	0.0	11.8	56.1	27.8	91.5	610.0	627	-0	-323	68	28	

STATISTICS	
Total	23224.7
Years	37
Average	627.7
DEFICIT /SURPLUS Offset for zero mean	
	323.1

Financial Analysis of Sport Hunting Potential for Roan, Sable and Tsessebe

This Appendix consists of seven tables all of which are linked as spreadsheets. The analysis of sport hunting potential is carried out as follows –

In **Table A** a hypothetical wildlife population which might be typical of an area of 1,000km² the Caprivi is set up in two stages. Firstly, the various species are assigned nominal densities which are used to calculate the total metabolic biomass in Livestock Unit equivalents (LSUs). The densities of the grazers are then adjusted by a multiplier to make the overall stocking rate exactly equal to 10ha/LSU – which has arbitrarily been chosen as the carrying capacity for such an area. In this first table there are no roan, sable or tsessebe.

In **Table B**, roan, sable and tsessebe are included at densities which they might achieve in an area of suitable habitat with average annual rainfall above 500mm. In order to preserve the carrying capacity of 10ha/LSU, the densities of the other grazing species are reduced to accommodate the newcomers (again by using a simple multiplying factor).

Hunting quotas for the two scenarios are calculated and transferred to **Table C** where the species are organised into groups based on the value of the trophy fees.

In **Tables D and E** the quotas for the two scenarios (with and without roan, sable and tsessebe) are automatically packaged into a set of hunts and the total number of hunter-days which this generates is calculated together with the gross income from daily rates and trophy fees.

In **Table F** the operating costs for a single safari operator hunting an area of 250km² are calculated. The reason that an area of 250km² has been chosen is that the number of hunter days generated in Table E are sufficiently large to justify 4 separate safari operators in an area of 1,000km².

Table G is the cost and land use summary. It is assumed that roan, sable and tsessebe numbers on which the calculation is based could be achieved over a core area in the Caprivi of about 4,000km². The quota generated by the scenario without roan sable and tsessebe is dominated by ‘low-value’ species which cannot easily be packaged into hunts which would attract international clients and, to maximise the use of the quota, most of the animals would have to be taken by ‘biltong hunters’ within the southern African region. The number of international hunting client days is sufficient to warrant only one safari operator in 1,000km² and it is assumed in the final table that the operating costs for a single safari operator in the larger area would remain much the same as those calculated for an operator in 250km² (mileage costs would increase but the remainder of the costs are linked more to the number of hunter days).

In the scenario which includes roan, sable and tsessebe, the number of international hunting client days increases to over 700 which would justify the allocation of four separate hunting concessions in 1,000km². This causes the overall operating costs to increase and, despite the fact that the gross income per hectare doubles when roan, sable and tsessebe are part of the hunting quota, the net income is only increased by about 20% over the scenario without these key species.

Financial Analysis of Sport Hunting Potential for Roan, Sable and Tsessebe

Table A: Optimal stocking rates and hunting quotas for an area of 1,000km² in the Caprivi assuming no roan, sable and tsessebe present

Species	Factor 1.99071 Density /km ²	Initial population	Model population	Unit LSU	Total LSUs	QUOTA	
						%	N
Buffalo	0.25	250	250	1.00	250.0	2.5	6
Bushbuck	0.05	50	100	0.12	11.9	3	3
Duiker	1.00	1,000	1,991	0.08	159.3	3	60
Eland	0.25	250	498	1.00	497.7	2	10
Elephant	1.00	1,000	1,000	3.33	3330.0	0.5	5
Giraffe	0.05	50	50	1.34	67.0	5	3
Impala	10.00	10,000	19,907	0.14	2787.0	3	597
Kudu	3.00	3,000	3,000	0.40	1200.0	2	60
Leopard	0.05	50	50	–	0.0	6	3
Lion	0.03	30	30	–	0.0	6	2
Reedbuck	0.10	100	199	0.14	27.9	3	6
Steenbok	2.00	2,000	3,981	0.10	398.1	3	119
Waterbuck	0.25	250	498	0.45	224.0	2	10
Warthog	2.00	2,000	3,981	0.18	716.7	3	119
Wildebeest (Blue)	0.10	100	199	0.40	79.6	3	6
Zebra (Burchell's)	0.20	200	398	0.63	250.8	5	20
TOTAL LSU					10000.0	1029	
Ha/LSU					10.0		

NOTES

1. The aim of this table is to create a 'model population' of large mammal species which would be typical for a well-stocked savanna system in an area of 1,000km² where rainfall is 500-600mm per annum and the habitats are suitable for roan, sable and tsessebe (although these species are not included in this first table). On this basis, lechwe, sitatunga and hippo have been excluded. The population is that which might be expected in the central area of the Caprivi around the Kwando River and relates to the 'core' wildlife range.
2. The assigned densities are based on crude potential for the Caprivi and experience from similar savannas.
3. Unit livestock biomass values are the same as those used by Barnes and de Jager (1995)
4. The rows which have been shaded are those species likely to be affected by any inter-specific competition with roan, sable and tsessebe.
5. The 'Model Population' is obtained by multiplying the initial population numbers in the shaded rows upwards by the factor in the top left hand corner of the table which has been selected so that the total stocking density is 1LSU/10ha.
6. The quota percentages are typical for safari hunting in southern Africa being adjusted upwards when trophy quality is less critical (e.g Zebra - 5%) and downwards where high trophy quality is important (e.g. elephant 0.5%).

The numbers of the grazing species have to be more or less double those assumed at the start of this exercise in order to achieve an overall stocking density of 1LSU/10ha.

Table B: Optimal stocking rates and hunting quotas for an area of 1,000km² in the Caprivi assuming roan, sable and tsessebe are present

Species	Factor 1.327 Density /km ²	Initial population	Model population	Unit LSU	Total LSUs	QUOTA	
						%	N
Buffalo	0.25	250	250	1.00	250.0	2.5	6
Bushbuck	0.05	50	50	0.12	6.0	3	2
Duiker	1.00	1,000	1,327	0.08	106.2	3	40
Eland	0.25	250	332	1.00	331.8	2	7
Elephant	1.00	1,000	1,000	3.33	3330.0	0.5	5
Giraffe	0.05	50	50	1.34	67.0	5	3
Impala	10.00	10,000	13,270	0.14	1857.8	3	398
Kudu	3.00	3,000	3,000	0.40	1200.0	2	60
Leopard	0.05	50	50	–	0.0	6	3
Lion	0.03	30	30	–	0.0	6	2
Reedbuck	0.10	100	133	0.14	18.6	3	4
ROAN	1.00	1,000	1,000	0.65	650.0	2	20
SABLE	2.00	2,000	2,000	0.40	800.0	2	40
TSESSEBE	1.00	1,000	1,000	0.27	270.0	2	20
Steenbok	2.00	2,000	2,654	0.10	265.4	3	80
Waterbuck	0.25	250	332	0.45	149.3	2	7
Warthog	2.00	2,000	2,654	0.18	477.7	3	80
Wildebeest (Blue)	0.10	100	133	0.40	53.1	3	4
Zebra (Burchell's)	0.20	200	265	0.63	167.2	5	13
TOTAL LSU					10000.0		794
Ha/LSU					10.0		

NOTES

1. In this table the 'model population' includes roan, sable and tsessebe populations and the numbers of other grazing animals have been reduced to allow for the new populations.
2. The assigned densities for roan, sable and tsessebe are based on crude potential for the Caprivi and experience from similar savannas.
3. The 'Model Population' is obtained by multiplying the initial population numbers in the shaded rows by the factor in the top left hand corner of the table which has been selected so that the total stocking density is 1LSU/10ha. The roan, sable and tsessebe numbers are assumed constant.
4. The quota percentages for roan, sable and tsessebe are based on the findings of the population model (Appendix 1) and for the other species they remain the same as those in Table A.

In order to accommodate the roan, sable and tsessebe populations at the assumed densities, the numbers of other grazers given in Table A are reduced by about one-third in order to maintain a stocking level of 1 LSU/10ha.

Table C: Trophy fees and quota value

Species	Trophy Fee	SCENARIO A			SCENARIO B		
		N	Quota value	#	N	Quota value	#
Premier Species							
ELEPHANT	7,500	5	37,500	5	5	37,500	5
BUFFALO	5,000	6	30,000	6	6	30,000	6
LION	4,000	2	8,000	2	2	8,000	2
LEOPARD	3,000	3	9,000	3	3	9,000	3
Plains Game - A				0			60
Roan	2,000	–	0		20	40,000	
Sable	2,000	–	0		40	80,000	
Plains Game - B				23			37
Eland	1,500	10	15,000		7	10,500	
Giraffe	1,500	3	4,500		3	4,500	
Tsessebe	1,500	–	0		20	30,000	
Waterbuck	1,500	10	15,000		7	10,500	
Plains Game - C				95			83
Bushbuck	750	3	2,250		2	1,500	
Kudu	750	60	45,000		60	45,000	
Reedbuck	750	6	4,500		4	3,000	
Wildebeest (Blue)	750	6	4,500		4	3,000	
Zebra (Burchell's)	750	20	15,000		13	9,750	
Plains Game - D				895			598
Impala	375	597	223,875		398	149,250	
Duiker	375	60	22,500		40	15,000	
Steenbok	375	119	44,625		80	30,000	
Warthog	375	119	44,625		80	30,000	
TOTAL TROPHY FEE VALUE US\$			525,875			546,500	

NOTES

1. Trophy fee values are based on Himavundu (2001) but have been adjusted in some cases to be more closely aligned with the regional averages.
2. Roan, sable and tsessebe are absent from **Scenario A** and included in **Scenario B**. The quotas for the two scenarios are taken from **Table A** and **Table B** respectively.
3. For simplicity in packaging the hunts (see following tables), the trophy fees have been averaged and rounded over groups of animals.

The total 'book' value of the quotas from the two scenarios is not greatly different. However, because the majority of hunts under Scenario A ending up being sold as 'biltong hunts' at half of the international trophy fee, the full value of the trophies under Scenario A is not realised.

HUNT PACKAGING

Table D: Packaging of Hunts with Roan, Sable and Tsessebe not available

	Elephant	Buffalo	Lion	Leopard	Plains Game			
					A	B	C	D
Overall Quota	5	6	2	3	0	23	95	895
Trophy fee	7,500	5,000	4,000	3,000	2,000	1,500	750	375
Hunt composition								
Big Game Safari	1	1	1	1	1	2	3	6
Elephant hunt	1					2	3	6
Buffalo hunt		1				2	3	6
Cat Hunt				- 1 -		2	3	6
Premier Plains Game					1	1	2	5
Plains Game						1	2	5
Biltong hunt							- 10 -	
Quota remaining after –								
Big Game Safaris	3	4	0	1	0	17	89	883
Elephant hunts	0	4	0	1	0	11	80	865
Buffalo hunts	0	0	0	1	0	3	68	841
Cat hunts			0	1	0	1	65	835
Premier Plains Game hunts				0	0	1	65	835
Plains Game hunts						0	63	830
Biltong hunts							0	0
	Big Game Safaris	Elephant Hunts	Buffalo Hunts	Cat Hunts	Premier Plains Game	Plains Game Hunts	Biltong Hunts	TOTALS
Number of hunts	2	3	4	1	0	1	89	100
Elephant	2	3						5
Buffalo	2		4					6
Lion	2			0				2
Leopard	2			1				3
Plains Game species - A	0				0			0
Plains Game species - B	6	6	8	2	0	1		23
Plains Game species - C	6	9	12	3	0	2	63	95
Plains Game species - D	12	18	24	6	0	5	830	895
Safari days	21	16	14	12	10	8	6	
Total Hunter days	42	48	56	12	0	8	534	700
Daily rates	1,500	1,250	1,000	750	500	250	100	
Gross income daily rates	63,000	60,000	56,000	9,000	0	2,000	53,400	243,400
Trophy fees	57,000	45,000	50,000	10,500	0	4,875	179,250	346,625
GROSS INCOME	120,000	105,000	106,000	19,500	0	6,875	232,650	590,025
Gross income/hectare								5.90

HUNT PACKAGING

Table E: Packaging of Hunts with Roan, Sable and Tsessebe available

	Elephant	Buffalo	Lion	Leopard	Plains Game			
					A	B	C	D
Overall Quota	5	6	2	3	60	37	83	598
Trophy fee	7,500	5,000	4,000	3,000	2,000	1,500	750	375
Hunt composition								
Big Game Safari	1	1	1	1	1	2	3	6
Elephant hunt	1					2	3	6
Buffalo hunt		1				2	3	6
Cat Hunt				- 1 -		2	3	6
Premier Plains Game					1	1	2	5
Plains Game						1	2	5
Biltong hunt								- 10 -
Quota remaining after –								
Big Game Safaris	3	4	0	1	58	33	77	586
Elephant hunts	0	4	0	1	58	27	68	568
Buffalo hunts	0	0	0	1	58	19	56	544
Cat hunts			0	1	58	17	53	538
Premier Plains Game hunts				0	0	0	0	144
Plains Game hunts						0	0	144
Biltong hunts							0	0
	Big Game Safaris	Elephant Hunts	Buffalo Hunts	Cat Hunts	Premier Plains Game	Plains Game Hunts	Biltong Hunts	TOTALS
Number of hunts	2	3	4	1	58	0	14	82
Elephant	2	3						5
Buffalo	2		4					6
Lion	2			0				2
Leopard	2			1				3
Plains Game species - A	2				58			60
Plains Game species - B	4	6	8	2	17	0		37
Plains Game species - C	6	9	12	3	53	0	0	83
Plains Game species - D	12	18	24	6	394	0	144	598
Safari days	21	16	14	12	10	8	6	
Total Hunter days	42	48	56	12	580	0	84	822
Daily rates	1,500	1,250	1,000	750	500	250	100	
Gross income daily rates	63,000	60,000	56,000	9,000	290,000	0	8,400	486,400
Trophy fees	58,000	45,000	50,000	10,500	329,000	0	27,000	519,500
GROSS INCOME	121,000	105,000	106,000	19,500	619,000	0	35,400	1,005,900
Gross income/hectare								10.06

Tables D and E are fully automated and will package any given hunting quota provided the first assumption below is met. The overall quotas from Table C are automatically transferred to the first row of Tables D and E and the hunt packaging follows from these quotas. The composition of each of the types of hunt (e.g. Big Game Safari, Elephant hunt etc) can be adjusted by entering the number of animals in each species category which will be taken on the particular type of hunt and the hunt packaging process will take this into account.

ASSUMPTIONS for Table D and Table E

1. There are more Category D animals than will be needed to complement the main hunts and the balance of these animals will end up in biltong hunts.
 2. A Big Game Safari is a 21 day hunt at a daily rate of US\$1,500/day. It includes an elephant, 1 buffalo, both large cats, 1 Category A, 2 Category B, 3 Category C and 6 Category D Plains Game species (which includes an allowance for baits for the cats). If there are insufficient Category A or Category B Plains Game animals, the deficit is made up with Category B and Category C Plains Game animals.
 3. An Elephant Hunt is an 16 day safari at a daily rate of US\$1,250 day. It includes an elephant, 2 Category B, 3 Category C and 6 Category D Plains Game species. If there are insufficient Category B Plains Game animals, the deficit is made up with Category C Plains Game animals.
 4. A Buffalo Hunt is a 14 day safari at a daily rate of US\$1,000 day. It includes a buffalo, 2 Category B, 3 Category C and 6 Category D Plains Game species.
 5. A Cat hunt is a 12 day safari at a daily rate of US\$750/day. It includes a lion or a leopard, 2 Category B, 3 Category C and 6 Category D Plains Game species (which includes an allowance for baits for the cat). If there are insufficient Category B animals to make up the quota, Category C animals are used.
 6. A Premier Plains Game safari is a 10 day hunt at US\$500/day which includes 1 Category A, 1 Category B, 2 Category C and 5 Category D Plains Game species. If the needed number of Category B species are not available, then they are substituted with Category C species. If the needed number of Category C species are not available, then they are substituted with Category D species.
 7. A Plains Game safari is a 8 day hunt at US\$250/day which includes 1 Category B, 2 Category C and 5 Category D Plains Game species.
 8. The remaining animals are sold on 6 day Biltong Hunts at US\$100/day. The typical number of animals expected to be taken on a Biltong Hunt is 10 and the trophy fees are halved.
 9. All hunts are assumed to be carried out by a single client.
 10. All financial amounts are in United States dollars.
-

Table F: Calculation of Operating Costs for 250 km² *All figures are in United States dollars*

CAPITAL (Capital costs are depreciated over 5 years and added to operating costs)					
#	ITEM	Quantity	Unit Cost US\$	Amounts	Totals
1	Vehicles				
2	4x4	5	20,000	100,000	
3	Fuel Storage	1	250	250	
4	Tools	1	500	500	
5	Vehicle Spares	1	1,000	1,000	101,750
6	Accommodation				
7	Clients	3	1,000	3,000	
8	Staff - senior	4	500	2,000	
9	Staff - junior	17	200	3,400	
10	Bathrooms	6	200	1,200	
11	Kitchen	1	500	500	
12	Dining Room	1	300	300	10,400
13	Equipment				
14	Refrigerators	2	300	600	
15	Deep Freeze	1	300	300	
16	Furniture	1	300	300	
17	Pots, pans, cutlery, crockery	1	500	500	
18	Lighting	1	1,500	1,500	3,200
19	Water supply	1	1,500	1,500	1,500
20				TOTAL	116,850
21	CAPITAL: Amount to be recovered annually				23,370
22	OPERATING COSTS				
23	Staff salaries (costs are for 6 month hunting season)				
24	Professional Hunter US\$/day	180	200	36,000	
25	Learner Hunter	1	4,000	4,000	
26	Camp Manager	1	3,000	3,000	
27	Cooks	2	1,000	2,000	
28	Waiters	2	500	1,000	
29	Scouts	5	300	1,500	
30	Skinners	2	400	800	
31	Trackers	2	400	800	
32	Driver	1	500	500	
33	PR/Community relations	1	2,000	2,000	
34	General workers	3	200	600	52,200
35	Vehicles				
36	Fuel (litres)	10000	1	10,000	
37	Lubricants (litres)	100	5	500	10,500
38	Camp				
39	Annual refurbishment	1	5,000	5,000	
40	Gas (kg)	200	5	1,000	
41	Miscellaneous	1	1,000	1,000	7,000
42	Food and drink				
43	Clients (2) US\$/day	180	50	9,000	
44	Senior staff (4)	180	20	3,600	
45	Junior Staff (17)	180	20	3,600	16,200
	Total annual operating costs for 250km ² , including capital replacement				109,270
	Annual operating costs/ha				4.37

NOTES on Table F (row number references)

The scenario with roan, sable and tsessebe present (**Table E**) results in a total of 822 hunter days of which 738 entail international safaris and the remainder (84) are biltong hunts. This would justify 4 separate operators in an area of 1,000km² each with approximately 180 days of hunting (ignoring the biltong hunts).

Without roan, sable and tsessebe, there are only 166 international safari hunter days which would only justify one safari operator. The remaining hunt days (534) are all in the biltong hunt category. This theoretically reduces the operating costs per hectare by a factor of 4.

2. Vehicles are for (a) Professional hunter (b) Learner hunter (c) Camp manager (d) PR officer (e) Standby
4. All vehicle maintenance is done on site. Provision for tyres, tubes etc is included under spares.
6. Rustic or tented accommodation will be used for the hunting camp
7. Provision has been made for up to three clients
18. Lighting includes a 25kva generator
19. Water supply includes 2 pumps, watertank and piping
21. All capital items are written off over 5 years and the total capital cost is included in the operating costs
24. The professional hunter is paid on a daily rate of US\$200/day for actual days hunted
29. Scouts are used for anti-poaching in 250km²
36. Fuel provision includes generator and water pumps.
39. 'Annual refurbishment' includes camp maintenance costs during the hunting season and at start-up each year
40. Gas is used in the kitchen for cooking
41. 'Miscellaneous' camp operating costs includes cleaning materials, toilet paper, cooking oils, salt, pepper, sauces, napkins etc.
42. All staff are fed whilst in the field

For the purposes of this exercise, all extra charges such as government tourism levies, CITES tags and documentation, transfers to hunting camps etc. are assumed to be passed on to the client with no mark-ups.

Table G: COST AND LAND USE SUMMARY

	WITHOUT ROAN, SABLE AND TSESSEBE	WITH ROAN, SABLE AND TSESSEBE
Area	4,000 km ²	4,000 km ²
Gross income US\$/hectare	5.90	10.06
Operating costs US\$/hectare	1.09	4.37
Net income US\$/hectare	4.81	5.69
Potential net earnings from 4,000sq.km	1,923,020	2,275,280

NOTES

1. It is assumed that these values could be achieved over a core area of about 4,000km² in the Caprivi.
2. Gross income/hectare is calculated in Tables D & E.
3. Operating costs are calculated in Table F. The operating costs for the scenario without roan, sable and tsessebe have been reduced to one quarter of the amount when these species are present because it would require only one safari operator to realise the full value of the available hunting – as opposed to 4 safari operators in the other case.
4. This net income includes no payments to government or local communities for the safari concession. However, the net income to the safari operator indicates the margins available for these payments.

Protected Area Requirements in Southern Africa

Martin (1996) empirically derived the relationship that the number of men required for effective patrolling against illegal hunting in any park was approximately equal to the square root of the area of the park. The relationship was based on the relative success of the different protected areas in Zimbabwe using the criterion that, under effective patrolling, illegal hunters will be found within less than two days.

$$\text{Number of men: } N_s = \sqrt{A}$$

— where A is expressed in square kilometres

PARK SIZE km ²	1	5	10	50	100	500	1,000	5,000	10,000
NUMBER OF MEN	1	2	3	7	10	22	32	71	100

Martin (1997) developed standard spreadsheets for calculating the operating costs and capital requirements based on this relationship. The number of men determines the annual running cost for any park. The budget is made up of salaries, field allowances, equipment, transport and maintenance costs and includes provisions for senior field and research staff. Allowing for variations in salaries and fuel costs from country to country in the region, the operational costs are approximately given by the formula —

$$\text{Annual Recurrent Expenditure/km}^2 \quad C_R = \text{US\$}50 \left(1 + \frac{2}{A} + \frac{3}{\sqrt{A}} \right)$$

The capital requirements to set up a new park from scratch are also dependent on the total staff complement in the park and vary slightly depending on building costs across the region. The required capital per unit area is approximately given by —

$$\text{Total Capital Expenditure/km}^2 \quad C_C = \text{US\$}500 \left(1 + \frac{1}{A} + \frac{1}{\sqrt{A}} \right)$$

— where A is expressed in thousands of square kilometres in both formulas

STAFF NUMBERS, OPERATING COSTS AND CAPITAL REQUIREMENTS FOR VARIOUS PARK SIZES

PARK SIZE km ²	Number of Field Staff Required	Operating Costs US\$/km ² /year	Total Operating Cost US\$/year	Capital Required US\$/km ²	Total Capital US\$
1	1	104,793	104,793	516,311	516,311
2	1	53,404	106,808	261,680	523,361
5	2	22,171	110,857	107,571	537,855
10	3	11,550	115,500	55,500	555,000
20	4	6,111	122,213	29,036	580,711
50	7	2,721	136,041	12,736	636,803
100	10	1,524	152,434	7,081	708,114
200	14	885	177,082	4,118	823,607
500	22	462	231,066	2,207	1,103,553
1,000	32	300	300,000	1,500	1,500,000
2,000	45	206	412,132	1,104	2,207,107
5,000	71	137	685,410	824	4,118,034
10,000	100	107	1,074,342	708	7,081,139
20,000	141	89	1,770,820	637	12,736,068
50,000	224	73	3,660,660	581	29,035,534
100,000	316	66	6,600,000	555	55,500,000

BIBLIOGRAPHY

Numerous people have kindly contributed to this bibliography and not all of the papers listed below have been read by the author. Papers which are referred to in the text are shown in black print.

- AME (1982). **República Popular de Angola: Atlas Geográfico, Volume 1**. Ministério da Educação, República Popular de Angola, Impresso por Esselte Map Service, Estocolmo, Suécia 1982. 49pp
- Ansell (1972). **Part 2. 15 Family Artiodactyla**. In: *The Mammals of Africa: an identification manual*. Eds. J. Meester and H.W. Setzer. Smithsonian Institution Press, Washington D.C. 1-84
- ASG (1989). **Antelopes - Global Survey and Regional Action Plans. Part 2. Southern and Central Africa**. IUCN/SSC Antelope Specialist Group, Compiler R. East, IUCN, Gland, Switzerland. 96pp
- ASG (1998). **African Antelope Data Base 1998**. IUCN/SSC Antelope Specialist Group, Compiler R. East, IUCN, Gland, Switzerland. 350pp
- Astley Maberly, C.T. (1952). **The roan antelope**. African Wildlife 6(4). 287-292
- Barnes J.I. (2001). **Economic returns and allocation of resources in the wildlife sector of Botswana**. South African Journal of Wildlife Research 31(3&4): 141-153
- Barnes J.I. (2002a). **Financial and Economic analysis of Mayuni Conservancy**. Spreadsheet data. Environmental Economics Unit, Directorate of Environmental Affairs, Ministry of Environment and Tourism.
- Barnes J.I. (2002b). **Financial and Economic analysis of Salambala Conservancy**. Spreadsheet data. Environmental Economics Unit, Directorate of Environmental Affairs, Ministry of Environment and Tourism.
- Barnes J.I. and J.L.V. de Jager (1995). **Economic and financial incentives for wildlife use on private land in Namibia and the implications for policy**. Research Discussion Paper Number 8, Directorate of Environmental Affairs, Ministry of Environment and Tourism, Namibia. 20pp
- Barnes, Jonathan, James Cannon and Karl Morrison (2001). **Economic Returns to selected Land Uses in Ngamiland, Botswana**. Consultancy for Conservation International. 60pp + Appendices
- Barnes, Jonathan I., James Macgregor and L. Chris Weaver (2002). **Economic efficiency and incentives for change within Namibia's community wildlife use initiatives**. World Development Vol 30 (4): 667-681
- Bell, R.H.V.(1986) **The Workshop Theme: Adaptive Management**. In: *Conservation and Wildlife Management in Africa*. Proc. Workshop sponsored by the US Peace Corps, Kasungu National Park, Malawi. Edited by R.H.V. Bell and E. McShane-Caluzi. Peace Corps, Washington, 1986.
- Best, A.A. and T.G.W. Best (1977). **Rowland Ward's Records of Big Game**. XVII Edition (Africa), Rowland Ward Publications Ltd., Sussex, England.
- Bigalke, R. (1955). **Notes on the former and present occurrence of the sable antelope in the southern part of the Kruger National Park**. Fauna and Flora 6. 117-123
- Brown, C.J. (2000). **Historic Distribution of Large Mammals in the Gondwana Canyon Park area of southern Namibia**. Unpublished report, Namibia Nature Foundation. 7pp + 1 table
- CARACAL (2002). **Botswana-namibia Transboundary Ungulate Migrations: Range Movement Corridors, Resource Limitations and Population Viability**. Project Proposal submitted to African Wildlife Foundation. 10pp

- CARACAL (2002). **Transboundary Natural Resource Management: Wildlife Conflict with Communities in the Four Corners Transboundary Conservation Area.** Project Proposal submitted to African Wildlife Foundation. 24pp
- Castley J.G., D. Zimmerman & D. Joubert (1999). **Roan antelope management strategy within South African National Parks.** Draft unreferenced document provided by C.J. Brown
- Child, G., H. Robbel and C.P. Hepburn (1972). **Observations on the biology of the tsessebe, *Damaliscus lunatus lunatus*, northern Botswana.** Mammalia 36: 342-388.
- Coe M.J., Cumming D.H. and J. Phillipson (1976). **Biomass and Production of large African Herbivores in relation to Rainfall and Primary Production.** Oecologia (Berl.) 22, 341-354.
- Coetsee B.J. (1980). **Classification is instructive – comments on a published table of antelope habitat preferences.** Koedoe 23: 33-34
- Corbett, Andrew and Brian T.B. Jones (2000). **The Legal Aspects of Governance in CBNRM in Namibia.** Directorate of Environmental Affairs Research Discussion Paper Number 41, October 2000, Ministry of Environment and Tourism, Republic of Namibia. 25pp
- Craig, G.C. (1998). **Aerial Survey of Northern Namibia.** Survey carried out for Directorate of Scientific Services, Ministry of Environment and Tourism, Republic of Namibia.
- Craig, G.C. (2000). **Aerial Survey of North Western Namibia.** Survey carried out for Directorate of Scientific Services, Ministry of Environment and Tourism, Republic of Namibia.
- Cumming, David H.M. (1999). **Study on the Development of Transboundary Natural Resource Management Areas in Southern Africa – Environmental Context: Matural Resources, Land Use and Conservation.** Biodiversity Support Programme. Washington D.C. USA. 56pp + 30 maps
- DAHSD (Undated). **The Veterinary Cordon Fence: Past, Present and Future.** Information booklet published by the Division of Animal Health Surveillance & Development, Directorate of Veterinary Services, Private Bag 12022, Ausspannplatz, Namibia. 14pp
- DFID (1997a). **Ngamiland Fences Scoping Exercise – Botswana.** Consultancy Report funded by the UK Department for International Development, Gaborone, carried out by Mouchel Consulting Limited, U.K. and Price Waterhouse, Zimbabwe, for the Department of Animal Health and Production, Botswana, July 1977. 59pp
- DFID (1997b). **Initial Recommendations for Mitigation Measures applicable to the Caprivi Border Fence.** Consultancy Report funded by the UK Department for International Development, Gaborone, carried out by Mouchel Consulting Limited, U.K. and Price Waterhouse, Zimbabwe, for the Department of Animal Health and Production, Botswana, July 1977. 14pp
- Dörgeleh W.G. (1998). **Habitat selection of a roan antelope (*Hippotragus equinus*) population in mixed bushveld, Nylsvlei Nature Reserve.** S.Afr.J.Wildl.Res. 28(2):47-57
- DSS (2002). **Aerial Survey Summary (1968 - 2001).** Compiler: Uatjavi Uanivi, Directorate of Scientific Services, Ministry of Environment and Tourism, Republic of Namibia. 54pp
- DSS (2003). **National Estimates for Roan, Sable and Tsessebe, 2003.** Compiler: P. Erb, Directorate of Scientific Services, Ministry of Environment and Tourism, Republic of Namibia. 3pp
- Du Toit, R.F. (1992). **Status, distribution and management of sable in Zimbabwe.** Symposium on the Sable Antelope as a Game Ranch Animal, Wildlife Group, South African Veterinary Association, Onderstepoort: 79-87

- Dunham, K. M. and E.F. Robertson (2001). **Trends in rainfall explain the rise and fall of tsessebe in Kruger National Park.** Scientific Report 01/01, South African National Parks. 33pp
- Dunham, Kevin M., E.F. Robertson and Clive M. Swanepoel (2003). **Population decline of tsessebe antelope (*Damaliscus lunatus lunatus*) on a mixed cattle and wildlife ranch in Zimbabwe.** Biological Conservation 113(1): 111-124
- DVS (2002). **Purposes, Structure and Functions.** Directorate of Veterinary Service, Ministry of Agriculture, Water and Rural Development, Republic of Namibia. 10pp
- DWNP (2002a). **Controlled Hunting Areas – Botswana.** Map supplied by courtesy of Department of Wildlife and National Parks, Botswana.
- Erb, Karl Peter (1993). **The Roan antelope (*Hippotragus equinus*, Desmarest 1804), its ecology in the Waterberg Plateau Park.** Msc Thesis, University of Stellenbosch. 109pp
- Erb, Peter (1992). **Notes on the roan, sable and tsessebe populations in the Waterberg Plateau Park.** Personal research notes made available for this study. 24pp
- Gaerdes, J. (1963). **Fahrt durch das ‘Kung-Kaukau-Veld zu den Aha-Bergen.** Mitteilungen, S.W.A. Wissenschaftliche Gesellschaft IV (2), February 1963: 3-6
- Gaerdes, J. (1969). **Kuhantilopen in Suedwestafrika.** Mitteilungen, S.W.A. Wissenschaftliche Gesellschaft IX (12), March 1969: xii-xiv
- Garstang, Richard (1982). **An analysis of home range utilisation by the tsessebe, *Damaliscus lunatus lunatus* (Burchell), in P W Willis Private Nature Reserve.** Msc Thesis, University of Pretoria. 165pp + appendices, tables and maps.
- Grant C.C. and J.L. van der Walt (2000). **Towards an adaptive management approach for the conservation of rare antelope in the Kruger National Park – outcome of a workshop held in May 2000.** Koedoe 43(2): 103-111
- Grant C.C., T. Davidson, P.J. Funston and D.J. Pienaar (2002). **Challenges faced in the conservation of rare antelope: a case study on the northern basalt plains of the Kruger National Park.** Koedoe 45(2):45-66
- Grobler, J.H. (1973). **Biological data on the tsessebe, *Damaliscus lunatus lunatus*, (Mammalia: Alcelaphinae) in Rhodesia.** Arnoldia 6:1-16
- Grobler, J.H. (1974). **Aspects of the biology, population ecology and behaviour of the sable *Hippotragus niger niger* (Harris, 1838) in the Rhodes Matopos National Park, Rhodesia.** Arnoldia Rhod. 7(6):1-36
- Grobler, J.H. (1978). **Population dynamics of sable *Hippotragus niger niger* (Harris, 1838) in Rhodesia.** D.Phil Thesis, Univ. of Rhodesia.
- Grobler J.H. (1979). **The use of horn sections to determine the age of sable *Hippotragus niger*.** Koedoe 22: 219-223
- Grobler J.H. (1980). **Body growth and age determination of the sable *Hippotragus niger* (Harris 1838).** Koedoe 23: 131-156
- Grobler J.H. (1981a). **Parasites and mortality of sable *Hippotragus niger niger* (Harris 1838) in the Matopos, Zimbabwe .** Koedoe 24: 119-123
- Grobler, J.H. (1981b). **Feeding behaviour of sable *Hippotragus niger niger* (Harris, 1838) in the Rhodes Matopos National Park, Zimbabwe.** S.Afr.J.Zool. 16:50-58

- Grobler, J.H. (1984). **The roan antelope**. Quagga 8. 5-11
- Hahn, H.S. (1925). **Big Game: Distribution in Ovamboland**. Report to the Secretary for South West Africa from the Officer in Charge of Native Affairs, Ondonga, 10th September 1925. 36pp
- Harrington, R. (1995). **Herbivore and habitat change associated with the roan antelope decline in the Northern Kruger National Park**. M.Sc. Dissertation, University of the Witwatersrand, Johannesburg. 148 pp.
- Harrington, R., N. Owen-Smith, P.C. Viljoen, H.C. Biggs, D.R. Mason and P. Funston (1999). **Establishing the causes of the roan antelope decline in the Kruger National Park, South Africa**. Biol. Cons. 90: 69-78
- Hatting H.J. (1988). **Comparative quantitation of the physiological response to acute stress in impala and roan antelope**. Comparative Biochemistry and Physiology A 89(4). 547-551
- Holling C.S. (1978) Ed. **Adaptive Environmental Assessment and Management**. John Wiley & Sons, New York.
- Humavindu, Michael N. (2001) **Trophy Hunting in the Namibian Economy: an Assessment**. Environmental Economics Unit, Directorate of Environmental Affairs, Ministry of Environment and Tourism. 6pp + Tables and Appendices
- Huntly B.J. (1972). **Observations on the Percy Fyfe Nature Reserve tsessebe population**. Ann. Transv. Mus. 27:225-239
- Huntly B.J. (1973). **Ageing Criteria for Tsessebe (*Damaliscus lunatus lunatus*)**. J. South African Wildlife Management Association, 3: 24-27
- IUCN (1997). **Conservation Assessment and Management Plan (CAMP)**. Appendix II: Taxon Data Sheet: Organisation and Definitions. The World Conservation Union, Gland, Switzerland. 14pp
- IUCN ROSA (1992). **Angola: Environment Status Quo Assessment Report. Main Report**. IUCN Regional Office for Southern Africa, October 1992. Study funded by the Commission of the European Community (PN B-7-5040/92/10). 256pp
- Joubert, E. and P.K.N. Mostert (1975). **Distribution Patterns and Status of some Mammals in South West Africa**. Madoqua 9(1): 5-44.
- Joubert, S.C.J. (1972). **Territorial behaviour of the tsessebe (*Damaliscus lunatus lunatus* Burchell) in the Kruger National Park**. Zoologica Africana 7(1). 141-156
- Joubert, S.C.J. (1975). **The mating behaviour of the tsessebe (*Damaliscus lunatus lunatus*) in the Kruger National Park**. Zeitschrift für Tierpsychologie 37. 182-191
- Joubert, S.C.J. (1976). **The population ecology of the roan antelope, *Hippotragus equinus equinus* (Desmarest, 1804), in the Kruger National Park**. D.Sc.thesis, University of Pretoria.
- Joubert S.C.J. and P.J.L. Bronkhorst (1977). **Some aspects of the history and population ecology of the tsessebe *Damaliscus lunatus lunatus* in the Kruger National Park**. Koedoe 20: 125-145
- Lange, Glenn-Marie, Jon I. Barnes and Daniel J. Motinga (1997). **Cattle Numbers, Biomass, Productivity and Land Degradation in the Commercial Farming Sector of Namibia**. Research Discussion Paper Number 17, Directorate of Environmental Affairs, Ministry of Environment and Tourism, Namibia. 35pp
- Leslie, P.H. (1984). **Some Further Notes on the Use of Matrices in Population Analysis**. Biometrika 35: 213-245.

- LIFE (2002). **Game Population Estimates for Nyae Nyae Conservancy, January 2002.** Annex One of an annual report on Nyae Nyae Conservancy from the WWF LIFE programme, Windhoek, Namibia.
- Martin, R.B. (1986). **Communal Areas Management Programme for Indigenous Resources (CAMPFIRE).** Publication of the Department of National Parks and Wild Life Management, Zimbabwe. 34pp
- Martin R.B. (1996). **Costs of Conserving State Protected Areas in Southern Africa.** Fact Sheet No.6, Africa Resources Trust, P.O. Box A860, Avondale, Harare, Zimbabwe.
- Martin R.B.(1997). **The Revised Investment Programme for Forestry and Wildlife in Mozambique.** Consultancy for Ministry of Agriculture and Fisheries, Maputo, Mozambique. FAO (Rome).
- Martin R.B. (1999). **Adaptive Management: The Only Tool for Decentralised Systems.** Paper presented at a Norway/UN Conference titled *The ecosystem approach for sustainable use of biological diversity* held in Trondheim, Norway, 6-10th September 1999. 14pp
- Martin R.B. (2002a). **Common Property on a Grand Scale: The “Four-Corners” Transfrontier Conservation Area.** Article for the Centre for Applied Social Sciences Trust on the Conference of the International Association for the Study of Common Property in Victoria Falls, June 2002.
- Martin, R.B. (2002b) **Conditions for Effective, Stable and Equitable Conservation at the National Level in Southern Africa.** Paper prepared for a Workshop titled *Local Communities, Equity and Protected Areas* to be held from 24-26th October in Durban, South Africa, hosted by The Programme for Land and Agrarian Studies (PLAAS), Africa Resources Trust (ART) and IUCN. 23pp
- Martin, R.B. (2002c). **Species Report for Southern Savanna Buffalo.** Study conducted in support of *The Transboundary Mammal Project* of the Ministry of Environment and Tourism, Namibia facilitated by The Namibia Nature Foundation and the World Wildlife Fund LIFE Programme (Living in a Finite Environment). 97pp
- Matthee,C.A. & T.J.Robinson (1999). **Mitochondrial DNA population structure of roan and sable antelope: implications for the translocation and conservation of the species.** *Molecular Ecology* 8: 227-238
- McLoughlin, C.A. (2001). Population viability assessment of rare antelope in the Kruger National Park. M.Sc. thesis, University of the Witwatersrand, Johannesburg.
- Mendelsohn, John, Selma el Obeid and Carole Roberts (1997). **A Profile of North-Central Namibia.** Environmental Profiles Project, Directorate of Environmental Affairs, Ministry of Environment and Tourism, Republic of Namibia. Gamsberg Macmillan Publishers. 79pp
- Mendelsohn, John and Carole Roberts (1997). **An Environmental Profile and Atlas of Caprivi.** Directorate of Environmental Affairs, Ministry of Environment and Tourism, Republic of Namibia. 45pp
- MET (2000). **Atlas of Namibia.** Notes on the Atlas website. Ministry of Environment and Tourism, Republic of Namibia. 5pp
- MET (2002). **Conservation and the Environment in Namibia.** Eds. Sem Shikongo and Phoebe Barnard, Publication for the World Summit on Sustainable Development (WSSD), August/September 2002, Johannesburg, Ministry of Environment and Tourism, Republic of Namibia. 56pp
- Mills M.G.L., H.C. Biggs and I.J. Whyte (1995). **The relationship between rainfall, lion predation and population trends in African herbivores.** *Wildl. Res.* 22:75-88

- Murphree, Marshall W. (1998). **Incentives for Sustainability**. Keynote Address at the Symposium Workshop on *Conservation*, 50th Anniversary Celebration of IUCN, Fontainebleau, 3-5 November 1998. 16pp
- Murphree, Marshall W. (2000). **Boundaries and Borders: The Question of Scale in the Theory and Practice of Common Property Management**. Paper presented at the Eighth Biennial Conference of the International Association for the Study of Common Property (IASCP), Bloomington, Indiana, USA 31 May-4 June 2000.
- Murphree, Marshall W. (2002). **Protected Areas and the Commons**. CPR Forum Commentary, The Common Property Resource Digest No.60, March 2002. pp1-3.
- Nelson, E. (1926). **Distribution of Game in South West Africa**. Government Report from the Game Warden of Namutoni Game Reserve, 23rd March 1926.
- Newton da Silva, S. (1970). **A Grande Fauna Selvagem de Angola**. Edição da Direcção Provincial dos Serviços de Veterinaria. 151pp.
- el Obeid, Selma and John Mendelsohn (2001). **A Preliminary Profile of the Kavango Region in Namibia**. Publ. Namibia Nature Foundation, P.O. Box 245, Windhoek, Namibia. 45pp
- van Oudtshoorn, Fritz and Eben van Wyk (1999). **Guide to the Grasses of Southern Africa**. Briza Publications, Pretoria. 288pp
- Penzhorn, B.L. (1968). **Saving the roan antelope**. *Oryx* 9(5). 333-334
- Penzhorn R.L. and N.J. van der Merwe (1993). **Testis size and spermatogenesis of tsessebes from Nylovley Nature Reserve, Transvaal**. *Koedoe* 36/1:95-98
- Perrin M.R. and C.L. Taolo (1999). **Diet of introduced roan antelope at Weenen Nature Reserve**. *S.Afr.J.Wildl. Res.* 29(2):43-51
- Pienaar U. De V. (1961). **A second outbreak of anthrax amongst game animals in the Kruger National Park**. *Koedoe* (4), 4-17
- Pienaar, U.de V (1963). **The large mammals of the Kruger National Park – their distribution and present-day status**. *Koedoe* 6: 1-37
- Pienaar U. De V. (1974). **Habitat preferences in South African antelope species and its significance in natural and artificial distribution patterns**. *Koedoe* 17: 185-195
- PW (1998). **Namibia North-East Parks Project. Volume 1: Background and Inventory**. Price Waterhouse Consultants' Report, 15th April, 1998. 108pp + photographs
- PWC (1999a). **Namibia North-East Parks Project. Volume 2: Kwando Core Area Management Plan**. Price Waterhouse Coopers Consultants' Report, March 1999. 54pp + Appendices
- PWC (1999b). **Namibia North-East Parks Project. Volume 2: Mamili National Park Management Plan**. Price Waterhouse Coopers Consultants' Report, May 1999. 54pp + Appendices
- PWC (1999c). **Namibia North-East Parks Project. Volume 2: Khaudom National Park Management Plan**. Price Waterhouse Coopers Consultants' Report, May 1999. 49pp + Appendices
- PWC (1999d). **Namibia North-East Parks Project. Volume 2: Mudumu National Park Management Plan**. Price Waterhouse Coopers Consultants' Report, May 1999. 48pp + Appendices
- PWC (1999e). **Namibia North-East Parks Project. Okavango National Park Integrated Development Plan**. Price Waterhouse Coopers Consultants' Report, May 1999. 36pp + Appendices

- RARE (2002). **The Rare Antelope Research Enterprise (RARE): Ecology and conservation biology of threatened species.** Project proposal of the Centre for African Ecology, School for Animal, Plant and Environmental Sciences, University of the Witwatersrand, 14 March 2002. 6pp
- Rodwell, T.C., J. Tagg and M. Grobler (1995). **Wildlife Resources in the Caprivi, Namibia: Results of an Aerial Census in 1994 and Comparisons with past Surveys.** Research Discussion Paper No. 9, Ministry of Environment and Tourism, Republic of Namibia. 29pp
- RN (1975). **Nature Conservation Ordinance, 1975.** Legislative Act No.4 of 1975 (with amendments), Government of the Republic of Namibia.
- Ruitenbeek, Jack and Cynthia Carter (2001). **The Invisible Wand: Adaptive Co-management as an Emergent Strategy in Complex Bio-Economic Systems.** Occasional Paper No.34, Centre for International Forestry Research, Bogor, Indonesia. 47pp
- Scott Wilson (2000). **Environmental Assessment of Veterinary Fences in Ngamiland – Summary Report.** Consultants' Report to the Government of Botswana, Scott Wilson Resource Consultants in association with The Environment and Development Group (EDG). 106pp
- Shortridge, G.C. (1934). **The Mammals of South West Africa.** Vol I & II. William Heineman Ltd., London.
- Sinclair, A.R.E. (1974a). **The Natural regulation of Buffalo Populations in East Africa. I. Introduction and Resource Requirements.** East African Wildlife Journal 12(2): 135-154
- Sinclair, A.R.E. (1974b). **The Natural regulation of Buffalo Populations in East Africa. III. Population Trends and Mortality.** East African Wildlife Journal 12(3): 185-200
- Sinclair, A.R.E. (1974c). **The Natural regulation of Buffalo Populations in East Africa. IV. The Food Supply as a Regulating Factor, and Competition.** East African Wildlife Journal 12(4): 291-311
- Skead, C.J. (1982). **Historical Mammal Incidence in the Cape Province.** Volume I, The Western and Northern Cape. 2nd Impression . 605-606.
- Smithers, Reay H.N. (1983). **The Mammals of the Southern African Subregion.** University of Pretoria, Pretoria, RSA. 736pp
- Tomkinson, R. (1957). **Sable at Shingwedzi.** Oryx 4(2). 137-139
- Travel News Namibia (2002). **Conservation 2002 and the Environment in Namibia.** Special edition produced for the World Summit on Sustainable Development, Venture Publications in collaboration with the Ministry of Environment and Tourism, Windhoek. 56pp
- ULG (1994). **Aerial Census of Animals in the Caprivi Strip, Namibia, Dry Season 1994.** Consultants' Report to the Department of Wildlife and National Parks, Botswana. 21pp
- ULG (1995). **Final Report – Aerial Surveys.** Consultants' Report to the Department of Wildlife and National Parks, Botswana. ULG Consultants, Birmingham Road, Saltisford. Warwick CV34 4TT, England. 59pp
- Viljoen, P.C. & D.J. Pienaar. 1993. **Management proposals for the northern basalt plains in the Kruger National Park with special reference to roan antelope.** Scientific Report, Skukuza 12/93
- de Vos, V., G.L. van Rooyen and J.J. Kloppers (1973). **Anthrax immunization of free-ranging roan antelope *Hippotragus equinus* in the Kruger National Park.** Koedoe (16); 11-26
- de Vos, V. and G.D. Imes (1976). **An outbreak of dermatophilosis in sable *Hippotragus niger* and roan *Hippotragus equinus* in the Kruger National Park.** Koedoe 19: 1-15

- de Vos, V. and H.B. Bryden (1995). **The epidemiology of a major anthrax outbreak in the Kruger National Park.** International Workshop on Anthrax 19-21 Sept 1995, Winchester, UK 1995. 25-26
- de Vos, V. and H.B. Bryden (1998). **Anthrax in the Kruger National Park, South Africa: The role of roan (*Hippotragus equinus*) and sable (*H. niger*) in the epidemiology of anthrax: A preliminary report.** Pp. 33-36. *Proceedings of the ARC-Onderstepoort OIE International Congress with WHO-Cosponsorship on Anthrax, Brucellosis, CBPP, Clostridial and Mycobacterial Diseases, Berg-en-Dal, Kruger National Park, South Africa, 9-15 August 1998.* Onderstepoort: Onderstepoort Veterinary Institute.
- Wilson D.E. (1975). **Factors affecting roan and sable populations on nature reserves in the Transvaal with particular reference to ecophysiological aspects.** D.Sc. Thesis, University of Pretoria.
- Wilson, David E. and Stanley M. Hirst (1977). **Ecology and Factors limiting Roan and Sable Antelope Populations in South Africa.** Wildlife Monographs, No. 54, April 1977. 111pp
- Wilson V.J. (1969). **The large mammals of the Rhodes Matopos National Park.** *Arnoldia Rhod.* 4(12):1-9
- Young, E. (1972). **Overstraining disease (capture myopathy) in the tsessebe *Damaliscus lunatus* and oribi *Ourebia ourebi*.** *Koedoe* 15. 143-144
-