

# *African Herp News*

**Newsletter of the  
Herpetological Association of Africa**



**Number 59**

**APRIL 2013**

## HERPETOLOGICAL ASSOCIATION OF AFRICA

<http://www.wits.ac.za/haa>

### FOUNDED 1965

The HAA is dedicated to the study and conservation of African reptiles and amphibians. Membership is open to anyone with an interest in the African herpetofauna. Members receive the Association's journal, *African Journal of Herpetology* (which publishes review papers, research articles, and short communications – subject to peer review) and *African Herp News*, the Newsletter (which includes short communications, natural history notes, geographical distribution notes, herpetological survey reports, venom and snakebite notes, book reviews, bibliographies, husbandry hints, announcements and news items).

### NEWSLETTER EDITOR'S NOTE

Articles shall be considered for publication provided that they are original and have not been published elsewhere. Articles will be submitted for peer review at the Editor's discretion. Authors are requested to submit manuscripts by e-mail in MS Word '.doc' or '.docx' format.

**COPYRIGHT:** Articles published in the Newsletter are copyright of the Herpetological Association of Africa and may not be reproduced without permission of the Editor.

**The views and opinions expressed in articles are not necessarily those of the Editor.**

### COMMITTEE OF THE HERPETOLOGICAL ASSOCIATION OF AFRICA

#### *CHAIRMAN*

**Aaron Bauer**, Department of Biology, Villanova University, 800 Lancaster Avenue, Villanova, Pennsylvania 19085, USA. [aaron.bauer@villanova.edu](mailto:aaron.bauer@villanova.edu)

#### *SECRETARY*

**Jeanne Tarrant**, African Amphibian Conservation Research Group, NWU. 40A Hilltop Road, Hillcrest 3610, South Africa. [jeannetarrant@ymail.com](mailto:jeannetarrant@ymail.com)

#### *TREASURER*

**Abeda Dawood**, National Zoological Gardens, Corner of Boom and Paul Kruger Streets, Pretoria 0002, South Africa. [abeda@nzg.ac.za](mailto:abeda@nzg.ac.za)

#### *JOURNAL EDITOR*

**John Measey**, Applied Biodiversity Research, Kirstenbosch Research Centre, South African Biodiversity Institute, P/Bag X7, Claremont 7735, South Africa. [john@measey.com](mailto:john@measey.com)

#### *NEWSLETTER EDITOR*

**Bryan Maritz**, Department of Ecology & Evolutionary Biology, Cornell University, Corson Hall, Ithaca, NY 14853, USA. [bryanmaritz@gmail.com](mailto:bryanmaritz@gmail.com)

#### *ADDITIONAL MEMBERS*

**Graham Alexander**, School of Animal, Plant and Environmental Sciences, University of the Witwatersrand, Johannesburg 2050, South Africa. [graham.alexander@wits.ac.za](mailto:graham.alexander@wits.ac.za)

**Michael Bates**, Department of Herpetology, National Museum, P.O. Box 266, Bloemfontein 9300, South Africa. [herp@nasmus.co.za](mailto:herp@nasmus.co.za)

**William Branch**, Curator of Herpetology, Bayworld, P.O.Box 13147, Humewood 6013, South Africa. [wrbranch@bayworld.co.za](mailto:wrbranch@bayworld.co.za)

**Andrew Turner**, Scientific Services, Western Cape Nature Conservation Board, Private Bag 5014, Stellenbosch, 7600, South Africa. [aaturner@capenature.co.za](mailto:aaturner@capenature.co.za)

**COVER PHOTOGRAPH:** *Naja (Boulengerina) annulata annulata* from Lagoa Carumbo, Lunda Norte, Angola. Photograph by: Bill Branch.

## LACERTIDAE

### *Meroles ctenodactylus* (Smith 1838)

#### SMITH'S DESERT LIZARD

On 20 April 2007 three species of *Meroles* were found in close proximity on the eastern edge of the southern dune sea west of Sossusvlei Mountain Lodge, NamibRand Reserve, Southern Namibia (24°50'03.3"S, 15°49'36.7"E 856m). They included two specimens of *Meroles ctenodactylus* (Fig. 1), that were accessioned into the Port Elizabeth Museum (PEM) herpetological collection (PEM R17290-91). They were sympatric with *Meroles cuneirostris* (Strauch 1867) and *Meroles anchietae* (Bocage 1867), also accessioned into the PEM as PEM R17288-89 and PEM R17286 respectively.



**Figure 1:** *Meroles ctenodactylus* (PEM R17290) Sossusvlei Mountain Lodge, Namib-Rand Reserve, Southern Namibia (Photo: Bill Branch).

Microhabitat differences between the species were evident. *Meroles ctenodactylus* were active in the well-vegetated dune slacks and dune

hummocks (left, Fig. 2), *M. cuneirostris* on the more sparsely vegetated dune tops (right, Fig. 2). Both species were found in microsympatry in intermediate habitats. *M. anchietae* was less common and only active on the largest mobile dune slipfaces (bottom right, Fig. 2).



**Figure 2:** Southern dune sea west of Sossusvlei Mountain Lodge, Namib-Rand Reserve, Southern Namibia (Photo: Bill Branch).

Branch (1994) recorded *M. ctenodactylus* from Rooilepel Depression (28° 16' 56" S, 16° 37' 23.5" E 440m) in the Sperrgebiet, southern Namibia, and noted that it appeared to be the northern record for the species. The Sossusvlei Mountain Lodge records extend the species' range a further 300km north, and is the first time that sympatry between these three psammophilic *Meroles* species has been documented.

## ACKNOWLEDGEMENTS

My wife and I thank Peter Dunning and Vernon Swanepoel (Sossusvlei Desert Lodge) and the other staff at the “&Beyond” Lodge for access and hospitality during fieldwork in the region. Specimens were covered under Namibian Research/Collecting Permit 1068/2006.

## REFERENCES

BRANCH, W. R. 1994. Herpetofauna of the Sperrgebiet region of southern Namibia. *Herpetological Natural History* 2(1): 1-11.

## SUBMITTED BY:

**WILLIAM R. BRANCH**, Department of Herpetology, Bayworld (Port Elizabeth Museum), P.O. Box 13147, Humewood 6013, South Africa. E-mail: wrbranch@bayworld.co.za.

\*\*\*\*\*

## GEKKONIDAE

### *Pachydactylus rangei* (Andersson, 1908)

#### Web-footed Gecko

Small dune hummocks along the Orange River, Sperrgebiet National Park (ca. 80km east of Oranjemund and 30km southwest of Rosh Pinah, S28.10 E16.51), Namibia. Sparsely vegetated – mainly stunted *Euclea pseudebenus* trees and shrubs – sandy area between the Orange River riparian vegetation and ancient gravel terraces in December 2012. Observed by Peter Cunningham. 6 specimens – photographed (Fig. 1).

According to Koch (1962), Haacke (1976) and Visser (1984) *Pachydactylus* (formerly *Palmatogeecko*) *rangei* occurs from the mouth of the Orange River to southern Angola and never further inland than 160 km. They are common in the sand sea south of the Kuiseb River as well as the coastal dunes and ephemeral river beds in the Namib Desert (Haacke 1976, Stuart 1980, Russell & Bauer 1990), mostly associated with fine sand and rainfall below 125mm p.a. (Haacke 1976) and fog (Louw 1972, Seely & Griffin 1986). It is estimated that 80% of the potential proportion of the taxon's range falls within Namibia (Griffin 1985) and it is viewed as a Namib Desert endemic (Herrman & Branch 2012).