

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/289522647>

Namib research: its development at Gobabeb and implications for Namibia

Article · January 2000

CITATIONS
5

READS
84

4 authors, including:





Joh R. Henschel
South African Environmental Observation Network

179 PUBLICATIONS 4,027 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:

 **Project** Ecology of tenebrionid beetles in arid environments [View project](#)

 **Project** Spotted hyena biology [View project](#)

NAMIB RESEARCH: ITS DEVELOPMENT AT GOBABEB AND IMPLICATIONS FOR NAMIBIA

MARY K. SEELY, JOH R. HENSCHER, JULIANE ZEIDLER and
E. S. CHARLIE SHANYENGANA

Desert Research Foundation of Namibia, Gobabeb
P.O. Box 20232, Windhoek, e-mail: drfn@drfn.org.na

Abstract

The aridity, variability and biodiversity of the Namib Desert have attracted researchers for over a century. The character of that research started as biophysical exploration, shifted towards biodiversity and economic geology, followed the emphasis of descriptive geomorphology and ecological systems. More recently, the focus is on long-term ecological research, biodiversity processes, restoration ecology, resource management and other applied topics. Since its establishment in 1962, the Gobabeb Training and Research Centre and its predecessors have generated over 1300 publications on the Namib while a similar number have emanated from other institutions world-wide.

Much of the Namib research reflects the unusual character of the desert, its longevity, shape and position, extreme aridity, prevalence of fog and wind, massive dunes and its ephemeral rivers forming linear oases. Approaches have ranged from observation, to natural and contrived experiments, and laboratory analyses of Namib-derived materials. One strength of Namib research has been the frequent use of a variety of approaches by collaborating investigators focusing efforts on specific questions. The collective results of this have ensured that the Namib is one of the best known deserts in the world.

While many research themes continue to be investigated a decade after Namibian Independence, there has been a major shift towards applied research in support of development. Fog collection as an alternative water source and the productivity of the *!nara* plant are just two examples. Application of existing results for land-use and regional planning, tourism development, combating desertification, monitoring global climate change and for environmental impact assessments is another recent development. A third direction is the use of research results for education and training. The Namib is particularly suited for this latter application because of the existing understanding of the environment as well as the stark clarity of its environmental systems that are reflected in other, less arid, regions of Namibia. Although more than a century of research has been focused on the Namib, there are many questions left and applications to be realised. Namib research can continue to make a major contribution to the development of Namibia.

Introduction

Environmental research has made the Namib one of the best known hyperarid deserts of the world. Situated at the central point of the Namib, the Gobabeb Training and Research Centre has contributed significantly towards this status in terms of conducting, coordinating, or collating research, as well as training young researchers. After Independence, this relatively good foundation of knowledge has facilitated the development of capacity, skills and knowledge to manage arid lands throughout Namibia and elsewhere in Southern Africa. By supporting development in this way, Namib research has gained local and regional significance in addition to its international reputation.

The Desert Research Foundation of Namibia (DRFN) that grew out of the Desert Ecological Research Unit (DERU) has been responsible for much of the above-mentioned activities. Appropriate for this 75-year Festschrift of the Namib Scientific Society, the DRFN is currently half as old as the society. This paper shows the course of the development of the DRFN, the research background from which it grew, and how it relates to Namib, Namibia, as well as Desert research in general. Today the DRFN is a leading Namibian NGO (non-government organisation) engaged in research, training and information exchange on the biophysical and the human environments. Its foundation for these national and regional roles rests on Namib research as outlined in this paper.

Early Explorations

The aridity, climatic variability and biodiversity of the Namib Desert have attracted researchers with varying interests for over a century. The character of that research started, before the turn of the century, as biophysical exploration. Often this early research was based on written comments from early traders with respect to the surrounding biophysical and socio-economic environment during the course of their activities (e.g., THOMPSON, 1786; ALEXANDER, 1838; WELWITSCH, 1861; BAINES, 1864, 1866; PEARSON, 1907). Although these very first observations can not be considered research as it is known today, they provide the basis for much investigative research that has followed. Moreover, these early accounts served to make information concerning the Namib Desert accessible to a limited reading public that would never visit Namibia.

The next group of people contributing to the foundations of research in the Namib emerged as Germany's colony 'Deutsch-Südwest-Afrika' became established. These included economic geologists looking for precious stones and metals as well as farmers, soldiers and others who found their new surroundings too interesting to ignore (VON FRANCOIS, 1893; GRIMME, 1910; TRENK, 1910; REUNING, 1913; KAISER and BEETZ, 1919; KAISER, 1920), as well as the first scientists (STAPFF, 1887a, b, 1888; GULLAND, 1907; MARLOTH, 1909). Following the First World War and the transfer of Namibia into South Africa's control, the basis for further fieldwork and good documentation broadened (e.g., KORN and MARTIN, 1937). One of the special features of the Namib, abundant and diverse tenebrionid beetles, was elucidated by GEBIEN (1937, 1939). With the broad ecological perspectives published by WALTER (1936a, b, c), the Namib Desert gained world renown.

Founding of Gobabeb as a research centre

The history of the Namib Desert Research Station, the first of many names for the centre at Gobabeb, dates back to 1948. In that year, the University of California made an expedition to the south-western coast of Africa and invited as their entomologist the world renowned tenebrionid beetle expert, Dr. Charles Koch (KOCH, 1950). Dr Koch was very familiar with other arid areas of the world and in particular their invertebrate fauna. When he encountered the diversity of species and forms and habitats of the Namib Desert, he gave up his museum work in Europe and dedicated the remainder of his life to their study (KOCH, 1960, 1961a, b, 1962a, b). After exploring the Namib throughout its length, from Angola south to South Africa, he selected a site on the Kuiseb River in the central Namib for establishing a research centre (KOCH, 1959; LAWRENCE, 1960). Although chosen for different reasons at the time, this site at Gobabeb has proved to be excellent for the research and training which has taken place over the years. It is located at a point of easy access to the dunes, the ephemeral Kuiseb River and the gravel plains and at a place where rainfall diminishes towards the west and where fog from the west ceases to be a regular phenomenon. What better place to study and learn about the effects of habitat and climatic variability, and resultant population dynamics and biodiversity, the Leitmotif of the Namib and most of Namibia.

To generate interest, and funding for the research centre, many popular articles were written about the Namib and its biota. These articles ranged from an account of the big Kuiseb River flood of 1963 (KOCH, 1963) through accounts of the discovery of the elusive golden mole (HAACKE, 1963) and overall descriptions of the unusual sand dune fauna (LAWRENCE, 1959, 1965; FITZSIMONS, 1963). Accompanying these popular articles were the basic taxonomic descriptions, written for the specialist, of the various tenebrionid beetles and other organisms discovered and described by Dr. Koch and his co-workers (e.g., in 1962 new species were described of tenebrionid beetles: KOCH, 1962c; tenebrionid larvae SCHULZE, 1962a, b; grasshoppers BROWN, 1962; spiders, scorpions, solifugids, and centipedes: LAWRENCE, 1962a, 1962b; pseudoscorpions: BEIER, 1962; geckos: BRAIN, 1962a; lizards: FITZSIMONS, 1962). Both types of publications were essential to provide the research base for further work, to inform the scientific community about opportunities in the Namib and to share discoveries with the Namibian and international public.

Several years after the establishment of the Namib Desert Research Association at Gobabeb in 1962, as an independent research entity under the auspices of the Southern African Museums Association and the Transvaal Museum, research activities accelerated. The focus was on making species inventories in different habitats, particularly of tenebrionid beetles, but also of all the other unexpected fauna. What has continued to fascinate many scientists until today (>180 publications), is the high diversity of tenebrionid beetles in the Namib, despite the extremely low productivity (SEELY and LOUW, 1980). Over 200 kinds of tenebrionids occur in the Namib Desert (KOCH op cit., 1967; PENRITH, 1975, 1977, 1979, 1986), with over 80 species near Gobabeb, and up to 42 sympatric species occurring together in one habitat (HOLM and SCHOLZ, 1980; WHARTON and SEELY, 1982; CRAWFORD and SEELY, 1987; HENSCHL, SEELY and POLIS, 1998).

Why does the Namib have this rich faunal assemblage when neither the Sahara nor any other sand massifs have comparable fauna? How does this wide variety of plants and animals live in such a seemingly hostile environment? These questions captured the attention of a number of scientists using a variety of approaches through the years (e.g., BRAIN, 1960, 1962b; BRAIN and PROZESKY, 1962; COETZEE, 1969; HOLM, 1970; CHANNING, 1976; HOLM and SCHOLTZ, 1980; GRIFFIN, 1981; HAACKE, 1982; IRISH and MENDES, 1988; GRIFFIN, 1990). The remarkable *Welwitschia* garnered a lot of this attention, then and now (e.g., BORNMAN, 1978; VON WILLERT, 1985; VON WILLERT and WAGNER-DOUGLAS, 1994; HENSCHER and SEELY, in press), as 300 scientific papers indicate (HENSCHER *et al.*, in press).

Desert Ecological Research Unit (DERU)

By the 1970s, the Council for Scientific and Industrial Research of South Africa (CSIR) was funding a majority of the research of the Desert Ecological Research Unit (DERU) and the then Directorate of Nature Conservation had taken over the infrastructure at Gobabeb. Nature Conservation was responsible for issuing all research permits for the Namib Park, where Gobabeb was situated, and conducted some own research in the Namib, or, more commonly, in collaboration with DERU. Emphasis of Namib research focused on descriptive geomorphology and ecological systems. It was during this period, when Dr. Mary Seely became Director of DERU, that the approach to research and training for which Gobabeb is widely known was established (SEELY, 1972, 1975, 1978a, b, 1979a, b; DERU, 1976-1990).

DERU has always been based on two main pillars of operation. First has been the research of the DERU 'staff', very limited in number and often in the process of undertaking post-graduate or post-doctoral studies. In addition to their own work, this nucleus of people supports the research of a variety of visiting scientists at Gobabeb who either work on their own topic or, more often, in collaboration with DERU staff. Since its establishment, DERU and its successor, DRFN, have generated over 1300 publications on the Namib (HENSCHER *et al.*, 2000). Eighteen doctoral (cited in this paper) and 38 masters students have successfully completed their degrees under the auspices of DERU/DRFN.

A similar number of publications on the Namib has emanated from other institutions world-wide, including many Namibian institutions, for example the Ministry of Environment and Tourism, Department of Water Affairs, Geological Survey, National Museum of Namibia, Namibian National Botanical Research Institute, Society for Scientific Development, and the Namibia Scientific Society. Important among this research by other institutions are archaeological investigations that indicated that nomadic people were hunters and gatherers in the Namib Desert over the course of some ten millenia, becoming pastoralists during the past two millenia (e.g., SANDELOWSKY, 1976, 1977; JACOBSON, 1977, 1978, 1985; SHACKLEY, 1985; KINAHAN, 1991). With its high concentration of rock paintings, the Brandberg and nearby mountains attracted considerable international attention (e.g., PAGER, 1980; BREUNIG, 1989).

The responsibilities of the DERU staff, while focused on research, range more widely. The core staff, mainly degree students or post-docs, undertook their defined research projects. To enhance their research, however, most soon recognised that it was important to collaborate with and involve researchers from other institutions. Staff members were strongly encouraged to adopt this approach. In this way, Gobabeb-based researchers could undertake and support field investigations while relying on their collaborators in other institutions to provide the sophisticated instrumentation and perform sensitive analyses. Initiation of these collaborations was varied, from joining up with a visiting scientist while the researcher was in the Namib to seeking out persons and negotiating procedures for collaboration. In addition to enhancing research possibilities and their outputs, this approach served to multiply many-fold the always limited funding of the DERU.

The collaborative approach also served to enhance the publication output of the DERU and its associates. As researchers, the staff and visitors are dependent on publications in international peer-reviewed journals. In addition, however, publication in popular journals was encouraged and often pursued. This might be in the local journals for tourists as well as international conservation magazines. Another vehicle for popularisation of research is through films for a variety of audiences. DERU staff and visiting scientists often assisted with the story line and gathering actors for a variety of films, some of which have an intercontinental viewing audience. Yet another method of popularising research is the granting of interviews, often to overseas journalists gathering information about this, ever more famous, coastal desert. When necessary, the DERU staff could refer interested persons to their many collaborators scattered throughout the globe.

The collaborative approach extended to incorporate students on various levels from a variety of institutions into the overall research programme. Professors sent their students to carry out parts of their study programme in the Namib, and these people would be incorporated into the overall programme. Opportunities for summer vacation work were cultivated so that undergraduates could learn research techniques and, at the same time, contribute to ongoing research programmes. Soon the opportunities presented by Gobabeb were recognised in many institutions from which persons interested in field biology and geomorphology came to the Namib. Of note was that DERU maintained a high proportion of female students, who were generally excluded from fieldwork elsewhere in southern Africa. At the same time, collaborating colleagues identified Gobabeb as a suitable venue to hold field courses. University classes from southern Africa and overseas would spend several days or weeks taking advantage of the site, the information and the expertise provided by Gobabeb residents.

During this twenty year span, most of the current research directions were initiated (SEELY, 1990a). Climate is a prevailing theme for research in the long, narrow Namib desert with its steep west-east gradient of precipitation encompassing winter rainfall in the south and summer in the north. Since 1962, weather observations have been made at the First Order Weather Station at Gobabeb and reported thrice daily to the Namibian Meteorological Services (e.g., SEELY and STUART, 1976; LANCASTER, LANCASTER and SEELY, 1984). These long term records, augmented by shorter term records from elsewhere in the Namib document the episodic events and variability of climate that

characterise all of arid Namibia. Analyses of the dynamics of the desert climate facilitated broader understanding in southern Africa (TYSON and SEELY, 1980; TYSON, 1986; LINDESAY and TYSON, 1990). Namib climate was also the key for elucidating geomorphological processes, and this connection allowed analyses of palaeoclimate to be made (SCHOLZ, 1963; BESLER, 1972; GOUDIE, 1972; BLÜMEL, 1976; HARMSE, 1980; VOGEL, 1981; WARD, SEELY and LANCASTER, 1983; WARD, 1984, 1988a, b; LIVINGSTONE, 1985; LANCASTER, 1989; EICHLER, 1990; HEINE, 1990, 1998; WARD and CORBETT, 1990; WALTER, 1994; ECKARDT, 1996).

Winds that shift surface material about blow for >50% of the time (LANCASTER, *et al.* 1984), shaping the landscape and the ecosystem. Wind continuously redistributes dead or dormant plant material and causes it to be buried and remobilised, thereby driving the slow, steady availability of detritus to detritivores (CRAWFORD and SEELY, 1994; HANRAHAN and KIRCHNER, 1997). It is therefore not surprising that detritivores, such as tenebrionid beetles, ants, termites, fish moths, and gerbils, are major consumers in the Namib (SEELY and LOUW, 1980; LOUW and SEELY, 1982; CURTIS, 1983; SEELY, 1983; CRAWFORD and TAYLOR, 1984; MARSH, 1985a; WEHNER, 1987; CRAWFORD and SEELY, 1987; WATSON, 1989; HANRAHAN and SEELY, 1990). Much of the research therefore concerned the functioning of these small animals and their consumers in such extremely arid conditions (e.g., WILLOUGHBY, 1967; COUTCHIÉ and CROWE, 1979; HADLEY and LOUW, 1980; WITHERS, LOUW and HENSCHÉL, 1980; DE VILLIERS, 1984; MCCLAIN *et al.*, 1985; SEELY and GRIFFIN, 1986; DOWNS, 1989; FIELDEN, 1989; HENSCHÉL, 1990; NICOLSON, 1990; ROBERTS, 1991; CLARKE, 1992; NAIDU, 1992; GRUBE, 1993). There is often little food, and tolerance of hunger is important (e.g., SEELY and LOUW, 1980; NAGY, SEELY and BUFFENSTEIN, 1993; HENSCHÉL, 1994; LUBIN and HENSCHÉL, 1996; SEYMOUR, WITHERS and WEATHERS, 1998). This is compounded by daily hot surface conditions and extreme daily temperature fluctuations (e.g., HENWOOD, 1974; HAMILTON, 1975; CURTIS, 1985; MARSH, 1985b, 1987; SEELY and MITCHELL, 1987; PIETRUSZKA, 1988; LOMBARD, 1989; CLOUDSLEY-THOMPSON, 1990; LUBIN and HENSCHÉL, 1990; ROBINSON, 1990, 1993; WARD, 1990; ROBERTS, 1991). Studies of the ecophysiology of arthropods and small vertebrates in the Namib established world prominence in this field during the 1970s and 1980s (MALOY, 1972; HADLEY, 1974; EDNEY, 1977; LOUW and SEELY, 1982; LOUW, 1990, 1993; SEELY, 1990a; CLOUDSLEY-THOMPSON, 1991; HEINRICH, 1993; LOVEGROVE, 1993; SOMME, 1995; HEATWOLE, 1996; BRADSHAW, 1997; DEGEN, 1997).

Studies of the spatial and temporal effects of variable climate on biotic communities of the desert contributed to understanding of functioning of natural and man-made ecosystems throughout Namibia and other arid regions of the world. A most important contribution of Namib research was the relationship that was established between productivity, biodiversity and climate. Fog has long been identified as a key component of the Namib (e.g., WALTER, 1936; BOSS, 1941; NAGEL, 1962) and is predictable (PIETRUSZKA and SEELY, 1985). Its precipitated water increases survival of plants and animals (HAMILTON and SEELY, 1976a, b; SEELY and HAMILTON, 1976; SEELY, DE VOS and LOUW, 1977; SEELY, 1979c; LOUW and SEELY, 1980; NOTT, 1985; LORIS, 1990; POLIS and

SEELY, 1990; HENSCHHEL, 1997a; HENSCHHEL, SEELY and POLIS, 1998; SEELY, HENSCHHEL and ROBERTSON, 1998). Fog explains the extraordinary growth and abundance of lichens in the Namib (SCHIEFERSTEIN and LORIS, 1992).

By contrast, rainfall is unpredictable to the degree that the occurrence of effective rain (affecting plant growth) cannot be predicted for any year (PIETRUSZKA and SEELY, 1985) and is rare. Nevertheless, temporal patterns of rainfall are of fundamental importance to the growth, abundance and diversity of Namib plants and animals. While fog enables many populations to sustain long periods without rain, effective rain is necessary for production (e.g., HOLM, 1970; SEELY, 1973, 1989, 1990b, 1991; SEELY and LOUW, 1980; NEL, 1983; HAMILTON, 1985, 1986; YEATON, 1988; BOYER, 1989; BERRY and SIEGFRIED, 1991; GÜNSTER, 1992, 1993; SOUTHGATE, MASTERS and SEELY, 1996; HENSCHHEL, SEELY and POLIS, 1998; HENSCHHEL and SEELY, in press). Distribution of rainfall affects spatial patterns within the ecosystem (ROBINSON, 1976; SEELY, 1978c, d; TILSON and HENSCHHEL, 1986; GÜNSTER, 1992, 1995; BRAIN, 1993; KILIAN, 1995; KOK and NEL, 1996; BURKE, 1997). Although big rainfall events of >100 mm per annum have occurred only three times during the 20th century in the western Namib, they are believed to be of fundamental importance for the Namib ecosystem. Such rains elicit extremely strong, rapid increases in productivity of many plants and animals, effects that last for decades (WALTER, 1971; SEELY and LOUW, 1980; SOUTHGATE, MASTERS and SEELY, 1996; HENSCHHEL, SEELY and POLIS, 1998).

In order to better understand the mechanisms, implications and applications of climatic variability, DERU initiated Long-term Ecological Research (LTER) at Gobabeb from its inception. Continuous monitoring grew partly out of short-term studies of microclimate, ecophysiology and behaviour (SEELY, 1975, 1979b, 1990a; SEELY and WARD, 1988), and continued to be an important activity after independence (SEELY and SGUAZZIN, 1992; SEELY *et al.*, 1999a; HENSCHHEL, 1999). The Gobabeb LTER program now encompasses collaborative projects on climate, ephemeral and perennial plants, tenebrionids, spiders, dune movements, weathering, ephemeral river dynamics and natural resource use (HENSCHHEL, SEELY and ZEIDLER, this volume).

Desert Research Foundation of Namibia (DRFN) at Gobabeb

At Namibian Independence, the controlling and funding functions of the CSIR and Transvaal Museum over DERU ceased (SEELY, 1990c). The umbrella organisation that now incorporated DERU was named Desert Research Foundation of Namibia (DRFN). This non-government organisation (NGO) became an independent research and training institution. DRFN transformed the knowledge and understanding gained in the Namib in the previous thirty years and extended it country-wide, particularly focusing on ways of applying it to development involving the environment (SEELY, 1990d; SEELY *et al.*, 2000). In the meantime, training and research in the Namib continued, with DRFN continuing to be based at Gobabeb besides its new offices in Windhoek and Oshakati and a temporary office in Khorixas (DRFN, 1990-2000).

In the early 1990s, Gobabeb housed the rapidly expanding DRFN where new directions were formalised. The Enviroteach project, funded by the Swedish International Development Agency (Sida) developed an innovative approach to cross-curricular, activity-based, learner-centred environmental education in co-operation with the Ministry of Basic Education and Culture. The 'old lab' was transformed into a book production centre and titles ranging from energy, to water, population and environmental education methodologies were compiled (DU TOIT *et al.*, 1995a-j). These were tested in 25 pilot schools throughout the country and, when the project moved to Windhoek after four hectic years, emphasis shifted to the four Colleges of Education.

An income generating arm of the DRFN was established under the name of Environmental Evaluation Associates of Namibia (Pty) Ltd (EEAN) which used the knowledge and understanding gained at Gobabeb to undertake environmental assessments throughout Namibia. Previous research and current assessments were incorporated into books for decision makers and awareness materials for extension personnel, students and decision-makers (MARSH and SEELY, 1992; Update series, DRFN 1997-2000). The ephemeral rivers project epitomises the approach pioneered at Gobabeb where scientific research (e.g. HUNTLEY, 1985; JACOBSON *et al.*, 1999; SEELY *et al.*, 1999b) is applied to sustainable development in Namibia (book, video, map by JACOBSON, JACOBSON and SEELY, 1995a, b). This project, and its successor project that concentrated on the Hoanib catchment (LEGGETT, 2000), was funded by Sida and carried out in close collaboration with the Department of Water Affairs, with whom a number of other projects were also undertaken with funds from various sources. These projects include information on water sharing in southern Africa (PALLETT, 1997), the decision-maker's guide to water use in Namibia (HEYNS *et al.*, 1998), and regional water books (FORBES-IRVING and WARD, 1999; WARD and FORBES-IRVING, 1999).

As circumstances forced the move of most of the DRFN staff to Windhoek, the stage was set at Gobabeb for future directions. While continuing to incorporate the ongoing activities at Gobabeb, the DRFN formed partnerships with a variety of projects, institutions and Ministries. The DRFN's participation in Namibia's Programme to Combat Desertification originated at Gobabeb (SEELY and JACOBSON, 1994; SEELY, 1998; ZEIDLER, 2000) as did various projects taking place throughout Namibia. Gobabeb is now particularly known for the Summer Desertification Project (Sida funded) wherein young Namibians from the University and Polytechnic are taught skills of environmental problem solving, including planning, field and analytical methods, presenting and publishing (DEALIE *et al.*, 1993; DAUSAB *et al.*, 1994; JOBST *et al.*, 1995; GUIDAO-OAB *et al.*, 1996; JOBST *et al.*, 1997; HAININGA *et al.*, 1998; GAZZA *et al.*, 1999; AMOOMO *et al.*, 2000).

At Gobabeb, graduates from the University of Namibia and Polytechnicon can gain hands-on experience in the environmental field (HENSCHERL, 1997b). This facilitates the ability of such students to obtain higher degrees at foreign universities or relevant employment. The DRFN offers intensive week-long courses, intern courses of 6-12 months, and coaches postgraduate studies. Recently, the "Tropenökologisches Begleitprogramm" (TÖB) of the "Gesellschaft für Technische Zusammenarbeit"

(GTZ), has served as a good example of how students become integrated as in-service trainees (HAMUKWAYA, 1998), or post-graduates (PARENZEE, 2000), or staff (ZEIDLER, 2000). Where appropriate, the DRFN at Gobabeb also involves visiting scientists as trainees. The situation of Gobabeb with its "outdoor classroom" and relative isolation from urban distractions makes it an ideal training centre that can contribute substantially to the goal of a well-trained cadre of Namibians.

Besides continuing basic research (some recent activities have been described with previous activities in the above section on DERU), a shift in the approach to Namib research has been to provide immediate relevance to development in Namibia (SEELY, 1998; SEELY, ZEIDLER and PARENZEE, 2000; SEELY *et al.*, in press). Both quantitative analytical and qualitative approaches are used to assess environmental conditions and to monitor and evaluate resource use in the Namib. People become directly involved in the research process, rather than being research subjects only, as was the case previously. The resulting information is not only prepared for scientific audiences, but is fed back by appropriate means to the people concerned, and is transmitted to decision-makers. Examples are projects on the development of an Index of Biological Integrity (ZEIDLER *et al.*, 1998; ZEIDLER, 1999; PARENZEE, 2000; ZEIDLER, SEELY and PARENZEE, 2000; PARENZEE, ZEIDLER and SEELY, 2000), the connection between resource management and socio-politics (SULLIVAN, 1998), and environmental consequences of communal and commercial farming (WARD *et al.*, 1999). Similar approaches are used in the interaction with the Topnaar community living at and near Gobabeb, who have requested the DRFN to become involved in resource management, such as the use of water (DAUSAB, 1994; HENSCHER *et al.*, 1998; MTULENI, HENSCHER and SEELY, 1998; AMOOMO *et al.*, 2000) and !nara fruit (several separate studies on plant biology, GOLDBERG and IWANEK, 2000; information exchange concerning harvesting and marketing, SHILOMBOLENI, 1998; BÜTTENDORF, 2000). Other projects concern Gobabeb itself, such as using it as a model in developing and applying appropriate technology in arid lands (SHANYENGANA, 1997a, b). The latter project is partly based on the opportunities for research and testing of equipment at Gobabeb, and partly out of the need to change resource use at Gobabeb with its ageing infrastructure and currently limited water and energy resources.

The Ministry of Environment and Tourism (MET) also continues to conduct important research in the Namib, including research on mammals and reptiles (KILIAN, 1995; GRIFFIN, 1997; ongoing aerial censuses), coastal birds (BRABY, BRABY and SIMMONS, 1992; SIMMONS 1996; SIMMONS, CORDES and BRABY, 1998; ongoing wetland censuses), and the conservation of lichen fields (BRABY, 1990; ongoing monitoring). MET has been instrumental in bringing relevant information together to assist decision-making for development in the region (BARNARD, 1998; BENDER, 1999). Based on these and further internal studies, as well as interactions with other organisations and people of the Namib, the MET is adapting the management of its Namib parks to current conditions and needs. In this way, the Namib-Naukluft Park, the Skeleton Coast Park, and the new Walvis Bay Park incorporating the Walvis Bay lagoon and its surroundings, are benefiting from the current relatively good status of knowledge of the Namib.

In 1998, the MET and DRFN signed a Joint Venture Agreement forming a Board of Trustees for the Gobabeb Training and Research Centre (GTRC). The Board of Trustees leased the facilities of the GTRC for 20 years, assuring its continued operation. Part of DRFN continues to be based at the GTRC, and continues to play an active role through its ongoing training and research activities. The GTRC is a natural meeting place for students, scientists, community members, government institutions, and other decision makers, who contribute to and receive information and experiences of sustainable natural resource management from the ongoing programmes based at Gobabeb (SEELY, 2000). The DRFN at Gobabeb has developed an active rapport with indigenous people in Namibia. Positive interaction is ensured by active participation and ongoing consultation by all parties involved. This approach is proving to be a powerful means of transforming knowledge and understanding gained in research towards enhancing the capacity of Namibians towards sustainable management of their natural resources.

In the year 2000, the DRFN at Gobabeb focuses on long-term ecological research (LTER), biodiversity processes, restoration ecology, resource management and other applied topics. The Namibian capacity in environmental research is actively furthered through courses to students, intern programmes, learning opportunities for staff, as well as facilitating post-graduate studies by Namibians. Examples of novel or topical research directions that grew out of a wealth of previous knowledge generated at or near Gobabeb are:

- * small-scale low-technology solar distillation experiments that incorporate findings dating back to the beginning of this century
- * collecting of fog water by using a principle first identified for Namib animals and plants 25 years ago
- * enhancing the ability of the Topnaars to manage utilisation of !nara products, based on extensive information exchange as well as participatory research in socio-economic and cultural fields, and ecological and ecophysiological research that focus on the plant's needs and its importance to its environment
- * improving natural resource management along the ephemeral rivers of western Namibia, such as the Kuiseb, by forming partnerships with information networks, resource users and decision makers, and thereby increasing our collective ability to contribute to and use environmental knowledge
- * monitoring and understanding extreme environmental variability of the hyperarid Namib with the Gobabeb Long Term Ecological Research Programme in synergy with related national and international research networks so as to facilitate the application of this understanding to other variable environments to reduce the latter's susceptibility to degradation and improve sustainable livelihoods (HENSCHER, SEELY and ZEIDLER, this volume)
- * interlinking Namib research of ecology, natural resource management and socio-economics in close collaboration with the Namibian Economic Policy Research Unit (Nepu) as a model for Namibia
- * contributing actively towards networking between various government and non-government organisations, so as to improve information exchange, developing supportive partnerships, inte-

gration of research and information into decision making and policy formulation, as well as developing the appropriate methodologies for current research and related functions, for example via the internet-based Netwise (<http://www.netwise.drfn.org.na/>).

The shift of research emphasis has echoed not only the growing understanding of the Namib environment but also advances in international science as well as the needs of Namibia's development process. DRFN faces the future by building on its rich foundation, by identifying opportunities for fruitful outreach, and addressing the challenges given by Namibia's need for environmental information and improved capacity to wisely manage natural resources in variable, arid environments.

Conclusion

Much of Namib research reflects the unusual character of the desert, its longevity, its shape and position, its extreme aridity, the prevalence of fog and wind, its ephemeral rivers forming linear oases and its massive dunes derived from sandstones across the continent. Approaches have ranged from simple observation, to experiments, and laboratory analyses of Namib-derived materials. One strength of Namib research has been the use of a variety of approaches by collaborating investigators focusing efforts on specific questions. The collective results of this extensive, frequently co-operative research have ensured that the Namib is one of the best known deserts in the world.

While many research themes continue to be investigated a decade after Namibian Independence, there has been a major shift toward applied research in support of development. Application of existing results for land-use and regional planning, for tourism development, for combating desertification, for monitoring global climate change and for environmental impact assessments is another recent development. A third fresh direction is the use of research results for education and training. The Namib is particularly suited for this latter application because of the existing understanding of the environment as well as the stark clarity of its environmental systems that are reflected in other, less arid, regions of Namibia. Although more than a century of research has been focused on the Namib, there are many questions to be asked and applications to be realised. The contribution of the Namib to the development of Namibia is a long way from being exhausted.

Acknowledgements

The Ministry of Environment and Tourism gave permission for research to be conducted in the Namib-Naukluft Park. We thank numerous donors for their generosity, which has enabled the Gobabeb Training and Research Centre to continue. In particular, Sida, GTZ, Norad, USAID, British Council and many other donors for their contribution toward the ongoing programmes of the DRFN. Over the years, many people have assisted diligently to ensure the functioning of the centre; currently these include ANNA BEES, DAWID DAMASEB, INGE HENSCHER, ED HOGG, ALEX KHURISAB, VILHO MTULENI, JAKOB MAKEMBE, SHARON MONTGOMERY, RAUNA NAWA, LESLEY PARENZEE, CHRIS PARK, DEON SHARURU and the current authors. SHARON MONTGOMERY and MARK ROBERTSON kindly read previous drafts of this manuscript.

REFERENCES

- ALEXANDER, J.E. 1838. *An expedition of discovery into the interior of Africa*, 2 volumes. Henry Colburn, London.
- AMOOMO, H., GASEB, N., HOVEKA, V., MBANGULA, E., MUHARUKUA, E. and ZAARUKA, B. 2000. *Determining the water reserve of the Kuiseb River - sustainable use of Namibia's ephemeral rivers, life-lines for development*. Desert Research Foundation of Namibia, Summer Desertification Programme 8.
- BAINES, T. 1864. *Explorations in South-West Africa*. London: Longman, Roberts & Green. 535 pp.
- BAINES, T. 1866. *The Welwitschia mirabilis* (plant of Hykamkop), South West Africa. *Nature and Art*, August 1: 69-74. (reprinted in 1972 in *Excelsa* 2: 41-49).
- BARNARD, P. (Ed.) 1998. *Biological diversity in Namibia - a country study*. Windhoek, Namibia: Namibian National Biodiversity Task Force, Directorate of Environmental Affairs. 325 pp.
- BEIER, M. 1962. Pseudoscorpioniden aus der Namib-Wüste. *Annals of the Transvaal Museum* 24: 223-230.
- BENDER, K. (Ed.) 1999. *Coastal profile of the Erongo Region*. Swakopmund, Namibia: Integrated Coastal Zone Management Project, Erongo Regional Council. 214 pp.
- BERRY, H.H. and SIEGFRIED, W.R. 1991. Mosaic-like events in arid and semi-arid Namibia. In: *The mosaic-cycle concept of ecosystems*. Remmert, H. (Ed). Berlin: Springer. pp.147-160.
- BESLER, H. 1972. Klimaverhältnisse und klimageomorphologische Zonierung der zentralen Namib (Südwestafrika). *Stuttgarter Geographische Studien* 83: 1-209.
- BLÜMEL, W.D. 1976. Kalkkrustenvorkommen in Südwestafrika. *Mitteilungen der Basler Afrika Bibliographien* 15: 17-50.
- BORNMAN, C.H. 1978. *Welwitschia: paradox of a parched paradise*. Cape Town: C. Struik Publishers.
- BOSS, G. 1941. Niederschlagsmenge und Salzgehalt des Nebelwassers an der Küste Deutsch-Südwestafrikas. *Bioklimatische Beiblätter der Meteorologischen Zeitschrift* 8(1): 1-14.
- BOYER, D.C. 1989. Some characteristics of the plant communities of three dunes situated across a climatic gradient in the Namib Desert. *Madoqua* 16: 141-148.
- BRABY, R. 1990. Off road vehicles in Desert Environments; minimizing the impact and rehabilitating the damage of vehicle tracks in the Namib. In: *Environmental Implications of Prospecting and Mining in the Namib*, DANEEL, J. (ED.). Geological Society of Namibia and Desert Ecological Research Unit of Namibia. pp. 14-17.
- BRABY, R., BRABY, S.J. and SIMMONS, R.E. 1992. 5000 Damara Terns in the northern Namib Desert: a reassessment of world population numbers. *Ostrich* 63: 133-135.
- BRADSHAW, S.D. 1997. *Homeostasis in desert reptiles*. Berlin: Springer Verlag. 213 pp.
- BRAIN, C.K. 1960. Observations on the locomotion of the South West African adder, *Bitis peringueyi* (Boulenger), with speculation on the origin of sidewinding. *Annals of the Transvaal Museum* 24: 19-24.
- BRAIN, C.K. 1962a. A review of the gecko *Ptenopus* with the description of a new species. *Cimbebasia*: 11-18.
- BRAIN, C.K. 1962b. Observations on the temperature tolerance of lizards in the central Namib Desert, South West Africa. *Cimbebasia* 4: 1-5.

- BRAIN, C.K. and PROZESKY, O.P.M. 1962. On the body temperatures of birds collected on the Carp-Transvaal Museum Namib Desert expedition. *Cimbebasia* 5: 13-20.
- BRAIN, C. 1993. *The eco-physiology of baboons living in the Kuiseb River Canyon*, Namibia. Ph.D. thesis, University of the Witwatersrand, Johannesburg. 198 pp.
- BREUNIG, P. 1989. Archaeological investigations into the settlement history of the Brandberg. In: *The rock paintings of the Upper Brandberg I: Amis Gorge*. PAGER, H. (Ed.). Heinrich Barth Institut, Köln: 17-33.
- BROWN, H.D. 1962. New species of the genus *Acocksacris* (Orthoptera: Acridoidea). *Annals of the Transvaal Museum* 24: 181-196.
- BURKE, A. 1997. Is the grass always greener at certain sites? A contribution to patch dynamics in the central Namib desert. *Tropical Ecology* 38: 125-127.
- BÜTTENDORF, U. 2000. The value of the !nara. *Namib Bulletin* 15: 25.
- CHANNING, A. 1976. Life histories of frogs in the Namib Desert. *Zoologica Africana* 11: 299-312.
- CLARKE, B.C. 1992. *Water and energy balance in the Namib sand dune lizard, Anglosaurus skoogi*. PhD thesis, University of Cape Town, Cape Town. 211 pp.
- CLOUDSLEY-THOMPSON, J.L. 1990. Thermal ecology and behaviour of *Physadesmia globosa* (Coleoptera: Tenebrionidae) in the Namib Desert. *Journal of Arid Environments* 19: 317-324.
- CLOUDSLEY-THOMPSON, J.L. 1991. *Ecophysiology of desert arthropods and reptiles*. Berlin: Springer-Verlag. 203 pp.
- COETZEE, C.G. 1969. The distribution of mammals in the Namib Desert and adjoining inland escarpment. *Scientific Papers of the Namib Desert Research Station* 4: 23-36.
- COUTCHIÉ, P.A. and CROWE, J.H. 1979. Transport of water vapor by tenebrionid beetles II. Regulation of the osmolarity and composition of the hemolymph. *Physiological Zoology* 52: 88-100.
- CRAWFORD, C.S. and SEELY, M.K. 1987. Assemblages of surface-active arthropods in the Namib dunefield and associated habitats. *Revue de Zoologie Africaine* 101: 397-421.
- CRAWFORD, C.S. and SEELY, M.K. 1994. Detritus mass loss in the Namib Desert dunefield: influence of termites, gerbils and exposure to surface conditions. *Journal of African Zoology* 108: 49-54.
- CRAWFORD, C.S., TAYLOR, E.C. 1984. Decomposition in arid environments: the role of the detritivore gut. *South African Journal of Science* 80: 170-176.
- CURTIS, B.A. 1983. *Behavioural and physiological ecology of the Namib Desert dune ant, Camponotus detritus Emery*. MSc thesis, University of Cape Town. 212 pp.
- CURTIS, B.A. 1985. Temperature tolerances in the Namib desert dune ant, *Camponotus detritus*. *Journal of Insect Physiology* 31: 463-466.
- DAUSAB, F., FRANCIS, G., JÖHR, G., KAMBATUKU, J., MOLAPO, M., SHANYENGANA, E. and SWARTZ, S. 1994. *Water usage patterns in the Kuiseb catchment area - with emphasis on sustainable use*. Desert Research Foundation of Namibia, Summer Desertification Programme 2, Occasional Paper 1. 243 pp.
- DEALIE, F.F., HAMATA, S.A., KAMBATUKU, J.R., MOLAPO, M., PARENZEE, L.G. and SOROSES, F.W. 1993. *Rainfall in Namibia: What is normal?* Desert Research Foundation of Namibia, Gobabeb, Summer Desertification Programme 1. 13 pp.

- DEGEN, A.A. 1997. *Ecophysiology of small desert mammals*. Berlin: Springer Verlag. 296 pp.
- DE VILLIERS, P.S. 1984. *Aspects of the reproductive behaviour of the Namib tenebrionid beetle, Onymacris unguicularis* (Haag). PhD thesis, Faculty of Science, University of the Witwatersrand, Johannesburg. 185 pp.
- D.E.R.U. 1976-1990. *Namib Bulletin* volumes 1-9.
- DOWNS, C.T. 1989. *The ecophysiology of four Gerbillurus species with special reference to their temperature and water regulation*. PhD thesis, University of Natal, Pietermaritzburg, South Africa. 210 pp.
- D.R.F.N. 1990-2000. *Namib Bulletin* volumes 10-15.
- D.R.F.N. 1997-2000. *Update: a regular briefing on aspects of sustainable development in Namibia*. Desert Research Foundation of Namibia, Windhoek, Volumes 1-4 (56 numbers).
- DU TOIT, D., KARITA, A. and SGUAZZIN, T. 1995a. *Lives in the balance - people and the Namibian environment*. Windhoek, Namibia: Desert Research Foundation of Namibia. 221 pp.
- DU TOIT, D., KARITA, A. and SGUAZZIN, T. 1995b. *A beginner's guide to outdoor teaching*. Windhoek, Namibia: Desert Research Foundation of Namibia. 23 pp.
- DU TOIT, D., KARITA, A. and SGUAZZIN, T. 1995c. *Fun with the sun - activities to demonstrate solar power*. Windhoek, Namibia: Desert Research Foundation of Namibia. 34 pp.
- DU TOIT, D., KARITA, A. and SGUAZZIN, T. 1995d. *Wholly ground*. Windhoek, Namibia: Desert Research Foundation of Namibia. 39 pp.
- DU TOIT, D., KARITA, A. and SGUAZZIN, T. 1995e. *My land my life*. Windhoek, Namibia: Desert Research Foundation of Namibia. 115 pp.
- DU TOIT, D., KARITA, A. and SGUAZZIN, T. 1995f. *Tools of the trade - skills and techniques for environmental education in Namibia*. Windhoek, Namibia: Desert Research Foundation of Namibia. 199 pp.
- DU TOIT, D., KARITA, A. and SGUAZZIN, T. 1995g. *Sink or swim...water and the Namibian environment*. Windhoek, Namibia: Desert Research Foundation of Namibia. 226 pp.
- DU TOIT, D., KARITA, A. and SGUAZZIN, T., BURKE, A. and MANNHEIMER, C. 1995h. *Let's look at seaweeds*. Windhoek, Namibia: Desert Research Foundation of Namibia. 98 pp.
- DU TOIT, D., KARITA, A. and SGUAZZIN, T., GÜNSTER, A. and MANNHEIMER, C. 1995i. *Our growing wealth*. Windhoek, Namibia: Desert Research Foundation of Namibia. 85 pp.
- DU TOIT, D. and SGUAZZIN, T. 1995j. *Playing with fire - energy and the Namibian environment*. Windhoek, Namibia: Desert Research Foundation of Namibia. 204 pp.
- ECKARDT, F.D. 1996. *The distribution and origin of gypsum in the Central Namib Desert, Namibia*. PhD thesis, University of Oxford, Hertford College, Oxford. 320 pp.
- EDNEY, E.B. 1977. *Water balance in land arthropods*. Berlin: Springer-Verlag. 282 pp.
- EICHLER, H. 1990. Von Wasser und Wind in der zentralen Namib (Namibia) - Bemerkungen zum wenig beachteten Feuchtehaushalt in den Dünensanden einer Nebelwüste. *Mainzer Geographische Studien* 34: 61-76.
- FIELDEN, L.J. 1989. *Selected aspects of the adaptive biology and ecology of the Namib Desert golden mole (Eremitalpa granti namibensis)*. PhD thesis, University of Natal, Dept. of Zoology and Entomology, Pietermaritzburg, South Africa. 247 pp.

- FITZSIMONS, V. 1962. A new worm snake (*Leptotyphlops*) from South West Africa. *Annals of the Transvaal Museum* 24: 239-240.
- FITZSIMONS, V. 1963. The Namib Desert. *African Wildlife* 17: 215-227.
- FORBES-IRVING, T. and WARD, V. 1999. *Managing water points and grazing areas in Namibia: Cuvelai, Northern Kunene, Erongo*. (3 volumes in English & Oshiwambo) Windhoek, Namibia: Desert Research Foundation of Namibia. 100 pp. each.
- GAZZA, A. LUBANDA, P., MAHESHE, L., MUHARUKUA, V., NAWA, R., NICKANOR, N., SHIMALI, P. and SHIPONENI, N. 1999. *Development options for resettlement in northeast Otjozondjupa*. Desert Research Foundation of Namibia, Summer Desertification Programme 7, Occasional Paper 9. 93 pp.
- GEBIEN, H. 1937. Die *Tenebrioniden* (*Coleoptera Heteromera*) der Namibwüste in Südwestafrika. 7: 20-107.
- GEBIEN, H. 1939. Körperbau und Lebensweise der Wüstentenebrioniden. *International Kongress der Entomologie*, Berlin: 118-132.
- GOLDBERG, K. and IWANEK, S. 2000. The !nara - (Natural Resource for the #Aonin) project. *Namib Bulletin* 15: 24.
- GOUDIE, A.S. 1972. Climate, weathering, crust formation, dunes, and fluvial features of the central Namib desert, near Gobabeb, South West Africa. *Madoqua, Series II* 1(54-62): 15-31.
- GRIFFIN, M. 1981. *Contributions to the Kuiseb Environmental Program: Namib game population studies*. Windhoek: Department of Agriculture and Nature Conservation. pp. 1-20.
- GRIFFIN, M. 1997. *Checklist and provisional national conservation status of amphibians, reptiles, and mammals known or expected to occur in Namibia*. Memorandum, Ministry of Environment and Tourism, Windhoek.
- GRIFFIN, R.E. 1990. Seasonal activity, habitat selection and species richness of Solifugae (Arachnida) on the gravel plains of the central Namib desert. In: *Namib ecology: 25 years of Namib research*. SEELY, M.K. (Ed). Transvaal Museum Monograph 7 Transvaal Museum, Pretoria. pp. 77-82.
- GRIMME, C. 1910. 'Narras', ein wichtiges Eingeborenen-Nahrungsmittel in Deutsch-Südwestafrika. *Der Tropenpflanzer Zeitschrift für tropische Landwirtschaft* 14: 297-302.
- GRUBE, S. 1993. *Untersuchungen zum Wassergleichgewicht bei Termiten in ariden Lebensräumen am Beispiel von Psammotermes allocerus Silvestri*. Diplom Thesis, Freie Universität Berlin, Berlin, Germany. 69 pp.
- GUIDAO-OAB, J., HUMAVINDU, M., KAKUKURU, E., NGAIRORUE, T., NGHISHOONGELE, I.O., NGHITILA, T.M. and NGOLOLO, E.N. 1996. *Interactions between people and elephants as one component of desertification in the Kunene Region*. Desert Research Foundation of Namibia, Summer Desertification Programme 4. 73 pp.
- GULLAND, A. 1907. Aus dem deutsch-südwestafrikanischen Schutzgebiete. Das Klima von Swakopmund. *Litt. dtsh. Schutzgeb.* 20(131): 17-164.
- GÜNSTER, A. 1992. Aerial seed banks in the central Namib: distribution of serotinous plants in relation to climate and habitat. *Journal of Biogeography* 19: 563-572.
- GÜNSTER, A. 1993. *Seed bank dynamics of serotinous plants in the central Namib*. Ph.D. thesis, University of Münster, Münster, Germany. 103 pp.

- GÜNSTER, A. 1995. Grass cover distribution in the central Namib - a rapid method to assess regional and local rainfall patterns of arid regions? *Journal of Arid Environments* 29: 107-114.
- HAACKE, W.D. 1963. First find of the Namib Golden Mole (*Eremitalpa granti namibensis*). IUCN Bulletin 9: 8.
- HAACKE, W.D. 1982. Distribution patterns of reptiles as indicators of geomorphological events in the Namib desert. In: *Palaeoecology of Africa* 15. COETZEE, J.A., VAN ZINDEREN BAKKER, E.M.Sr. (Eds). Rotterdam: A.A.Balkema. pp. 225-226.
- HADLEY, N.F. 1974. *Environmental physiology of desert organisms*. Stroudsburg: Halted Press. 283 pp.
- HADLEY, N.F. and LOUW, G.N. 1980. Cuticular hydrocarbons and evaporative water loss in two tenebrionid beetles from the Namib Desert. *South African Journal of Science* 76: 298-301.
- HAININGA, S., HASHIPALA, M., HEYMAN, I., MUFITA, V., NGHILEBENDELE, P., NHULEIPO, O., PARENZEE, L. and RÖHL, C. 1998. *Environmental and resource economic implications of fencing in the communal areas of eastern Oshikoto*. Desert Research Foundation of Namibia, Windhoek, Summer Desertification Programme 6. 58 pp.
- HAMILTON III, W.J. 1975. Coloration and its thermal consequences for diurnal desert insects. In: *Environmental Physiology of Desert Organisms*. HADLEY, N.F. (Ed.). USA: Halstead Press. pp. 67-89.
- HAMILTON III, W.J. 1985. Demographic consequences of a food and water shortage to desert Chacma baboons, *Papio ursinus*. *International Journal of Primatology* 6: 451-462.
- HAMILTON III, W.J. 1986. Namib Desert Chacma baboon (*Papio ursinus*) use of food and water resources during a food shortage. *Madoqua* 14: 397-408.
- HAMILTON III, W.J. and SEELY, M.K. 1976a. Fog basking by the Namib Desert beetle, *Onymacris unguicularis*. *Nature* 262(5566): 284-285.
- HAMILTON III, W.J. and SEELY, M.K. 1976b. Fog trap construction and fog basking by Namib Desert tenebrionid beetles. *Bulletin of the Ecological Society of America* 57: 57.
- HAMUKWAYA, P.N. 1998. *Historical and present land management and land-uses at three selected farms in southern Kunene region*. Third year Diploma in-service training thesis, Department of Natural Resource Management of the Polytechnic of Namibia & Desert Research Foundation of Namibia, Windhoek. 37 pp.
- HANRAHAN, S.A. and KIRCHNER, W.H. 1997. The effect of wind on foraging activity of the tenebrionid beetle *Lepidochora discoidalis* in the sand dunes of the Namib Desert. *South African Journal of Zoology* 32: 136-139.
- HANRAHAN, S.A. and SEELY, M.K. 1990. Food and habitat use by three tenebrionid beetles (Coleoptera) in a riparian desert environment. In: *Namib ecology: 25 years of Namib research*. SEELY, M.K. (Ed.). Transvaal Museum Monograph 7, Transvaal Museum, Pretoria. pp. 143-147.
- HARMSE, J.T. 1980. *Die noordwaartse begrensing van die duinsee van die sentrale Namib langs die Benede-Kuiseb*. Ph.D thesis, University of Stellenbosch, Stellenbosch, South Africa. 229 pp.
- HEATWOLE, H. 1996. *Energetics of desert invertebrates*. Berlin: Springer-Verlag. 266 pp.
- HEINE, K. 1990. Klimaschwankungen und klimagenetische Geomorphologie am Beispiel der Namib. *Berliner Geographische Studien* 30: 221-234.

- HEINE, K. 1998. Late Quaternary climate changes in the Central Namib Desert, Namibia. In: *Quaternary Deserts and Climatic Change*. ALSHARHAN, A.S., GLENNIE, K.W., WHITTLE, G.L., and KENDALL, C.G.S. (Eds.). Rotterdam: AA Balkema. pp. 293-304.
- HEINRICH, B. 1993. *The hot-blooded insects*. Berlin: Springer. 601 pp.
- HENSCHHEL, J.R. 1990. Spiders wheel to escape. *South African Journal of Science* 86: 151-152.
- HENSCHHEL, J.R. 1994. Diet and foraging behaviour of huntsman spiders in the Namib dunes (Araneae: Heteropodidae). *Journal of Zoology*, London 234: 239-251.
- HENSCHHEL, J.R. 1997a. Psammophily in Namib Desert spiders. *Journal of Arid Environments* 37: 695-707.
- HENSCHHEL, J.R. 1997b. International research cooperation with the Desert Research Foundation of Namibia (DRFN) benefits the natural environment in Namibia. In: *Wissenschaftler Austausch und Entwicklungszusammenarbeit vor der Jahrtausendwende Bestandsaufnahme, Probleme und Perspektiven*. Lehnert, G. and Duggan, J. (Eds.). Bonn-Bad Godesberg, Germany: Alexander von Humboldt-Stiftung. pp. 401-406.
- HENSCHHEL, J.R. 1999. Namibia moving towards networking LTER. *Abstract, Annual General Meeting of the International Long Term Ecological Research Network, August 1999, Skukuza, South Africa*.
- HENSCHHEL, J.R., BÖHME, E., DENTLINGER, L., HENSCHHEL, I.A., MÜLLER, M. and SEELY, M.K. 2000. DRFN *Bibliography 2000*. Windhoek: Desert Research Foundation of Namibia. 86 pp.
- HENSCHHEL, J.R., ELLER, B.M., SEELY, M.K. and VON WILLERT, D.J. in press. Bibliography *Welwitschia mirabilis* J.D.Hooker. *Plant Ecology*: in press.
- HENSCHHEL, J.R., MTULENI, V., GRUNTKOWSKI, N., SEELY, M.K. and SHANYENGANA, S.E. 1998. *Namfog: Namibian Application of Fog-Collecting Systems, Phase I: Evaluation of Fog-Water Harvesting*. Windhoek: Desert Research Foundation of Namibia, Occasional Paper No. 8. 81pp.
- HENSCHHEL, J.R. and SEELY, M.K. in press. Long-term growth patterns of *Welwitschia mirabilis*, a long-lived plant of the Namib Desert (including a bibliography). *Plant Ecology*: in press.
- HENSCHHEL, J.R., SEELY, M.K. and POLIS, G.A. 1998. Long term population dynamics of tenebrionid beetles in the Namib Desert. *Abstract, INTECOL: VII International Congress of Ecology, July 1998, Florence, Italy*.
- HENSCHHEL, J.R., SEELY, M.K. and ZEIDLER, J. 2000. Long-term ecological research in western Namibia identifies fundamental processes underlying biodiversity. *Namibia Scientific Society Journal Volume 48*.
- HENWOOD, K. 1974. I. *A field tested thermoregulation model for two diurnal desert tenebrionid beetles*. II. *Infrared relectivity as an alternative strategy to colour change*. Ph.D. thesis, University of California, Davis, USA. 39 pp.
- HEYNS, P., MONTGOMERY, S., PALLETT, J. and SEELY, M. (Eds.) 1998. *Namibia's water: a decision makers' guide*. Windhoek, Namibia: The Department of Water Affairs, Ministry of Agriculture, Water and Rural Development, and The Desert Research Foundation of Namibia. 173 pp.

- HOLM, E. 1970. *The influence of climate on the activity patterns and abundance of xerophilous Namib desert dune insects*. M.Sc. thesis, University of Pretoria, Pretoria. 44 pp.
- HOLM, E. and SCHOLTZ, C.H. 1980. Structure and pattern of the Namib Desert dune ecosystem at Gobabeb. *Madoqua* 12: 3-39.
- HUNTLEY, B.J. (Ed.). 1985. *The Kuiseb environment: the development of a monitoring baseline*. South African National Scientific Programmes Report 106. 138 pp.
- IRISH, J. and MENDES, L.F. 1988. New genera and species of ultraspammophilous Namib Desert Lepismatidae (Thysanura). *Madoqua* 15: 275-284.
- JACOBSON, L. 1977. Namib oriented archaeological research at the State Museum, Windhoek. *Namib Bulletin* 2: 9.
- JACOBSON, L. 1978. A Brandberg bibliography. S.W.A. *Wissenschaftliche Gesellschaft Supplement* 18(10-11): 1-4.
- JACOBSON, L. 1985. Review on: recent archaeological research between the Orange and Kavango rivers in south-western Africa. *South African Archaeological Bulletin* 40(142): 115-116.
- JACOBSON, P.J., JACOBSON, K.M., ANGERMEIER, P.L. and CHERRY, D.S. 1999. Transport, retention, and ecological significance of woody debris within a large ephemeral river. *Journal of the North American Benthological Society* 18: 429-444.
- JACOBSON, P.J., JACOBSON, K.M. and SEELY, M.K. (Eds.) 1995a. *Ephemeral rivers and their catchments: sustaining people and development in western Namibia*. Windhoek, Namibia: Desert Research Foundation of Namibia. 167 pp.
- JACOBSON, P.J., JACOBSON, K.M. and SEELY, M.K. (Eds.). 1995b. Video: *Lifelines of Namibia: Ephemeral rivers project*. Cape Town: Doxa Production.
- JOBST, P., KAKUKURU, E., KAMBATUKU, J.R., KAMWI, J., MOUTON, D., NETHA, L., NGHITILA, T.M. and SHANYENGANA, S.E. 1995. *Indicators of desertification in north-western Namibia, alternative land uses, and associated socio-economic and policy issues*. Desert Research Foundation of Namibia, Summer Desertification Programme 3, Occasional Paper 2. 93 pp.
- JOBST, P., NGHITILA, T., MAHERERO, J.H., MUFITA, V. and NGULEIPO, O. 1997. *Environmental economics and the influence of infrastructure on land use in the Oshana and Oshikoto Regions*. Desert Research Foundation of Namibia, Windhoek, Summer Desertification Programme 5.
- KAISER, E. and BEETZ, W. 1919. Die Wassererschließung in der südlichen Namib Südwestafrikas. *Zeitschrift für praktische Geologie* 11/12: 165-186.
- KAISER, E. 1920. Morphogenetische Ergebnisse auf Reisen während des Krieges in Südwestafrika. *Abhandlungen der Giessener Hochschulgesellschaft* 2: 159-175.
- KILIAN, J.W. 1995. *The ecology of gemsbok (Oryx gazella gazella) in the southern Namib*. Windhoek: Ministry of Environment and Tourism.
- KINAHAN, J. 1991. *Pastoral Nomads of the central Namib Desert: the people history forgot*. Windhoek: Namibia Archaeological Trust. 167 pp.
- KOCH, C. 1950. The Tenebrionidae of Southern Africa 1: first account of the Tenebrionidae collected on the University of California-Transvaal Museum expedition, 1948. *Annals of the Transvaal Museum* 21(3): 273-367.

- KOCH, C. 1959. Report on Transvaal Museum expedition to Namib Desert May 1959. *Memorandum, Transvaal Museum*: 1-2.
- KOCH, C. 1960. The tenebrionid beetles of South West Africa. *South African Museums Association Bulletin* 7(4): 73-85.
- KOCH, C. 1961a. Die Dünentenebrioniden der Namibwüste mit besonderer Berücksichtigung ihrer Ökologie (und Physiologie). *XI Internationaler Kongress für Entomologie, Vienna*: 655-657.
- KOCH, C. 1961b. Some aspects of abundant life in the vegetationless sand of the Namib Desert dunes. *Journal of the South West African Scientific Society* 15: 9-34.
- KOCH, C. 1962a. Zur Ökologie der Dünen-Tenebrioniden der Namibwüste Südwest-Afrikas und Angolas. *Lunds Universitets Arsskrift N.F.* 58(5): 3-25.
- KOCH, C. 1962b. The Tenebrionidae of southern Africa. 31: Comprehensive notes on the tenebrionid fauna of the Namib Desert. *Annals of the Transvaal Museum* 24(2/3): 61-106.
- KOCH, C. 1962c. The Tenebrionidae of southern Africa. 32: New psammophilous species from the Namib Desert. *Annals of the Transvaal Museum* 24(2/3): 107-159.
- KOCH, C. 1963. An illustrated account of a major flood in the Kuiseb River. *Der Kreis* 2/3: 39-51.
- KOCH, C. 1967. Beetles from the Namib Desert. *African Wildlife* 21(4): 305-312.
- KOK, O.B. and NEL, J.A.J. 1996. The Kuiseb river as a linear oasis in the Namib desert. *African Journal of Ecology* 34: 39-47.
- KORN, H. and MARTIN, H. 1937. Die jüngere geologische und klimatische Geschichte Südwestafrikas. *Zentralblatt für Mineralogie* 11: 456-473.
- LANCASTER, N. 1989. *The Namib sand sea*. Rotterdam: AA Balkema. 179 pp.
- LANCASTER, J., LANCASTER, N. and SEELY, M.K. 1984. Climate of the central Namib Desert. *Madoqua* 14: 5-61.
- LAWRENCE, R.F. 1959. The sand-dune fauna of the Namib Desert. *South African Journal of Science* 55: 233-239.
- LAWRENCE, R.F. 1960. The Namib Desert biological station. *South African Journal of Science* 56(4): 90.
- LAWRENCE, R.F. 1962a. Spiders of the Namib Desert. *Annals of the Transvaal Museum* 24: 197-212.
- LAWRENCE, R.F. 1962. Solifuges, scorpions and Chilopoda of the Namib Desert. *Annals of the Transvaal Museum* 24: 213-222.
- LAWRENCE, R.F. 1965. Dune spiders of the Namib Desert. *Animals* 7(10): 260-263.
- LEGGETT, K. 2000. The Hoanib River Catchment study. *Namib Bulletin* 15: 22.
- LINDESAY, J.A. and TYSON, P.D. 1990. Thermo-topographically induced boundary layer oscillations over the central Namib, Southern Africa. *International Journal of Climatology* 10: 63-77.
- LIVINGSTONE, I. 1985. *The dynamics of sand transport on a Namib linear dune*. Ph.D. thesis, University of Oxford, Oxford. 318 pp.
- LOMBARD, A.T. 1989. *The thermal implications and ecological consequences of coloration in selected species: Tenebrionid beetles (Onymacris bicolor and Onymacris unguicularis), Cape gannets (Morus capensis) and Cape cormorants (Phalacrocorax)*. Ph.D. thesis, University of Cape Town, Cape Town. 205 pp.

- LORIS, K. 1990. Zur Problematik der Wasserversorgung von *Arthraerua leubnitziae* in der Äusseren Namib, mit Daten des Nebelniederschlags. *Symposium zum Gedenken an Prof. Dr. h.c. H. Walter, Stuttgart-Hohenheim, University of Stuttgart-Hohenheim*: 1-15.
- LOUW, G.N. 1990. Physiological studies on the Namib fauna: a brief critique. In: *Namib ecology: 25 years of Namib research*. SEELY, M.K. (Ed.). Transvaal Museum Monograph 7, Transvaal Museum, Pretoria. pp. 203-207.
- LOUW, G. 1993. *Physiological animal ecology*. London: Longman. 281 pp.
- LOUW, G.N. and SEELY, M.K. 1980. Exploitation of fog water by a perennial Namib dune grass, *Stipagrostis sabulicola*. *South African Journal of Science* 76: 38-39.
- LOUW, G. and SEELY, M.K. 1982. *Ecology of desert organisms*. London: Longman. 184 pp.
- LOVEGROVE, B. 1993. *The living deserts of Southern Africa*. Cape Town: Fernwood Press. 224 pp.
- LUBIN, Y.D. and HENSCHHEL, J.R. 1990. Foraging at the thermal limit: burrowing spiders (Seothyra, Eresidae) in the Namib desert dunes. *Oecologia* 84: 461-467.
- LUBIN, Y.D. and HENSCHHEL, J.R. 1996. The influence of food supply on foraging behaviour in a desert spider. *Oecologia* 105: 64-73.
- MALOIJ, G.M.O. 1972. *Comparative physiology of desert animals*. London: Zoological Society of London. 413 pp.
- MARLOTH, R. 1909. The vegetation of the southern Namib. *South African Journal of Science* 6: 80-87.
- MARSH, A.C. 1985a. *Aspects of the ecology of Namib desert ants*. Ph.D. thesis, University of Cape Town, Cape Town. 254 pp.
- MARSH, A.C. 1985b. Microclimatic factors influencing foraging patterns and success of the thermophilic desert ant, *Ocymyrmex barbiger*. *Insectes Sociaux* 32: 286-296.
- MARSH, A.C. 1987. Thermal responses and temperature tolerance of a desert ant-lion larva. *Journal of Thermal Biology* 12: 295-300.
- MARSH, A. and SEELY, M.K. (Eds.) 1992. *Oshanas: Sustaining people, environment, and development in Central Owambo, Namibia*. Windhoek: Desert Research Foundation of Namibia. 55 pp.
- MCCLAIN, E., SEELY, M.K., HADLEY, N.F. and GRAY, V. 1985. Wax blooms in tenebrionid beetles of the Namib Desert: correlations with environment. *Ecology* 66: 112-119.
- MTULENI, V., HENSCHHEL, J.R. and SEELY, M.K. 1998. Evaluation of fog-harvesting potential in Namibia. In: *Proceedings: 1st International Conference on Fog and Fog Collection*, Vancouver, Canada. SCHEMENAUER, R.S. and BRIDGMAN, H. (Eds.). pp. 179-182.
- NAGEL, J.F. 1962. Fog precipitation measurements on Africa's Southwest Coast. *South African Weather Bureau, Notos* 11: 51-60.
- NAGY, K.A., SEELY, M.K. and BUFFENSTEIN, R. 1993. Surprisingly low field metabolic rate of a diurnal desert gecko, *Rhoptropus afer*. *Copeia* 3: 216-219.
- NAIDU, S.G. 1992. *Osmoregulation in tenebrionid beetles endemic to the Namib Desert*. Ph.D. thesis, University of Witwatersrand, Johannesburg. 215 pp.
- NEL, P.S. 1983. *Monitoring van die beskikbaarheid, gehalte en benutting van voer op die gruisflaktes van die Kuiseb-studiegebied*. M.Sc. thesis, University of the Orange Free State, Bloemfontein, South Africa. 233 pp.

- NICOLSON, S.W. 1990. Water relations of the Namib tenebrionid beetles In: *Namib ecology: 25 years of Namib research*. SEELY, M.K. (Ed). Transvaal Museum Monograph 7 Transvaal Museum, Pretoria. pp. 173-178.
- NOTT, K. 1985. *An autecological study of a Namib dune succulent, Trianthema hereroensis*. M.Sc. thesis, University of Natal, Pietermaritzburg, South Africa. 147 pp.
- PAGER, H. 1980. Felsbildforschung am Brandberg in Namibia: Beiträge, allgemeine und vergleichende Archäologie. *Deutsches Archäologisches Institut, Bonn* 2: 351-357.
- PALLET, J. (Ed.) 1997: *Sharing water in southern Africa*. Windhoek: Desert Research Foundation of Namibia. 121 pp.
- PARENZEE, L. 2000. *Use of tenebrionid beetles as indicators of habitat quality*. M.Sc. thesis, University of Witwatersrand, Johannesburg.
- PARENZEE, L., ZEIDLER, J. and SEELY, M. 2000. Testing biodiversity indicators for community use - a case study from Namibia. *Abstract, 13th Congress of the German Society for Tropical Ecology, March 2000, Würzburg, Germany*. p. 37.
- PEARSON, H.H.W. 1907. Some notes on a journey from Walfish Bay to Windhuk. *Bulletin of Miscellaneous Information, Royal Botanic Gardens, Kew* 1907: 339-360.
- PENRITH, M-L. 1975. The species of *Onymacris* Allard (Coleoptera: Tenebrionidae). *Cimbebasia, Series A* 4(3): 47-97.
- PENRITH, M-L. 1977. The Zophosini (Coleoptera: Tenebrionidae) of western southern Africa. *Cimbebasia Memoir*: 1-291.
- PENRITH, M-L. 1979. Revision of the western southern African Adesmiini (Coleoptera: Tenebrionidae). *Cimbebasia, Series A* 5(1): 1-94.
- PENRITH, M-L. 1986. Relationships in the tribe Adesmiini (Coleoptera: Tenebrionidae) and a revision of the genus *Stenodesia* Reitter. *Annals of the Transvaal Museum* 34(13): 275-302.
- PIETRUSZKA, R.D. 1988. Maxithermy and the thermal biology of an herbivorous sand dune lizard. *Journal of Arid Environments* 14: 175-185.
- PIETRUSZKA, R.D. and SEELY, M.K. 1985. Predictability of two moisture sources in the Namib Desert. *South African Journal of Science* 81: 682-685.
- POLIS, G.A. and SEELY, M.K. 1990. Imbibition of precipitated fog by Namib Desert scorpions. *Journal of Arachnology* 18: 362-363.
- REUNING, E. 1913. Eine Reise längs der Küste Lüderitzbucht-Swakopmund im Februar-März 1912. *Mitteilungen aus den Deutschen Schutzgebieten* 26: 118-126.
- ROBERTS, C.S. 1991. *The surface and subsurface environments and the physiological and behavioural ecology of dune-living beetles in the Namib Desert*. M.Sc. thesis, University of the Witwatersrand, Johannesburg. 122 pp.
- ROBINSON, E.R. 1976. *Phytosociology of the Namib Desert Park, South West Africa*. M.Sc. thesis, University of Natal, Pietermaritzburg, South Africa.
- ROBINSON, M.D. 1990. Summer field energetics of the Namib desert dune lizard *Aporosaura anchietae* (Lacertidae), and its relation to reproduction. *Journal of Arid Environments* 18: 207-216.
- ROBINSON, M.D. 1993. Death and dancing on the sun-baked dunes of Namibia. *Natural History* 102(8): 28-30.

- SANDELOWSKY, B.H. 1976. The beginning of archaeo-ethno-botany in the Namib. In: *Palaeoecology of Africa* 9. VAN ZINDEREN BAKKER, E.M. (Ed). Rotterdam: A.A.Balkema. pp. 136-143.
- SANDELOWSKY, B.H. 1977. Mirabib - an archaeological study in the Namib. *Madoqua* 10: 221-283.
- SCHIEFERSTEIN, B. and LORIS, K. 1992. Ecological investigation on lichen fields of the central Namib, I. Distribution patterns and habitat conditions. *Vegetatio* 98: 113-128.
- SCHOLZ, H. 1963. *Studien über die Bodenbildung zwischen Rehoboth und Walvis Bay*. D.Sc. thesis, Rheinische Friedrich-Wilhelm Universität, Bonn. 184 pp.
- SCHULZE, L. 1962a. The Tenebrionidae of Southern Africa. 33: Descriptive notes on the early stages of *Onymacris rugatipennis* Haag and *Lepidochora discoidalis* Gebien and keys to genera and species of Adesmiini and Eurychorini. *Annals of the Transvaal Museum* 24: 161-180.
- SCHULZE, L. 1962b. The Tenebrionidae of Southern Africa. 33: Description of the larvae of *Gonopus tibialis* Fabricius and *Gonopus agrestis* Fahraeus (*Gonopina*, sensu Koch 1956). *Cimbebasia* 5: 1-12.
- SHACKLEY, M. 1985. Palaeolithic archaeology of the central Namib Desert: a preliminary survey of chronology, typology and site location. *Cimbebasia Memoir* 6: 1-84.
- SHANYENGANA, E.S. 1997a. Baseline study: Gobabeb Training and Research Centre programme: appropriate technology for arid lands. Windhoek: Desert Research Foundation of Namibia. 56 pp.
- SHANYENGANA, E.S. 1997b. Appropriate technology at Gobabeb: a look into the future. *Namib Bulletin* 14: 18.
- SHILOMBOLENI, A. 1998. *The !nara and factors that lead to its decline in productivity*. Third year Diploma in-service training thesis, Department of Natural Resource Management of the Polytechnic of Namibia & Desert Research Foundation of Namibia, Windhoek. 17 pp.
- SEELY, M.K. 1972. Namib research. *South West African Annual* 1972: 169-176.
- SEELY, M.K. 1973. Factors controlling reproduction of certain Namib Desert tenebrionids. *Madoqua, Series II* 2(63-68): 63-65.
- SEELY, M.K. 1975. *Dr.-Charles-Koch-Namib-Wüsten-Forschungsstiftung der Namib-Wüsten-Forschungsstation*. Windhoek: John Meinert. 6 pp.
- SEELY, M.K. 1978a. Desert Ecological Research Unit. *Scientiae* 19(1): 37-39.
- SEELY, M.K. 1978b. Desert research. *South African Journal of Science* 74: 196.
- SEELY, M.K. 1978c. Grassland productivity: the desert end of the curve. *South African Journal of Science* 74: 295-297.
- SEELY, M.K. 1978d. Standing crop as an index of precipitation in the central Namib grassland. *Madoqua* 11: 61-68.
- SEELY, M.K. 1979a. Desert Ecological Research Unit. *Education and Culture* 2(6): 5-22.
- SEELY, M.K. 1979b. Ecology of a living desert: twenty years of research in the Namib. *South African Journal of Science* 75: 298-303.
- SEELY, M.K. 1979c. Irregular fog as a water source for desert dune beetles. *Oecologia* 42: 213-227.
- SEELY, M.K. 1983. Unsung heroes of the desert: detritivores. *Rössing Magazine*: 1-5.
- SEELY, M.K. 1989. Desert invertebrate physiological ecology: is anything special? *South African Journal of Science* 85: 266-270.

- SEELY, M.K. (Ed.). 1990a. *Namib ecology: 25 years of Namib research*. Transvaal Museum Monograph 7, Transvaal Museum, Pretoria. 223 pp.
- SEELY, M.K. 1990b. Patterns of plant establishment on a linear desert dune. *Israel Journal of Botany* 39: 443-451.
- SEELY, M.K. 1990c. DERU after independence: focus on education, funding. *Namib Bulletin* 10: 1.
- SEELY, M.K. 1990d. Ecology in Namibia at independence. *Intecol Newsletter* 20(2): 1-3.
- SEELY, M. 1991. *Namibia: drought and desertification*. Windhoek: Gamsberg Macmillan. 20 pp.
- SEELY, M.K. 1998. Can science and community action connect to combat desertification? *Journal of Arid Environments* 39: 267-277.
- SEELY, M.K. 2000. Gobabeb Training and Research Centre. *Namib Bulletin* 15: 10.
- SEELY, M.K., DE VOS, M.P. and LOUW, G.N. 1977. Fog imbibition, satellite fauna and unusual leaf structure in a Namib Desert dune plant, *Trianthema hereroensis*. *South African Journal of Science* 73: 169-172.
- SEELY, M.K. and GRIFFIN, M. 1986. Animals of the Namib Desert: interactions with their physical environment. *Revue de Zoologie Africaine* 100: 47-63.
- SEELY, M.K. and HAMILTON III, W.J. 1976. Fog catchment sand trenches constructed by tenebrionid beetles, *Lepidochora*, from the Namib Desert. *Science* 193(4252): 484-486.
- SEELY, M.K., HENSCHER, J.R., ABRAMS, M., JACOBSON, J. and ZEIDLER, J. 1999a. Long-term research in western Namibia identifies fundamental environmental processes. *Conference of the Ecological Society of America*, August 1999, USA.
- SEELY, M.K., HENSCHER, J.R. and ROBERTSON, M. 1998. The ecology of fog in Namib sand dunes. In: *Proceedings: 1st International Conference on Fog and Fog Collection, Vancouver, Canada*. SCHEMENAUER, R.S. and BRIDGMAN, H. (Eds.). pp. 183-186.
- SEELY, M.K. and JACOBSON, K.M. 1994. Desertification and Namibia: a perspective. *Journal of African Zoology* 108: 21-36.
- SEELY, M.K., JACOBSON, K.M., JACOBSON, P.J., LEGGETT, K. and NGHITILA, T. 1999b. Understanding integrated natural resource management at a catchment scale: the case of Namibia's ephemeral rivers. *Abstract, VIth International Rangelands Conference, Townsville, Australia*.
- SEELY, M.K. and LOUW, G.N. 1980. First approximation of the effects of rainfall on the ecology and energetics of a Namib Desert dune ecosystem. *Journal of Arid Environments* 3: 25-54.
- SEELY, M.K. and MITCHELL, D. 1987. Is the subsurface environment of the Namib Desert dunes a thermal haven for chthonic beetles? *South African Journal of Zoology* 22: 57-61.
- SEELY, M.K. and SGUAZZIN, T.M. 1992. Gobabeb: arid zone research in Namibia. *Journal of the Namibian Scientific Society* 43: 133-145.
- SEELY, M.K. and STUART, P. 1976. Namib climate: 2. The climate of Gobabeb, ten year summary 1962/72. *Namib Bulletin* 1: 7-9.
- SEELY, M.K. and WARD, J.D. 1988. The Namib Desert. In: *Long-Term data series relating to southern Africa's renewable natural resources*. MCDONALD, I.A.W. and CRAWFORD, R.J.M. (Eds.). South African National Scientific Programmes Report No. 157, Pretoria. pp. 268-279.
- SEELY, M.K., ZEIDLER, J., HENSCHER, J.R. and BARNARD, P. in press. Creative problem solving in support of biodiversity conservation. *Journal of Arid Environments*: in press.

- SEELY, M.K., ZEIDLER, J. and PARENZEE, L. 2000. Connecting science and community action. *Abstract, 13th Congress of the German Society for Tropical Ecology, March 2000, Würzburg, Germany.* p. 45.
- SEYMOUR, R.S., WITHERS, P.C. and WEATHERS, W.W. 1998. Energetics of burrowing, running, and free-living in the Namib Desert golden mole (*Eremitalpa namibensis*). *Journal of Zoology, London* 244: 107-117.
- SIMMONS, R.E. 1996. Population declines, viable breeding areas and management options for flamingos in southern Africa. *Conservation Biology* 10: 504-514.
- SIMMONS, R.E., CORDES, I. and BRABY, R. 1998. Latitudinal trends, population size and habitat preferences of Damara terns *Sterna balaenarum* on Namibia's desert coast. *Ibis* 140: 439-445.
- SOMME, L. 1995. *Invertebrates in hot and cold arid environments*. Berlin: Springer-Verlag. 265 pp.
- SOUTHGATE, R.I., MASTERS, P. and SEELY, M.K. 1996. Precipitation and biomass changes in the Namib Desert dune ecosystem. *Journal of Arid Environments* 33: 267-280.
- STAPFF, F.M. 1887a: Notiz über das Klima von Walfischbai. *Deutsche Kolonialzeitung* 4: 646-654.
- STAPFF, F.M. 1887b. Das untere Khuseibthal und sein Strandgebiet. *Gesellschaft für Erdkunde zu Berlin*: 45-66.
- STAPFF, F.M. 1888. Bodentemperaturbeobachtungen im Hinterlande der Walfischbay. *Sitzungsberichte der Mathematisch-Naturwissenschaftlichen Classe der kaiserlichen Akademie der Wissenschaften* 97: 119-141.
- SULLIVAN, S. 1998. *People, plants and practice in drylands: socio-political and ecological dimensions of resource-use by Damara farmers in north-west Namibia*. Ph.D. thesis, University College London, London. 448 pp.
- THOMPSON, T.B. 1786. Narrative of a voyage performed in His Majesty's Sloop Nautilus, under my command, for the purpose of investigating the western coast of South Africa, from latitude 33°00'00"S to 16°00'00" South. Kew Public Record Office Reference No. Adm 55/92. In: *The impenetrable shield: HMS Nautilus and the Namib coast in the late eighteenth century*, J. Kinahan (Ed.). *Cimbebasia* 12: 23-61.
- TILSON, R.L. and HENSCHER, J.R. 1986. Spatial arrangement of spotted hyaena groups in a desert environment, Namibia. *African Journal of Ecology* 24: 173-180.
- TRENK, W. 1910. In den Dünen der Namib. *Deutsches Kolonialblatt* 1910: 230-236.
- TYSON, P.D. 1986. *Climatic change and variability in southern Africa*. Oxford University Press, Cape Town. 217 pp.
- TYSON, P.D. and SEELY, M.K. 1980. Local winds over the central Namib. *South African Geographical Journal* 62: 135-150.
- VOGEL, J.C. 1981. Ancient climates and the age of the Namib. *South African Journal of Science* 77: 435-436.
- VON FRANCOIS, C. 1893. Das Küstengebiet zwischen Tsaokhaubmundung und Kap Cross. *Mitteilungen aus den Deutschen Schutzgebieten* (Berlin) 6: 299-302.
- VON WILLERT, D.J. 1985. *Welwitschia mirabilis* - new aspects in the biology of an old plant. *Advances in Botanical Research* 11:157-191.
- VON WILLERT, D.J. and WAGNER-DOUGLAS, U. 1994. Water relations, CO₂ exchange, water use efficiency and growth of *Welwitschia mirabilis* Hook. fil. in three contrasting habitats of the Namib Desert. *Botanica Acta* 107: 291-299.

- WALTER, H. 1936a. *Die ökologischen Verhältnisse in der Namib-Nebelwüste (Südwestafrika)*. Gebrüder Borntraeger, Leipzig. 222 pp.
- WALTER, H. 1936b. Die ökologischen Verhältnisse in der Nebelwüste Namib (Deutsch-Südwestafrika). *Berichte der Deutschen Botanischen Gesellschaft* 54: 39-44.
- WALTER, H. 1936c. Die Vegetationsverhältnisse in der Namib-Nebelwüste (Deutsch Südwestafrika). *Forschungen und Fortschritte (Berlin)* 12(24): 293-294.
- WALTER, H. 1971. *Ecology of tropical and subtropical vegetation*. Edinburgh: Oliver & Boyd. 539 pp.
- WALTER, R. 1994. *Beziehungen zwischen Krustenböden (Gips, Kalk) und der Morphologie der Flächennamib (Relationships between soil crusts (gypsum, calcium carbonate) and the morphology of the Namib plains)*. Diplom thesis, Institut für Geographie, Lehrstuhl für Physische Geographie, Universität Regensburg, Regensburg, Germany. 84 pp.
- WARD, D.M. 1990. A test of the 'maxithermy' hypothesis with three species of tenebrionid beetles. *Journal of Arid Environments* 21: 331-336.
- WARD, D., NGAIRORUE, B.T., KATHENA, J., SAMUELS, R. and OFRAN, Y. 1999. Land degradation is not a necessary outcome of communal pastoralism in arid Namibia. *Journal of Arid Environments* 40: 357-371.
- WARD, J.D. 1984. *Aspects of the Cenozoic geology in the Kuiseb Valley, central Namib Desert*. Ph.D. thesis, University of Natal, Durban, South Africa. 310 pp.
- WARD, J.D. 1988a. Eolian, fluvial and pan (playa) facies of the tertiary Tsondab Sandstone formation in the central Namib Desert, Namibia. *Sedimentary Geology* 55: 143-162.
- WARD, J.D. 1988b. On an interpretation of the Oswater conglomerate formation, Kuiseb Valley, Namib Desert. *Palaeoecology of Africa* 19: 119-125.
- WARD, J.D. and CORBETT, I. 1990. Towards an age for the Namib. In: *Namib ecology: 25 years of research*. SEELY, M.K. (Ed.). Transvaal Museum Monograph 7, Transvaal Museum, Pretoria. pp. 17-26.
- WARD, J.D., SEELY, M.K. and LANCASTER, N. 1983. On the antiquity of the Namib. *South African Journal of Science* 79: 175-183.
- WARD, V. and FORBES-IRVING, T. 1999. *Managing water points and grazing areas in Namibia: Caprivi, Southern Kunene, Omaheke, Karas & Hardap*. (4 volumes in English & Oshiwambo) Windhoek: Desert Research Foundation of Namibia. 100 pp. each.
- WATSON, R.T. 1989. Niche separation in Namib Desert dune Lepismatidae (Thysanura: Insecta): detritivores in an allochthonous detritus ecosystem. *Journal of Arid Environments* 17: 37-48.
- WEHNER, R. 1987. Spatial organization of foraging behavior in individually searching desert ants, *Cataglyphis* (Sahara Desert) and *Ocymyrmex* (Namib Desert). In: *Behavior in Social Insects*. PASTEELS, J.M. and DENEUBOURG, J.-L. (Eds). Basel: Birkhäuser Verlag. pp. 15-42.
- WELWITSCH, F. 1861. Extract from a letter addressed to Sir William J. Hooker on the botany of Benguela, Mossamedes etc. in Western Africa. *Journal of the Proceedings of the Linnean Society, Botany* 5: 182-187.
- WHARTON, R.A. and SEELY, M.K. 1982. Species composition of and biological notes on Tenebrionidae of the lower Kuiseb River and adjacent gravel plain. *Madoqua* 13: 5-25.

- WILLOUGHBY, E.J. 1967. *Desert adaptations of larks of the central Namib*. Ph.D. thesis, University Microfilms, London, England. 152 pp.
- WITHERS, P.C., LOUW, G.N. and HENSCHER, J.R. 1980. Energetics and water relations in Namib Desert rodents. *South African Journal of Zoology* 15: 131-137.
- YEATON, R.I. 1988. Structure and function of the Namib dune grasslands: Characteristics of the environmental gradients and species distributions. *Journal of Ecology* 76: 744-758.
- ZEIDLER, J. 1999. *Establishing indicators of biological integrity in western Namibian rangelands*. Ph.D. thesis, University of Witwatersrand, Johannesburg.
- ZEIDLER, J. 2000. Linking the environment to rural development - Napcod III. *Namib Bulletin* 15: 9.
- ZEIDLER, J., SEELY, M., HANRAHAN, S. and SCHOLES, M. 1998. Establishing indicators of biological integrity in western Namibian rangelands: combining an ecological with a participatory community-based research approach. *Abstract, Proceedings of the Kalahari Transect Meeting, Gaborone, Botswana, 10-13 June 1998, Start report No.2*.
- ZEIDLER, J., SEELY, M. and PARENZEE, L. 2000. Environmental indicators for community management. *Abstract, 13th Congress of the German Society for Tropical Ecology, March 2000, Würzburg, Germany*. p. 53.



Biography

Dr. MARY SEELY received her Ph.D. at the University of California, Davis, and a D.Sc. Honoris causa from the University of Natal, Pietermaritzburg. She is honorary professor at the University of Namibia and the University of Witwatersrand, and adjunct professor at the University of New Mexico and U.C.Davis. In 1970, she became director of the Desert Ecological Research Unit and in 1990 Executive Director of the Desert Research Foundation of Namibia. Apart from publishing 160 papers and 9 books, Dr. Seely has played a leading role in connecting science to development in Namibia.



Dr. JOH HENSCHEL received a M.Sc. in Marine Biology at the University of Cape Town and a Ph.D. in Mammal Ecology at the University of Pretoria. After completing a research fellowship of the Alexander von Humboldt Foundation in Germany, he became coordinator of research and information services at the DRFN. His association with this institution dates back to 1977. Dr. Henschel has published 60 scientific papers and 35 popular articles. His current research fields include population dynamics, food webs, climatology, natural resource management, and long-term ecological research, particularly in the Namib Desert.



Ms. JULIANE ZEIDLER submitted her Ph.D. thesis on Indicators of Biological Integrity during 1999 at the University of Witwatersrand. After travelling for one year by car from Germany through Africa, she became associated with the DRFN in 1991, and has since completed two M.Sc. degrees, one on Arthropod Ecology, and the second in Conservation Biology. Ms. Zeidler is currently the DRFN project manager of the Namibian Programme to Combat Desertification in which she is linking ecological research with community based research in the Namib and other areas of Namibia.



Mr. E.S. "CHARLIE" SHANYENGANA is a doctoral student in hydrochemistry at the University of Stellenbosch. The theme of his thesis is small-scale water desalination for rural areas. He received a M.Sc. in Environmental Engineering from Stockholm, Sweden, with a "best student award" from SIDA. Mr. Shanyengana obtained his B.Sc. in chemistry and zoology at the University of Namibia and has been directly associated with the DRFN since 1993. He manages DRFN's planning and research in appropriate technology.