African Herp News

Newsletter of the Herpetological Association of Africa



GEOGRAPHICAL DISTRIBUTIONS

AMPHIBIA: ANURA

MICROHYLIDAE

Phrynomantis affinis Boulenger, 1901 Spotted Rubber Frog

Namibia, Erongo Region, Brandberg, 2114BA, 16 July 1987, J.O. Wirminghaus, Angelo J. Lambiris collection, AJL2782.

The details of the *Phrynomantis affinis* tadpole illustrated in du Preez & Carruthers (2009, A Complete Guide to the Frogs of Southern Africa. Struik Nature, Cape Town, South Africa, Pp. 488) from the Brandberg extends the known distribution of this rare species by 275 km from Ombutjomatemba, Namibia (2017CA) in the north-east. This locality also extends the known range westward by 150 km for the species in Namibia as a whole from Oshigambo (1716CC), 19 km NNE of Odangwa and which is 375 km north of the Brandberg.

Acknowledgments

Louis du Preez answered queries on the source of the material for his and Carruthers' description of the Brandberg P. *affinis* tadpole. Angelo Lambiris sent details of the Brandberg P. *affinis* tadpole and a digital photograph of it which also served to verify his original identification. He generously gave permission to use his catalogued details for this note.

Submitted by:

John VISSER, P.O. Box 3599, Vineta, Swakopmund, Namibia. E-mail: jdvisser@iafrica.com.

REPTILIA: CHELONIA

TESTUDINIDAE

Kinixys spekii Gray,1863 Speke's Hinged Tortoise

Namibia, 110 km north of Grootfontein, 1818DB, S 18° 54', E 18° 47', 6 March, 2010, J.D. Visser.

The distribution of *Kinixys spekii* in Namibia has been recorded as the northern districts of Okavango and the Caprivi districts, specifically the Mamili and Mudumu National Parks, the Mahango Game Reserve and the Caprivi and Popa Falls Game