

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/227663576>

Phylogenetics, genome diversity and origin of modern leopard, *Panthera pardus*

Article in *Molecular Ecology* · November 2001

DOI: 10.1046/j.0962-1083.2001.01350.x

CITATIONS

211

READS

849

7 authors, including:



Olga Uphyrkina

Federal Biodiversity Center, Vladivostok, Russia

16 PUBLICATIONS 725 CITATIONS

[SEE PROFILE](#)



Howard Quigley

Panthera

156 PUBLICATIONS 4,805 CITATIONS

[SEE PROFILE](#)



Dale Miquelle

Wildlife Conservation Society

185 PUBLICATIONS 4,609 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Jaguar Corridor Initiative [View project](#)



reproductive parameters of wild femae amur tiegrs [View project](#)

Phylogenetics, genome diversity and origin of modern leopard, *Panthera pardus*

OLGA UPHYRKINA,*¶ WARREN E. JOHNSON,* HOWARD QUIGLEY,† DALE MIQUELLE,† LAURIE MARKER,‡ MITCHEL BUSH§ and STEPHEN J. O'BRIEN*

*Laboratory of Genomic Diversity, National Cancer Institute, Frederick, MD 21702–1201, USA, †Hornocker Wildlife Institute, Wildlife Conservation Society, Bozeman, MT 59719–0970, USA, ‡Cheetah Conservation Fund, Windhoek 9000, Namibia, Southwest Africa, §Conservation and Research Center, Front Royal, VA 22630, USA

Abstract

Leopards, *Panthera pardus*, are widely distributed across southern Asia and sub-Saharan Africa. The extent and phylogeographic patterns of molecular genetic diversity were addressed in a survey of 77 leopards from known geographical locales representing 13 of the 27 classical trinomial subspecies. Phylogenetic analysis of mitochondrial DNA sequences (727 bp of *NADH5* and control region) and 25 polymorphic microsatellite loci revealed abundant diversity that could be partitioned into a minimum of nine discrete populations, tentatively named here as revised subspecies: *P. pardus pardus*, *P. p. nimr*, *P. p. saxicolor*, *P. p. fusca*, *P. p. kotiya*, *P. p. delacouri*, *P. p. japonensis*, *P. p. orientalis* and *P. p. melas*. However, because of limited sampling of African populations, this may be an underestimate of modern phylogeographic population structure. Combined phylogeographic and population diversity estimates support an origin for modern leopard lineages 470 000–825 000 years ago in Africa followed by their migration into and across Asia more recently (170 000–300 000 years ago). Recent demographic reductions likely have led to genetic impoverishment in *P. p. orientalis* and in the island subspecies *P. p. kotiya*.

Keywords: evolution, microsatellites, mitochondrial DNA, *Panthera pardus*, subspecies

Received 14 September 2000; revision received 30 March 2001; accepted 8 May 2001

Introduction

The leopard, *Panthera pardus*, one of the most widely distributed and adaptable big cats, has pelage hues that vary from pale yellow to deep golden or tawny, and are patterned with black rosettes. Melanistic forms occur throughout its range, mostly in humid areas (Seidensticker & Lumpkin 1991; Nowell & Jackson 1996). The coat and colour patterns vary widely across various types of habitat. Pocock (1932) described four different colouration patterns that correspond to semidesert, savannah, rain forest and high mountain leopards. In the Russian Far East the leopard inhabits snowy temperate forests with winter temperatures reaching –25 °C, and displays a pale cream-

coloured long-hair winter coat that has led to its confusion with the snow leopard, *Panthera uncia* (Pocock 1930). Leopards occur at sea level (Africa, Arabia, India, Java), in foothill areas, in mountains, and on tops of volcanoes (Morocco, Turkmenistan, Iran, Russia, Java). The leopards are found in the Himalayas where they are sympatric with snow leopards up to 5200 m. Throughout their range the leopard feeds on a broad range of prey, including small rodents, birds, different species of ungulates and livestock (Hoogerwerf 1970; Nowell & Jackson 1996; Christen 2000).

The leopard's historic range spanned all of the sub-Saharan and north Africa, the Middle East and Asia Minor, South and Southeast Asia, and extended to the Amur Valley in the Russian Far East. Island ranges included Sri Lanka, Java, Zanzibar, and Kangean (Seidensticker & Lumpkin 1991; Nowell & Jackson 1996). Leopards are still found throughout most of their historic range (Fig. 1), although their numbers have been significantly reduced over the last hundred years due to increasing human population expansion, habitat loss, hunting, and poaching. In some

Correspondence: Stephen J O'Brien. Fax: 301 846 1686; E-mail: obrien@ncifcrf.gov

¶Present address: Laboratory of Evolutionary Zoology and Genetics, Institute Biology and Soil Sciences, Russian Academy of Sciences, Vladivostok, 690022, Russia




Fig. 1 Map showing: (i) historic (dark and light grey) and present (dark grey) geographical distribution of leopards; (ii) distribution of named classical leopard subspecies (big and small three-letter codes together) and distribution of revised subspecies classifications based on present work extending that of Miththapala *et al.* (1996) (big three-letter codes); and (iii) sample collection sites and number of samples from each site (circles). Full Latin trinomial subspecies names, subspecies codes, countries of their origins and collaborators that provided leopard samples are shown in Table 1.

areas, leopard populations have become heavily fragmented and isolated (Fig. 1). Except for central Africa and India, the leopard is endangered throughout its range. The Far Eastern (Amur, *P. p. orientalis*), Anatolian (*P. p. tulliana*), Arabian (*P. p. nimr*), and Barbary (*P. p. panthera*) leopards are considered to be almost extinct (listed as critically endangered by the IUCN Red List; Nowell & Jackson 1995). Caucasus (*P. p. ciscaucasica*), North Persian (*P. p. saxicolor*), Sri Lankan (*P. p. kotiya*), North-Chinese (*P. p. japonensis*), and Javan (*P. p. melas*) leopards are classified as endangered; and the Zanzibar (*P. p. adersi*) leopard is thought to be extinct (Nowell & Jackson 1995, 1996). The leopard species is listed in Appendix I of the Convention on International Trade of Endangered Species of Wild Fauna and Flora (CITES).

The leopard, along with the lion (*P. leo*), tiger (*P. tigris*), jaguar (*P. onca*) and snow leopard (*P. uncia*), comprise the relatively young felid genus *Panthera*, thought to have diverged from a common ancestor 2–3 million years ago (Ma) (Hemmer 1976; O'Brien *et al.* 1987; Wayne *et al.* 1993; Johnson & O'Brien 1997). The fossil records for the leopard as well as for other *Panthera* cats are controversial. Turner & Anton (1997) reported the earliest fossils for the leopard, along with the lion, at Laetoli in Tanzania at about 3.5 Ma. Prior to this report, the oldest leopard remains were reported from the Indian Siwaliks approximately 2 Ma; this primitive leopard resembled the jaguar (*P. onca*) and

the now extinct *P. gombazogensis* (Hemmer 1976; Kitchener 1991). During the earlier Middle Pleistocene (about 1 Ma), again along with the lion, the leopard was present in Africa and Europe, although in Europe it appeared earlier than the lion. European leopards at this time also resembled the Pleistocene North American jaguar (Hemmer 1976; Turner & Anton 1997). Teeth of ancient leopards found in southern China and dated from the Middle of Pleistocene were similar to the recent subspecies *P. p. sinensis*; this led to the hypothesis of local evolution in eastern and southeastern Asia (Hemmer 1976). In Java, the leopard was found in early Middle Pleistocene as well. Leopards in Java were thought to be very large in size (Hemmer 1976).

The leopard's extensive geographical distribution, its varied coat colour patterns and morphological characteristics led to the naming of 27 subspecies in early taxonomic treatments (Pocock 1930, 1932). Miththapala *et al.* (1996) analysed three molecular genetic methods [allozymes, mitochondrial DNA–restriction fragment length polymorphism (mtDNA–RFLP) and minisatellites] and morphological measurements to resolve six phylogeographic groups of leopards which corresponded to: (1) Africa; (2) central Asia; (3) India; (4) Sri Lanka; (5) Java; and (6) east Asia. Based on explicit subspecies definition criteria (Avise & Ball 1990; O'Brien & Mayr 1991), they recommended that the 27 classical leopard trinomials be reclassified into eight subspecies:

(1) *P. p. pardus* in Africa; (2) *P. p. saxicolor* in central Asia; (3) *P. p. fusca* in India; (4) *P. p. kotiya* in Sri Lanka; (5) *P. p. melas* in Java; (6) *P. p. orientalis* in Russian Far East; (7) *P. p. japonensis* in North China; and (8) *P. p. delacouri* in South China. The distinctiveness of the East Asian subspecies was not well supported due to limited sampling, and the molecular taxonomic definition of these subspecies remains provisional (Miththapala *et al.* 1996).

In the present study we revisit the assessment of molecular genetic variation and genetic differentiation in contemporary leopard populations. We examined leopard geographical partitioning with 25 feline-specific microsatellites, or short tandem repeat loci (STRs), and DNA sequence variation in two mtDNA regions, part of the *NADH-5* gene (611 bp) and the control region (CR, 116 bp). Our samples included 36 specimens from Miththapala *et al.* (1996) and an additional 41 specimens from other geographical locations (Table 1, Fig. 1). We used new methods and additional samples to test for subspecies/population differentiation and to compare the amount of genetic variation within identified leopard subspecies. The results were interpreted in terms of evolutionary history and phylogeography of the leopard in its natural habitat.

Materials and methods

Samples

Seventy-seven samples from leopards of known geographical origin were used (Table 1, Fig. 1). Subspecies *Panthera pardus saxicolor*, *P. p. delacouri*, *P. p. japonensis*, and *P. p. melas* were represented by captive bred individuals only. Samples were selected from unrelated (to the best of our knowledge) leopards with the exception of two *P. p. melas* which appeared to be relatives (A. Shoemaker, personal communication). Both traditional and revised subspecies names for each sampled leopard are listed in Table 1. Twenty-two wild tiger samples from five subspecies were used as the outgroup for microsatellite analysis (Wentzel *et al.* 1999). Individual samples of tigers, lions, jaguars, and snow leopards were used as outgroup species in mtDNA analysis.

DNA was extracted from whole blood, white blood cells, or fibroblast cultures from skin biopsies using a standard phenol–chloroform method (Sambrook *et al.* 1989). DNA from plasma was extracted using QIAamp DNA Blood Midi Kits (QIAGEN).

mtDNA sequence analysis

A fragment of 611 bp of the 5' end of the *NADH-5* mitochondrial gene corresponding to positions 12632–13242 in the complete *Felis catus* mtDNA sequence (Lopez *et al.* 1996) was amplified in two separate pieces that overlapped in approximately 140 bp, using primer pairs

F/RL2 (F: 5'-GTGCAACTCCAAATAAAAG-3' and RL2: 5'-TAAACAGTTGGAACAGGTT-3' and FL2/RL4 (FL2: 5'-CGTTACATGATCGATCATAG-3', and RL4: 5'-TTAGG-TTTCGTGTTGGGT-3'). All primers, except forward primer F (from Johnson *et al.* 1998), were developed from leopard (*P. pardus*) sequences. Forward primer FL2-nimr (5'-CGTTACATGGTCGATCATGG-3') was specifically designed for the Arabian leopard (*P. p. nimr*). Primers PDL-3 (Culver *et al.*, in preparation) and DLUP-4 (Hoelzel *et al.* 1993) were used to amplify 116 bp of the 5' variable region directly adjacent to the central conserved region of the mitochondrial CR.

Polymerase chain reactions (PCR) (25 µL) were performed using 2.5 ng of genomic DNA in 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 1.5 mM MgCl₂, 200 µM each of dATP, dCTP, cTTP, dGTP, 1 µM of each primer and 1 unit Taq-Gold DNA polymerase. For each reaction 35 cycles were performed with 0.5 min denaturation at 94 °C, 1.5 min annealing at 50 °C for *NADH-5* gene and 55 °C for CR, and 1 min extension at 72 °C. Products were checked in 2% agarose gel in TBE buffer. PCR products were purified with CENTRICON-100 filters (Amicon).

The *NADH-5* segment was sequenced in both forward and reverse directions using an ABI BigDye Terminator kit. The CR fragment was sequenced twice in the forward direction using an ABI FS Dye Primer kit (in this case PCR primers were designed to include M13 tails). Products were sequenced using Applied Biosystem 373 A and 377 Automated DNA sequencers. Sequences are deposited in GenBank (accession numbers AY035227–AY035292).

Sixty-nine from the total 77 leopards were taken to final analysis (only those successfully amplified and sequenced for total length of both mtDNA segments were taken). Preliminary results from separate analyses of *NADH-5* sequences produced phylogenetic associations similar to those when two segments were combined together [shown in Fig. 5 in form of linearized tree derived from neighbour-joining (NJ) tree topology]. The resolution of the maximum parsimony (MP) tree, however, was higher when both mtDNA segments, *NADH-5* and CR, were analysed together. The probability of estimating the correct tree seems to be higher when data from different genes are combined in the phylogenetic analysis (Bull *et al.* 1993; Huelsenbeck *et al.* 1996). Thus, we present our final data from combined sequenced analysis. To correct for different mutation rates in *NADH-5* and CR, the among site variation option (Yang & Kumar 1996) was applied throughout all mtDNA analysis.

Lion (*P. leo*), tiger (*P. tigris*), jaguar (*P. onca*), and snow leopard (*P. uncia*) individuals were sequenced for the homologous segments of *NADH-5* and CR to be used as outgroup species in the phylogenetic analysis. CR was not successfully amplified in lions using primers reported here; thus, only *NADH-5* sequences were used.

Table 1 Leopard sample collection used in the study

Subspecies Code ¹	Classical subspecies ²	Revised subspecies ³	Geographical area & Common name	Number individuals	Ppa number ⁴ (mtDNA Haplotype ⁵)	Status ⁶	Locale of origin	Sample Sources
Africa								
SUA	<i>Panthera pardus suahelicus</i>	<i>P. p. pardus</i> ⁷	East African	1	<u>155</u> (Sua1)	?		Chipangali Wildlife Trust, V. Wilson
REI	<i>P. p. reichenowi</i>	<i>P. p. pardus</i> ⁷	West African	1	<u>42</u> (Rei1)	?		Chipangali Wildlife Trust, V. Wilson
PAN	<i>P. p. panthera</i>	<i>P. p. pardus</i> ⁷	Barbary	1	<u>2</u> (Pan1)	W	Morocco	Carnivore Preservation Trust, M. Bleyman
SHO	<i>P. p. shortridgei</i>	<i>P. p. pardus</i> ⁷	Central African	14:				
				3	<u>801*</u> (Sho3), <u>802*</u> (Sho5), <u>803*</u> (Sho3)	W	Namibia	Cheetah Conservation Fund, L. Marker
				1	<u>804*</u> (Sho3)	W	Botswana	S. Osofsky
				5	<u>134*</u> (Sho6), <u>135*</u> (Sho7), <u>136*</u> (Sho8), <u>137*</u> (Sho1), <u>83*</u> (Sho2)	W	Kruger	Kruger National Park, M. Bush
				5	<u>33</u> , <u>35</u> (Sho4), <u>37</u> (Sho9), <u>38</u> (Sho4), 40	W	Zimbabwe	Chipangali Wildlife Trust, V. Wilson
Arabia								
NIM	<i>P. p. nimr</i>	<i>P. p. nimr</i> ⁷	South Arabian	1	<u>89*</u> (Nim1)	W	South Arabia	Tel Aviv University, H. Mendelssohn
Central Asia								
SIN	<i>P. p. sindica</i>	<i>P. p. saxicolor</i>	Baluchistan	1	<u>30</u> (Sin1)	W	Afghanistan	Lincoln Park Zoo, T. Meehan
SAX	<i>P. p. saxicolor</i>	<i>P. p. saxicolor</i>	Persian	9:				
				1	<u>45*</u> (Sax2)	C		Lowry Park Zoological Garden, D. Hansbury
				1	48	C		San Francisco Zoological Garden, C. Machado
				1	<u>49</u> (Sax1)	C		Berlin Zoological Garden, R. Goltenboth
				1	<u>75</u> (Sax2)	C		Koln Zoological Garden, O. Behlert
				1	76	C		Hannover Zoo, L. Dittrich
				1	<u>147*</u> (Sax2)	C		San-Petersburg Zoo, I. Korneev
				1	<u>148*</u> (Sax2)	C		Tallin Zoo, V. Fainstein
				1	<u>200</u> (Sax2)	C		Wilhelma Zoological-Botanical Garden, M. Holtkotter
				1	<u>203</u> (Sax2)	C		Welsh Mountain Zoo, N. Jackson
Java								
MEL	<i>P. p. melas</i>	<i>P. p. melas</i> ⁷	Javan	2				
				1	<u>50</u> (Mel1)	C		Berlin Zoological Garden, R. Goltenboth
				1	<u>195*</u> (Mel1)	C		Wuppertal Zoo, A. Sliwa
Sri Lanka								
KOT	<i>P. p. kotiya</i>	<i>P. p. kotiya</i>	Sri Lankan	11:				
				7	<u>102</u> (Kot1), <u>104</u> (Kot1), <u>105</u> (Kot1), <u>106</u> (Kot1), <u>116</u> (Kot2), <u>118</u> (Kot3), <u>128</u> (Kot1)	W	Sri Lanka	Sri Lanka National Zoological Garden
				4	<u>107</u> , <u>110</u> (Kot1), <u>112</u> (Kot1), <u>114</u> (Kot1)	C		Sri Lanka National Zoological Garden

Table 1 *Continued.*

Subspecies Code ¹	Classical subspecies ²	Revised subspecies ³	Geographical area & Common name	Number individuals	Ppa number ⁴ (mtDNA Haplotype ⁵)	Status ⁶	Locale of origin	Sample Sources
FUS	<i>P. p. fusca</i>	<i>P. p. fusca</i>	India Indian	9:	2 <u>87</u> (Fus1), <u>88</u> (Fus2) 7 <u>91*</u> (Fus5), <u>92*</u> (Fus3), <u>93*</u> (Fus5), <u>94*</u> (Fus5), <u>95*</u> (Fus6), <u>96*</u> (Fus4), <u>97*</u> (Fus5)	W W	South India North India	Nagarhole National Park, U. Karanth, India Sakkarbaug Zoo, Mr Rawal
DEL	<i>P. p. delacouri</i>	<i>P. p. delacouri</i>	East Asia South Chinese	4:	2 <u>108</u> (Del1), <u>115</u> (Del3) 2 <u>99*</u> (Del2), <u>211*</u>	C C		Sri Lanka National Zoological Garden Tierpark Berlin Zoological Garden, Dr Blaszkiewitz
JAP	<i>P. p. japonensis</i>	<i>P. p. japonensis</i>	North Chinese	11:	1 <u>18*</u> (Jap2) 5 <u>22*</u> (Jap2), <u>159*</u> (Jap2), <u>160*</u> (Jap2), <u>162*</u> , <u>163*</u> 2 <u>24</u> (Jap1), <u>26</u> (Jap1) 2 <u>52</u> (Jap2), <u>54</u> (Jap1)	C C C C		Toronto Metropolitan Zoo, G. Crawshaw EFBC's Feline Conservation Center Henry Doorly Zoo, D. Armstrong Royal Zoological Society of Antwerp, W. Meurichy San Diego Zoo, D. Janssen
ORI	<i>P. p. orientalis</i>	<i>P. p. orientalis</i>	Far Eastern (Amur)	12:	1 <u>79</u> (Jap1) 7 <u>149*</u> (Ori2), <u>150*</u> (Ori2), <u>151*</u> (Ori2), <u>152*</u> (Ori2), <u>153*</u> (Ori2), <u>156*</u> (Ori2), <u>157*</u> (Ori2) 3 <u>138*</u> (Ori1), <u>140*</u> (Ori2), <u>142*</u> (Ori2) 1 <u>144*</u> (Ori2) 1 <u>158*</u> (Ori2)	C W W W W	Russian Far East Korea Korea Korea	Hornocker Wildlife Institute, H. Quigley Moscow Zoo, V. Spitsin Tallin Zoo, V. Fainstein Tierpark Berlin Zoo, Dr Blaszkiewitz

¹Three letter code was assigned to each subspecies for convenient use throughout the figures.²Leopard subspecies as described in literature based on morphology and geographical distribution.³Leopard subspecies revised based on molecular genetic analysis presented in the paper.⁴Identification number of leopard individuals as they are listed in the exotic database collection at the Laboratory of Genomic Diversity, National Cancer Institute, Frederick, MD, 21702.⁵mtDNA haplotype assigned to each sample sequenced in the present study.⁶Status of each leopard: W-wild; C-captive bred; ?-status unknown.⁷Assigned to subspecies provisionally based on the current analysis.*Samples taken into analysis the first time (others were used in previous study; see Miththapala *et al.* 1996); underlined samples were sequenced.

Microsatellite loci genotyping

Leopard samples were analysed using 25 polymorphic microsatellite loci, originally isolated from the domestic cat, *Felis catus* (Menotti-Raymond *et al.* 1999). The selected loci (FCA008, FCA026, FCA043, FCA075, FCA077, FCA090, FCA094, FCA096, FCA097, FCA098, FCA105, FCA123, FCA126, FCA139, FCA161, FCA211, FCA220, FCA224, FCA229, FCA247, FCA310, FCA391, FCA441, FCA453, FCA678) were shown to be polymorphic in leopards and other nondomestic cat species (Driscoll 1998; Wentzel *et al.* 1999). Three loci (FCA 391, 441 and 453) were tetranucleotide repeats, and the others were dinucleotides. Twenty-three of 25 loci were mapped in the domestic cat (Menotti-Raymond *et al.* 1999); of these, two pairs of loci (FCA096-FCA075 and FCA224-FCA161) were linked at 9.0 cM (centimorgans, a unit of distance between genes on chromosomes) and 4.0 cM, respectively. A test for linkage disequilibrium performed for each pair of microsatellite loci revealed none across leopard populations.

Of 77 leopard samples, 75 were included in the microsatellite analysis. A DNA sample of a *P. p. nimr* (Ppa 89) extracted from a museum pelt that amplified only for 10 of the loci and DNA from a Javan leopard plasma sample (Ppa 195) with low yield were excluded from the analysis.

PCR amplifications for each microsatellite locus were performed as described in Menotti-Raymond *et al.* (1999). PCR amplification products were diluted with sterile deionized water in individual tubes, then multiplex pooled into groups of 4–5 loci based on product size and fluorescent dye label. Products were resolved by electrophoresis in an ABI Prism™ 310 Genetic Analyser. Data were analysed using ABI Prism™ GENESCAN 2.1 and GENOTYPER 2.0. Amplification products of at least two samples were electrophoresed in each run as standards to correct allele size differences if necessary. PCR product length was used as an actual repeat length (Ellegren 1995).

Data analysis

mtDNA. Leopard *NADH-5* and CR sequences were edited and aligned using SEQUENCHER (Gene Codes Co., Ann Arbor, Michigan) and CLUSTALX (Thompson *et al.* 1997) and checked visually. The *NADH-5* mtDNA gene sequences were translated into 203 codons. Thirteen sites in leopard sequences could not be unambiguously scored and were excluded from the analysis.

Phylogenetic relationships among mtDNA haplotypes were assessed using three methods implemented in PAUP* version 4.0 (Swofford 1998): maximum parsimony (MP), minimum evolution (ME), and maximum likelihood (ML). The MP analysis was conducted using a heuristic search, with random addition of taxa and tree-bisection-reconnection (TBR) branch swapping. In the ME approach,

neighbour-joining (NJ) trees (Saitou & Nei 1987) were generated with Kimura 2-parameter (Kimura 1980) γ -corrected distances; TBR branch swapping was used next to find a minimum evolution tree. The shape parameter (α) for the γ -distribution was estimated using the PAMP program in the PAML software package (Yang & Kumar 1996; Yang 1999), and was equal to 0.29. The ML analysis was performed using the HKY85 model (Hasegawa *et al.* 1985) with the among site variation option (α set to 0.29 as estimated by PAML). Reliability of all trees was tested by 100 bootstrap replications (Hillis & Bull 1993).

Five different scenarios of leopard geographical subdivision were tested by F_{ST} (with γ -corrected Kimura 2-parameter distances) using the AMOVA algorithm (Excoffier *et al.* 1992) as implemented in the ARLEQUIN 1.1 (Schneider *et al.* 1997). *P. p. nimr* and *P. p. melas* were excluded from subdivision analysis due to limited sample size. Parameters of genetic variability for leopard populations were assessed with MEGA 1.01 (Kumar *et al.* 1993) and ARLEQUIN 1.1, and were measured in terms of polymorphic sites, number of pairwise differences, and nucleotide diversity (genetic diversity of populations with a sample size of four or more was estimated).

The approximate age of modern leopard lineages was estimated using LINTRE (Takezaki *et al.* 1995). This program tests the molecular clock on a given topology of a phylogenetic tree and makes linearized trees re-estimating branch lengths under the assumption of a constant rate of evolution (Takezaki *et al.* 1995). A phylogenetic tree was constructed using the NJ method (Saitou & Nei 1987) and Kimura 2-parameter γ -corrected distances for *NADH-5* sequences (611 bp) only. The parameter α for γ -correction for the *NADH-5* sequences was estimated using PAML ($\alpha = 0.90$). Both the two-cluster and branch-length tests implemented in LINTRE were applied (Takezaki *et al.* 1995). The coalescence point between leopard and lion haplotypes was chosen to be a calibration point and two fossil dates were used. First, 3.5 Ma was used because it has been recorded as the earliest fossils in Tanzania, Africa, for both leopards and lions (Turner & Anton 1997). Second, 2 Ma was used because it is believed to be the lowest bound for the proposed time of split in the *Panthera* lineage (O'Brien *et al.* 1987; Wayne *et al.* 1993). Recent evidence suggests that the snow leopard is a basal divergence in the *Panthera* genus (Johnson & O'Brien 1997), therefore the snow leopard was used as an outgroup.

Microsatellites. Pairwise genetic distances among individual leopards using microsatellite data were estimated based on the proportion of shared alleles (Dps) and the kinship coefficient (Dkf) (Bowcock *et al.* 1994) with [1 – ps/kf] option as implemented in MICROSAT (Minch *et al.* 1995). The program NEIGHBOR (included in PHYLIP package; Felsenstein 1985a) was used to construct NJ phylogenetic

trees from the distance matrices. Nei's D_a genetic distances (Nei *et al.* 1983) computed with DISPAN (Ota 1993) were used for pairwise comparisons among leopard populations. D_a genetic distances are not proportional to evolutionary time, but have been shown to generate correct phylogenetic trees under various evolutionary conditions (Nei & Takezaki 1996; Takezaki & Nei 1996). One hundred bootstrap iterations were used to estimate the reliability of nodes uniting leopard individuals; one thousand iterations were used for the leopard population trees (Felsenstein 1985b).

Five different subdivision scenarios among leopard populations based on STR data were assessed by R_{ST} , sum of squared size differences (Slatkin 1995), in ARLEQUIN 1.1. A Mantel correlation test between pairwise F_{ST} and R_{ST} values was applied with 1000 iterations using MANTEL 2 (Liedloff 1999) to test whether or not 'subspecies' subdivisions estimated with mtDNA and microsatellite data are congruent. Tests for significance of deviation from Hardy-Weinberg equilibrium for each locus in each population (Guo & Thompson 1992), and tests for genotypic linkage disequilibrium for each pair of loci in each population were performed using GENEPOP software (version 3.1) (Raymond & Rosset 1995). Significant deviations from Hardy-Weinberg equilibrium ($\alpha = 0.05$) showing deficiency of heterozygotes were found at three loci in three populations: *P. p. japonensis* (FCA126), *P. p. pardus* (FCA097), and *P. p. kotiya* (FCA441) after a Bonferroni correction (Rice 1989).

Variability across 25 microsatellite loci for each leopard population (for those with a sample size of four or more) was measured in terms of percentage of polymorphic loci, average expected heterozygosity, average number of alleles per locus, percentage of unique alleles, average and maximum range of microsatellite repeats, and average variance of microsatellite repeats. Measures of genetic variation were estimated using MICROSAT and EXEL. Estimates of leopard microsatellite diversity may be biased since only polymorphic loci were used in the present study.

Results

Phylogenetic analysis

To test for evidence of leopard population/subspecies differentiation, the relationships among individual leopards were examined by phylogenetic analyses of mtDNA haplotypes and of composite genotypes from 25 microsatellite loci.

mtDNA. The two combined mtDNA regions of 69 leopards revealed 50 variable sites (44 in *NADH-5* and six in the CR), which defined 33 haplotypes in leopards (Table 2). The haplotypes of each leopard are listed in Table 1. All 13 classical leopard subspecies sampled had private mtDNA haplotypes, i.e. a haplotype found in only one subspecies

(Table 2). The phylogenetic analysis of haplotypes using MP, ME and ML produced concordant topologies (Fig. 2). African and Asian leopards assorted into separate monophyletic groups with two exceptions. The exceptions involved two haplotypes, one found in two *Panthera pardus melas* (Mel1) leopards and the other in *P. p. nimr* (Nim1) leopard. These differed from other haplotypes and did not consistently cluster with African or Asian clusters (Fig. 2).

Two clusters of African leopard haplotypes (labelled as PAR-I and PAR-II in Fig. 2) were resolved with relatively high bootstrap support (76% and 84% for the group I and II, respectively, in MP, 78% and 79% in ME, and 87% and 84% in ML). These two groups along with the *P. p. nimr* (Nim1) were basal in the MP analysis relative to a cluster of Asian leopards (Fig. 2a). The ME tree topology differed in that two clusters of African leopards, PAR-I and PAR-II grouped together (63%) and defined sister taxa with the *P. p. nimr* haplotype (67%) (Fig. 2b).

Within the cluster of Asian leopards, haplotypes belonging to a particular geographical subspecies tended to group together (Fig. 2a,b). A haplotype representative of *P. p. sindica* (Sin1) clustered closely to *P. p. saxicolor* leopards, differing by a single site (position 23) in the CR sequence (Table 2). Three contiguous east Asian subspecies, *P. p. delacouri* (Del1-Del3), *P. p. japonensis* (Jap1, Jap2), and *P. p. orientalis* (Ori1, Ori2) associated, albeit weakly, in all analyses (Fig. 2a,b).

Haplotypes from the African PAR-I group were represented by several *P. p. shortridgei* leopards from different countries of southern Africa (Table 1), and also by single representatives of other African classical subspecies: *P. p. panthera* (Pan1), *P. p. reichenowi* (Rei1) and *P. p. suahelicus* (Sua1) (Table 1, Fig. 1). The four African individuals with PAR-II haplotypes were restricted to *P. p. shortridgei*: three from Kruger National Park, South Africa (haplotypes Sho6-Sho8) and one from Zimbabwe (Sho9; Table 1).

Microsatellites. Using the composite genotypes of 25 microsatellite loci from 75 leopard individuals, NJ phylogenetic trees were constructed using Dps and Dkf genetic distances with [1 - (ps/kf)] option. Both distances produced a similar topology: in all trees leopard individuals tended to cluster together according to their geographical origins, forming eight groups (Fig. 3). In contrast to mtDNA data, there was no evidence from microsatellite genotypes for the PAR-I/PAR-II subdivision among the African leopards. Leopards from mitochondrial groups PAR-I and PAR-II clustered together and were not significantly distinctive with microsatellite R_{ST} value ($R_{ST} = 0.009$; $P = 0.297$). The single *P. p. sindica* associated with *P. p. saxicolor* leopards, forming a group of central Asian leopards. Sri Lankan leopards (*P. p. kotiya*) were grouped in the same cluster, but closely aligned with Indian leopards, *P. p. fusca* (Fig. 3). Among East Asian leopards three classical subspecies

Table 2 Haplotypes and variable sites in combined analysis of the NADH-5 (611 bp) and CR (116 bp) mtDNA in leopard (*P. pardus*) and outgroup species, lions (*P. leo*), tigers (*P. tigris*), jaguars (*P. onca*) and snow leopard (*P. uncia*)

		NADH5																																																CR											
Species	Haplotype	N***	*8	10	16	21	23	31	57	62	69	75	87	123	137	147	154	162	165	168	181	187	210	213	219	240	249	252	260	264	270	276	279	294	303	318	321	327	381	435	447	450	477	498	537	609	**10	18	22	23	40	92									
<i>P. uncia</i>	Pun	2	A	.	.	T	.	T	T	.		.	T	T	.	A	.	A	G	.	A	.	A	T	.	T	G	T	C	.	A	C	G	.	G	G									
<i>P. onca</i>	Pon	1	C	.	.		T	.	.	.	T	T	T	A	A	A	.	.	.	C	A	.	A	C	C	.	C	T	.	G	G									
<i>P. tigris</i>	Pti	2		T	.		.	.	T	.	A	A	A	T	A	C	C	A	C	T	.	G	G									
<i>P. leo</i>	Ple	2	T	T	.	.	.		T	A	.	A	T	T	.	C		.	.	T	.	.	.		G	-	-	-	-	-									
<i>P. p. pardus</i>																																																													
<i>P. p. panthera</i>	Pan	1	T	T	T	T	T	G	C	T	C	A	C	C	T	T	A	C	C	C	G	G	T	C	C	T	T	T	T	C	C	A	A	G	A	G	C	C	C	A	C	A	T	T	T	T	G	T	C	A	A	A									
<i>P. p. suahelicus</i>	Sua	1	T	.	.	.		T	C	.	.	T				C										
<i>P. p. reichenowii</i>	Rei	1		T	.	.	G	C	.	A	C										
<i>P. p. shortridgei</i>	Sho1	1		T										
<i>P. p. shortridgei</i>	Sho2	1		T									
<i>P. p. shortridgei</i>	Sho3	3		T								
<i>P. p. shortridgei</i>	Sho4	2		T								
<i>P. p. shortridgei</i>	Sho5	1		T								
<i>P. p. shortridgei</i>	Sho6	1	A	.	C	.	.		T	A	.	A	.	A	T	T	C	.	.	A	C							
<i>P. p. shortridgei</i>	Sho7	1	A	.	C	.	.		T	A	.	A	.	A	T	T	C	.	.	A	C	T	.	G							
<i>P. p. shortridgei</i>	Sho8	1	A	.	C	.	.		T	A	.	A	.	A	T	C	.	.	A	C	T	.	G								
<i>P. p. shortridgei</i>	Sho9	1	A	.	C	.	.		T	A	.	A	.	A	T	T	C	.	.	A	C	T	.	G							
<i>P. p. nimr</i>	Nim1	1	G	.	T	.		.	.	T	.	A	.	A	T	.	C	G	.	.	.	T	T	G	.	.	.	C	.	.	A	C	T	.	.	.						
<i>P. p. melas</i>	Mel1	2		T	.		.	.	T	T	A	.	A	T	T	.	C	.	C	A	.	A	C	.	.	A	C	T	.	.	G					
<i>P. p. saxicolor</i>																																																													
<i>P. p. saxicolor</i>	Sax1	1	T	T	.	C	.	.	T	.	A	.	A	T	C	G	A	.	A	.	T	C	.	.	A	C	T	.	.	.								
<i>P. p. saxicolor</i>	Sax2	6	.	C	T	T	.	C	.	.	T	.	A	.	A	T	C	G	A	.	A	.	T	C	.	.	A	C	T	.	.	.								
<i>P. p. sindica</i>	Sin	1	.	C	T	T	.	C	.	.	T	.	A	.	A	T	C	G	A	.	A	.	T	C	.	.	A	C	T	C	.	.	.							
<i>P. p. fusca</i>	Fus1	1	T	.	.	.	T	.	C	.	.	T	.	A	.	A	T	C	G	G	A	.	A	C	.	.	A	C	T	.	.	.								
	Fus2	1	T	.	C	.	.	T	.	A	.	A	T	C	G	G	A	.	A	C	.	.	A	C	T	.	.	.								
	Fus3	1	T	T	.	C	.	.	T	.	A	.	A	T	C	G	G	A	C	.	.	A	C	T	.	.	.							
	Fus4	1	T	T	.	C	.	.	T	.	A	.	A	T	C	G	G	A	.	A	.	T	.	.	.	C	.	C	.	A	C	T	.	.	.							
	Fus5	4	T	T	.	C	.	.	T	.	A	.	A	T	C	G	G	A	.	A	.	T	C	.	.	A	C	T	.	.	.							
	Fus6	1	T	T	.	C	.	.	T	.	A	.	A	T	C	G	G	A	.	A	.	T	.	T	.	.	.	C	.	.	A	C	T	.	.	.							
<i>P. p. kotiya</i>	Kot1	8	.	.	.	C	T	T	.	C	.	.	T	.	A	.	A	T	C	C	.	.	.	G	A	.	A	.	T	C	.	.	A	C								
	Kot2	1	.	.	.	C	T	T	.	C	.	.	T	.	A	.	A	T	C	C	.	.	.	G	A	.	A	.	T	.	T	.	.	.	C	.	.	A	C								
	Kot3	1	.	.	.	C	T	T	.	C	.	.	T	.	A	.	A	T	C	C	.	.	.	G	A	.	A	.	T	.	T	.	.	.	C	.	.	A	C	T	.	.	.								
<i>P. p. delacouri</i>	Del1	1	.	.	C	.	C	T	T	.	C	.	.	T	.	A	.	A	T	C	G	A	.	A	.	T	.	.	.	G	.	C	.	C	.	A	C	T	.	G	G							
	Del2	1	C	T	T	.	C	.	.	T	.	A	.	A	T	C	G	A	.	A	.	T	.	T	.	.	G	.	C	.	C	.	A	C	T	.	G	.						
	Del3	1	.	.	C	.	C	T	T	.	C	.	.	T	.	A	.	A	T	C	G	A	.	A	.	T	C	.	.	A	C	T	.	G	G								
<i>P. p. japonensis</i>	Jap1	4	T	T	.	C	.	.	T	.	A	.	A	T	C	C	.	.	.	G	A	.	A	.	T	C	.	.	A	C	T	.	G	.								
	Jap2	5	T	T	.	C	.	.	T	.	A	.	A	T	C	G	A	.	A	.	T	C	.	.	A	C	T	.	G	.								
<i>P. p. orientalis</i>	Ori1	1	C	T	T	.	C	.	.	T	.	A	.	A	T	C	G	A	.	A	.	T	C	.	.	A	C	T	.	G	.								
	Ori2	11	C	T	T	.	C	.	.	T	.	A	.	A	T	C	G	A	.	A	.	T	.	T	.	.	.	C	.	.	A	C	T	.	G	.								

*From the beginning of NADH-5 gene; 1 corresponds to 12634 in the complete *Felis catus* mtDNA sequence (Lopez *et al.* 1996).

**From the beginning of sequenced portion of the Control Region (116 bp).

***Number of individuals with each haplotype.

Haplotype of *P. p. panthera* is the reference sequence. Nucleotides diagnostic for a particular subspecies (or two phylogenetic groups, PAR




Fig. 2 Phylogenetic relationships among the leopard mtDNA haplotypes from combined *NADH-5* (611 bp) and control region (CR, 116 bp) mitochondrial regions (Table 2). Individual samples of *Panthera leo*, *P. tigris*, *P. onca*, and *P. uncia* are taken as outgroup species. (a) Maximum parsimony (MP) tree. MP tree constructed with PAUP* (Swofford 1998) and a general heuristic search; with random taxon addition and tree-bisection reconnection branch swapping; shown in majority rule consensus of 10 145 trees (length = 212, CI = 0.684). Numbers above branches represent bootstrap support (100 replicates); only those with > 50% are shown. Numbers below show number of steps/number of homoplasies. (b) Minimum evolution (ME) tree, constructed with PAUP* using Kimura 2-parameter γ -corrected distances ($\alpha = 0.29$ as defined by PAML) and neighbour-joining algorithm followed by tree-bisection-reconnection branch swapping. Values above branches represent support from 100 bootstrap replicates (only those with > 50% are shown). Maximum Likelihood (ML) approach performed using HKY85 model (Hasegawa *et al.* 1985) with the among site variation option (the α set to 0.29 and estimated with PALM) produced generally the same topology as ME tree (not shown). Bootstrap support from 100 replicates for ML tree is shown in ME tree (b) below branches (only those with > 50% are shown).

— *P. p. delacouri*, *P. p. japonensis*, *P. p. orientalis* — formed separate apparently monophyletic lineages. *P. p. melas* clustered separately and basal relative to all other leopards (Fig. 3).

Population subdivision and subspecies recognition

To evaluate the extent of population differentiation in leopards we tested five different geographical scenarios and compared them based on analysis of molecular

variance (AMOVA) with both mtDNA and microsatellite data. The first scenario considered only two groups: all African leopards (*P. p. shortridgii*, *P. p. panthera*, *P. p. reichenowi*, *P. p. suahelicus*) vs. all Asian leopards (*P. p. saxicolor*, *P. p. sindica*, *P. p. fusca*, *P. p. kotiya*, *P. p. delacouri*, *P. p. japonensis*, and *P. p. orientalis*); these major groups were proposed based on the topology of the mtDNA phylogenetic trees (Fig. 2). AMOVA performed using mtDNA found 68.9% of the variation (F_{ST} , Table 3) between the continents and 31.1% within the continents ($P < 0.0001$). With microsatellites,




Fig. 3 Phylogenetic relationships among the individual leopards based on 25 microsatellite loci. Branches of the same colour represent leopard individuals of the same classically named subspecies defined by three-letter codes (Table 1). Trees constructed based on proportion of shared alleles (Dps) and kinship coefficient (Dkf) genetic distances with $1 - (kf/ps)$ option in MICROSAT (Minch *et al.* 1995) produced the identical topologies; Dps tree is shown. Numbers are individual Ppa identification (Table 1).

36.1% of the variation (R_{ST} , Table 3) could be explained by continent subdivision, and 63.9% of the variation was retained within the continents ($P < 0.0001$).

In a second test, leopards were considered as three geographical groups: (1) African (as above); (2) Central Asian, Indian and Sri Lankan together (*P. p. saxicolor*, *P. p. sindica*, *P. p. fusca*, and *P. p. kotiya*); and (3) East Asian (*P. p. delacouri*, *P. p. japonensis*, and *P. p. orientalis*). This scheme was suggested by the topology of the microsatellite trees (Fig. 3). With this grouping, 63% of mtDNA and 31.6% of microsatellite variation can be explained by geographical partitioning (Table 3), and 37% and 68.4% of the variation, respectively, are retained within the groups ($P < 0.0001$). The third scenario involved four geographical groups

and differed from the second in that central Asian (*P. p. saxicolor*, *P. p. sindica*), and Indian and Sri Lankan (*P. p. fusca*, and *P. p. kotiya*) were analysed separately. Among group variation in this case was slightly higher than in previous scenario, 63.3% with mtDNA ($P < 0.0001$) and 32.9% with micro-satellites ($P < 0.0001$; Table 3).

In the fourth scenario applied, leopards were divided into seven different groups that correspond to the revised subspecies, based on the mtDNA and microsatellite phylogenetic analysis (Figs 2, 3 and 4). With this grouping, 76.04% of mtDNA variation was distributed among leopard populations and only 23.96% ($P < 0.0001$) within the populations (Table 3). With microsatellite data, 35.8% of the variation was found between the subspecies, while

Table 3 Measures of geographical subdivision in *Panthera pardus* based on analysis of molecular variance (AMOVA) with mtDNA and microsatellite data

Subdivision	mtDNA F_{ST}^{\dagger}	Microsatellites R_{ST}^{\ddagger}
2 groups:		
AF vs. AS	0.689	0.361
3 groups:	0.630	0.316
AF vs. CA, I, S	0.676	0.291
AF vs. EA	0.724	0.447
CA, I, S vs. EA	0.394	0.220
4 groups:	0.633	0.329
AF vs. CA	0.666	0.367
AF vs. I, S	0.655	0.250
AF vs. EA	0.724	0.447
CA vs. I, S	0.372*	0.211
CA vs. EA	0.499	0.234
I, S vs. EA	0.431	0.288
7 groups (subspecies§):	0.760	0.358
8 groups (subspecies¶):	0.838	0.363

F_{ST} and R_{ST} values were significant with $P < 0.0001$; *significant with $P < 0.002$.

Different subdivision scenarios are described in the text.

AF, African (*P. p. shortridgei*, *P. p. panthera*, *P. p. reichenowi*, *P. p. suahelicus*); AS, Asian (*P. p. saxicolor*, *P. p. sindica*, *P. p. fusca*, *P. p. kotiya*, *P. p. delacouri*, *P. p. japonensis*, and *P. p. orientalis*); CA, Central Asian (*P. p. saxicolor* and *P. p. sindica*); I, S, Indian and Sri Lankan (*P. p. fusca* and *P. p. kotiya*); ES, East Asian (*P. p. delacouri*, *P. p. japonensis*, and *P. p. orientalis*); CA, I, S, Central Asian, Indian and Sri Lankan together.

§Includes seven revised subspecies: *P. p. pardus*, *P. p. saxicolor*, *P. p. fusca*, *P. p. kotiya*, *P. p. delacouri*, *P. p. japonensis*, and *P. p. orientalis*.

¶The same as 7 groups, but African divided into *P. p. pardus* I (PAR-I) and *P. p. pardus* II (PAR-II). *P. p. nimr* and *P. p. melas* were excluded from all scenarios due to limited sampling.

†Calculated with Kimura 2-Parameter distances (Kimura 1980).

‡Calculated with R_{ST} option in ARLEQUIN 1.1 (Schneider *et al.* 1997).

64.2% of the variation occurred within the subspecies ($P < 0.0001$). When African leopards were considered as two groups, PAR-I and PAR-II (the last scenario), subdivision was even higher: $F_{ST} = 83.8\%$, and $R_{ST} = 36.3$ ($P < 0.0001$; Table 3). Based on these tests, variation among leopard

populations was best explained by grouping leopards on the basis of 'subspecies' scenarios (see below).

The statistical significance of pairwise population differentiation was tested by F_{ST} for mtDNA data and by R_{ST} for microsatellite data (Table 4). Each population was significantly different from the others by pairwise F_{ST} for mtDNA data. For microsatellites, two population comparisons of R_{ST} , PAR-I and PAR-II, and DEL and JAP, were not significantly different ($P = 0.297$ and 0.069 , respectively). R_{ST} pairwise comparisons performed with African leopards considered to be a single group (PAR) revealed all revised subspecies to be distinctive ($P < 0.001$). Mantel correlation analysis between F_{ST} and R_{ST} pairwise values (given in Table 4) revealed the two matrices to be significantly correlated ($P < 0.005$; $g = 5.661$, $Z = 20138$ and $r = 0.841$).

In summary, the position of each mtDNA haplotype (Fig. 2) and each leopard's composite microsatellite genotype in phylogenetic trees (Fig. 3) correlated well with their geographical origins; however, there was not strong bootstrap support for the phylogeographic clusters. The phylogeographic concordance of both mtDNA and microsatellite analyses plus the significant partitions of distinctive groups (Tables 3 and 4) would support the recognition and genetic distinctions for a minimum of nine groups: *P. p. pardus*-PAR, *P. p. saxicolor*-SAX, *P. p. nimr*-NIM, *P. p. fusca*-FUS, *P. p. kotiya*-KOT, *P. p. delacouri*-DEL, *P. p. japonensis*-JAP, *P. p. orientalis*-ORI, and *P. p. melas*-MEL, which we propose to recognize as revised subspecies of *P. pardus*.

A phylogenetic analysis of seven revised leopard subspecies (i.e. those with multiple individuals) was constructed using D_a genetic distances for microsatellite population data (Fig. 4). The deepest split separated African leopards, *P. p. pardus* (combined PAR-I and PAR-II) from other leopard groups. Central Asian leopards, *P. p. saxicolor*, followed Africans in the phylogenetic tree. Indian leopards (*P. p. fusca*) clustered with Sri Lankan (*P. p. kotiya*) individuals with relatively high bootstrap support (83%). The three East Asian subspecies formed a monophyletic lineage with high statistical support (99%): *P. p. orientalis* consistently grouped with *P. p. japonensis*; and the pair formed a sister taxon with *P. p. delacouri* (Fig. 4). This phylogenetic tree corresponds rather well with the geographical distribution of leopard populations.




Fig. 4 Relationships among revised leopard subspecies based on Neighbour-joining analysis of 25 microsatellite loci and D_a (Nei *et al.* 1983) genetic distances. Numbers indicate bootstrap percentages.

	PAR	PAR I	PAR II	SAX	FUS	KOT	DEL	JAP	ORI
PAR	—	—	—	0.436	0.296	0.336	0.409	0.511	0.530
PAR I	—	—	0.009	0.345	0.210	0.259	0.289	0.416	0.439
PAR II	—	0.690	—	0.554	0.383	0.417	0.518*	0.606	0.695
SAX	0.656	0.832	0.896	—	0.163	0.341	0.178	0.330	0.485
FUS	0.617	0.783	0.810	0.546	—	0.169	0.218	0.320	0.435
KOT	0.670	0.849	0.915	0.873	0.730	—	0.323	0.460	0.528
DEL	0.620	0.799	0.802	0.800	0.639	0.865	—	0.167	0.401
JAP	0.663	0.835	0.883	0.727	0.574	0.812	0.670	—	0.340
ORI	0.750	0.889	0.946	0.941	0.809	0.950	0.898	0.862	—

All populations were significantly different ($P < 0.01$) by F_{ST} values based on mitochondrial data. Two pairs of populations were not significantly different by R_{ST} based on microsatellite data: PAR I and PAR II ($P = 0.297$) and DEL and JAP ($P = 0.069$). The rest were significantly different ($P < 0.01$).

* $P < 0.02$.

Genetic variation

Estimates of population genetic variability calculated from mtDNA sequences and microsatellite loci for each revised leopard subspecies are summarized in Table 5. Genetic diversity varied appreciably among the leopard subspecies, and trends were similar for both mtDNA sequences and microsatellites. Estimates of genetic variation were the highest in African leopards, *P. p. pardus*, lowest in the Far Eastern leopard *P. p. orientalis*, substantially reduced in Sri Lankan *P. p. kotiya* leopards, and moderate in other populations. With mtDNA data, for example, there was 21 variable sites in *P. p. pardus* population while there was only one in *P. p. orientalis* or *P. p. japonensis*. Mean number of pairwise nucleotide differences in *P. p. pardus* (8.77) was more than 50 times higher than within *P. p. orientalis* (0.17) or about 15 times higher than in the *P. p. kotiya* (0.56). Nucleotide diversity (π) was high in *P. p. pardus* (1.22%), very low in *P. p. orientalis* (0.02%), and in between these two extremes in other subspecies populations (Table 5).

The African leopards, *P. p. pardus*, also had the highest diversity estimates for microsatellite loci. Heterozygosity in this population was the highest (0.803), somewhat lower in *P. p. fusca* (0.696) and in *P. p. saxicolor* (0.616), relatively low in *P. p. kotiya* (0.485), and lowest in *P. p. orientalis* (0.356). The average number of microsatellite alleles, range of microsatellite repeats and microsatellite variance had the same trends: highest in *P. p. pardus* (8.52, 9.72 and 7.28, respectively), lowest in *P. p. orientalis* (2.60, 2.84 and 1.71) and moderate in all others (Table 5). The microsatellite allele size distribution was most heterogeneous in African leopards (allele size distributions for each population and each locus are given as complementary information to this paper at <http://lgd.nci.nih.gov>). Alleles found in *P. p. orientalis* and in *P. p. kotiya* populations were almost always a subset of those seen in *P. p. japonensis* and *P. p. fusca*, respectively, and they were discontinuously distributed,

which may suggest a founder effect in history of *P. p. orientalis* and *P. p. kotiya* populations.

Diagnostic characteristics

Each revised leopard subspecies possessed population-specific mtDNA haplotypes, and/or microsatellite alleles (Table 6). Diagnostic mitochondrial sites were found in every revised subspecies except *P. p. japonensis*. Two mtDNA sites were specific for *P. p. orientalis*; one for *P. p. delacouri*, *P. p. fusca* and *P. p. kotiya*; three sites were unique for *P. p. melas* and five for *P. p. nimr* (Tables 2 and 6). Three fixed sites present in *P. p. pardus* leopards are not considered to be diagnostic for the African group since these sites were found in *P. p. nimr* as well, suggesting close evolutionary relatedness of these two subspecies (Table 2). Three sites were shared among *P. p. pardus*, *P. p. nimr* and *P. p. melas* leopards as well as with most outgroup species (*P. tigris*, *P. leo*, *P. onca*, and *P. uncia*) (Table 2). African leopards in general had the largest number of mitochondrial sites in common with outgroup species (Table 2).

All revised subspecies, except *P. p. saxicolor* and *P. p. melas*, revealed subspecies-specific microsatellite alleles (Table 6). Frequencies of such private alleles, however, were low in each population (2.83–5.80% of total number of alleles), with the exception of African leopards *P. p. pardus*, where they were 29.0% (Table 5). Number of diagnostic mtDNA sites and percentage of subspecies-specific microsatellite alleles should be considered relative to the sample sizes of leopard populations presented here.

Estimation of divergence times

The mtDNA sequence divergences were used to test the hypothesis of a molecular clock for different leopard haplotypes and to estimate approximate times of leopard divergence (Fig. 5, Takezaki & Nei 1996). The two-cluster

Table 4 Population pairwise F_{ST} estimates using the combined data from the mitochondrial regions and Kimura 2-parameter corrected distances (below the diagonal). Population pairwise R_{ST} estimates using data from 25 microsatellite loci (above the diagonal)

Table 5 Genetic variation across mtDNA gene segments (*NADH-5*, 611 bp, and Control Region, 116 bp) and 25 microsatellite loci in seven revised leopard subspecies

Subspecies	mtDNA				Microsatellites						
	Number leopards mtDNA/µsat	Number variable sites	Mean number pairwise differences (SE)	$\pi \times 10^2$ (SE)	% Polymorphic loci	Average H_E (SE)	Average number alleles/locus	% Specific alleles	Average range repeat/locus	Microsatellite variance	Maximum range
<i>Panthera pardus</i>	69/75	50	8.67 (4.40)	1.21 (0.62)	100	0.793 (0.073)	11.08	—	12.64	7.11	17
<i>P. p. pardus</i> (I + II)	15/17	21	8.77 (4.29)	1.22 (0.67)	100	0.803 (0.076)	8.52	29.1	9.72	7.28	15
PAR I	11/13	14	4.78 (2.53)	0.67 (0.40)	100	0.795 (0.099)	8.36	20.1	10.28	7.59	13
PAR II	4/4	3	3.75 (2.38)	0.52 (0.39)	100	0.675 (0.083)	4.08	3.92	6.00	5.24	9
<i>P. p. saxicolor</i>	8/10	2	0.50 (0.47)	0.07 (0.07)	100	0.616 (0.083)	4.24	2.83	5.12	4.28	7
<i>P. p. fusca</i>	9/9	8	2.61 (1.54)	0.36 (0.24)	100	0.696 (0.144)	5.52	5.80	6.2	5.38	9
<i>P. p. kotiya</i>	10/11	2	0.56 (0.50)	0.08 (0.08)	96	0.485 (0.202)	3.52	5.68	4.58	4.25	7
<i>P. p. delacouri</i>	3/4	5	3.41 (2.37)	0.48 (0.41)	100	0.674 (0.126)	4.20	5.71	5.56	5.70	6
<i>P. p. japonensis</i>	9/11	1	0.95 (0.71)	0.21 (0.15)	100	0.549 (0.171)	3.76	3.19	4.44	2.70	7
<i>P. p. orientalis</i>	12/12	1	0.17 (0.24)	0.02 (0.04)	92	0.356 (0.222)	2.60	3.07	2.84	1.71	4

Table 6 Diagnostic molecular genetic characters for nine revised leopard subspecies

Geographic Group	Revised Subspecies	mtDNA haplotypes	mtDNA sites ¹	Microsatellite alleles
Africa	<i>Panthera pardus pardus</i> * (I + II)	Pan, Sua, Rei, Sho1, Sho2, Sho3, Sho4, Sho5, Sho6, Sho7, Sho8, Sho9	None, or [252, 294, 318] ³ , or [87, 147, 279] ⁴	FCA008–122, –128, –130, –132; FCA026–114, –150; FCA043–116, –118, –120, –122, –124, –126; FCA075–131; FCA077–127, –141, –143, –145, –147; FCA090–123; FCA094–203; FCA096–183, –187, –189; FCA098–98, –102, –118, 124, –128, –132; FCA105–202, –206, 208; FCA123–127, –133, –153; FCA126–153, –157, –159, –161, –165; FCA139–130, –144, –150; FCA161–163, –187; FCA211–124, –126; FCA220–198, –206; FCA224–177; FCA229–155; FCA247–127, –137; FCA310–125, –127; FCA441–126; FCA453–169, –173; FCA678–212, –218, –222, –234
Arabia	<i>P. p. nimr</i> *	Nim	75, 240, 303, 327, 435	N. D.
Central Asia	<i>P. p. saxicolor</i>	Sax1, Sax2, Sin	23 ²	FCA097–124; FCA310–113
India	<i>P. p. fusca</i>	Fus1, Fus2, Fus3, Fus4, Fus5, Fus6	276	FCA026–124; FCA096–217; FCA098–116; FCA224–179; FCA229–169; FCA310–105; FCA391–190; FCA453–177
Sri Lanka	<i>P. p. kotiya</i>	Kot1, Kot2, Kot3	21	FCA008–150; FCA075–115; FCA096–177; FCA139–136, FCA247–137
South China	<i>P. p. delacouri</i>	Del1, Del2, Del3	23	FCA043–106; FCA-075–117; FCA090–105; FCA098–96; FCA224–157, –175; FCA391–222
North China	<i>P. p. japonensis</i>	Jap1, Jap2	None	FCA008–156; FCA026–148; FCA391–222; FCA441–134
Russian Far East	<i>P. p. orientalis</i>	Ori1, Ori2	8, 187	FCA026–148; FCA441–158
Java	<i>P. p. melas</i> *	Mel	168, 219, 249	None

¹Sites listed relative to beginning of *NADH-5* gene; 1 corresponds to 12634 in the complete domestic cat mtDNA sequence (Lopez *et al.* 1996).²From the beginning of sequenced portion of the Control Region (116 bp).³Together with *P. p. nimr*.⁴Together with *P. p. nimr* and *P. p. melas*.

*Subspecies that are considered to be as tentative under the present analysis.




Fig. 5 Linearized tree derived from Neighbour-joining tree topology constructed based on *NADH-5* mtDNA sequences (611 bp) and Kimura 2-Parameter α -corrected genetic distances ($\alpha = 0.90$ estimated by PAML). The scale represents the time scale in Ma (see text).

test did not reveal significant rate heterogeneity among leopard sequences (confidence probability; $CP < 95\%$); with the branch-length test two sequences (Del 1 and Del 2) have evolved faster than the rest (for both $CP = 95\%$). However, because these two sequences were only marginally deviated in the branch-length test, they were not excluded, and thus all sequences were used to construct a linearized tree (Fig. 5). Two dates were estimated, the divergence time of the first major node (node 2), and the time of origin of Asian leopards (node 3). Using the equation, $H = \mu T$, H was equal to the linearized

heights estimated by the two-cluster test, and μ was a substitution rate estimated based on known fossil dates (T). When T was set to be equal to 3.5 million years (Myr) as the time when leopards and lions split (node 1), μ was estimated to be 0.0142 per site per Myr, or approximately 1.4% per Myr. Applying this substitution rate, extant leopard lineages were estimated to have diverged approximately 0.825 ± 0.178 Ma and the age of Asian leopards was estimated to be about 0.297 ± 0.086 Ma. Using 2 Ma as a calibration point between leopards and lions, μ was estimated to be 0.024 per site per Myr, or approximately 2.5% per Myr. According to this substitution rate, modern leopards originated about 0.471 ± 0.102 Ma and Asian leopards at about 0.169 ± 0.049 Ma.

Discussion

A population genetic and phylogeographic assessment of leopards sampled from specific geographical origins throughout their current range was determined for 77 leopards representing 13 of 27 named classical subspecies. Using DNA sequence data from two mitochondrial gene segments, *NADH-5* and CR, and composite microsatellite genotypes for 25 microsatellite loci, evidence for nine revised subspecies was obtained (Fig. 1 and Table 6). Recognition of the nine revised subspecies was derived from phylogenetic analyses of mtDNA haplotypes (ME, MP and ML algorithms, Fig. 2) plus statistically significant distinct F_{ST} measures (Table 4). The mtDNA distinctions were supported by phylogenetic analysis of composite microsatellite genotypes that assorted individuals into the same groups (Fig. 3). Seven revised subspecies for which there were four or more representatives (all except *Panthera pardus melas* and *P. p. nimr*) consistently showed genetic differentiation based on R_{ST} estimates for microsatellite allele distributions among the population (Tables 3 and 4). Diagnostic mtDNA sites, haplotypes or subspecies specific microsatellite alleles form the basis for recognition of the nine subspecies (Table 6). The results affirm and extend the provisional subspecies categories proposed earlier (Miththapala *et al.* 1996), based on other molecular evolutionary markers.

The two revised subspecies with the fewest sampled individuals, *P. p. melas* ($n = 2$) and *P. p. nimr* ($n = 1$), should be considered as tentative at this stage, although highly distinctive mtDNA haplotypes were observed in multiple individuals from these regions. Populations of *P. p. nimr* appear to have been isolated for quite a long time, accumulating multiple diagnostic sites that distinguish it from any other subspecies (Table 2). Presently, not more than 200 leopards are thought to be left in the whole Arabian peninsula (Lagrot & Lagrot 1999).

Sampling in central, west and northern Africa was limited (one *P. p. suahelicus*, one *P. p. reichenowi*, and one

P. p. panthera, Fig. 1 and Table 1). These isolated individuals did not discriminate from the more numerous southern African samples (Figs 2 and 3). More extensive sampling in the future may reveal further partitions among north/central African leopards. The occurrence of divergent mtDNA haplotype lineages (PARI and PARII, Fig. 2) may reflect ancestral subdivisions consistent with this possibility. Nonetheless, until better evidence is developed, we consider African leopards to comprise a single revised subspecies, *P. p. pardus*.

In east Asia, we found significant differentiation and diagnostic markers that support the recognition of *P. p. orientalis*, *P. p. japonensis* and *P. p. delacouri*. Miththapala *et al.* (1996) had correctly deferred judgement on these three subspecies due to inadequate sampling, which is remedied here. The island population of Sri Lanka, *P. p. kotiya*, was distinctive, but was closely aligned to the mainland India subspecies *P. p. fusca*. This similarity is likely due to an historic origination of *P. p. kotiya* from *P. p. fusca* founders. The Javan leopard *P. p. melas* was highly distinctive from other Asian leopards for evolutionary reasons that remain uncertain (see Miththapala *et al.* 1996).

Genetic diversity

The amount of genetic diversity revealed in the leopards was comparable to or higher than those reported for other cat species, such as lions, cheetahs (Driscoll 1998), jaguars (Eizirik *et al.* 2001), and pumas (Culver *et al.* 2000). The genetic variation in leopards, however, varied significantly across their geographical range. There may be bias in estimation of actual genetic diversity among leopard populations due to different and sometimes insufficient sample sizes. Three populations (*P. p. saxicolor*, *P. p. japonensis* and *P. p. delacouri*) were represented only by captive-bred individuals, and thus, the reported genetic estimates reflect the status of captive populations of these revised subspecies.

The African leopards were the most genetically variable among all leopard subspecies by both mtDNA sequences and microsatellites (Table 5). Similar high diversity has been reported for Tanzanian leopards (Spong *et al.* 2000). The lowest level of genetic variation in both types of markers was observed in the Far Eastern leopard, *P. p. orientalis*. This population has a documented history of demographic and range reduction, and it is the most critically endangered leopard subspecies (Miquelle *et al.* 1996; Nowell & Jackson 1996; Uphyrkina *et al.*, unpublished data). Sri Lankan leopards, *P. p. kotiya*, which had previously been reported as showing diminished genetic diversity with several genetic metrics (Miththapala *et al.* 1991, 1996), also showed relatively low levels of microsatellite variation (Table 5). The mtDNA diversity was somewhat low in *P. p. saxicolor* (mean number of pairwise differences was 0.50 and nucleotide diversity was 0.07%). This may be explained by the sampling

of only captive individuals which are thought to be substantially inbred (A. Shoemaker, personal communication).

Radiation of modern leopards

African leopards were the first group to split off from the phylogenetic tree based on the microsatellite data (Fig. 4) and they were also basal relative to Asian leopards in the MP phylogenetic tree with mtDNA data (Fig. 2). African leopards possessed the broadest range of genetic variation by all molecular genetic techniques applied to date: allozymes, mtRFLP, minisatellites (Miththapala *et al.* 1996), microsatellites and mtDNA (Table 5). Further, *P. p. pardus* share more mitochondrial sites in common with outgroup species than other subspecies (Table 2). We interpret these observations as indicative of an African origin for leopard genetic diversity which we estimate as between 470 000 and 825 000 years ago depending on which fossil calibration dates were employed (Fig. 5).

The Asian lineages are estimated as somewhat younger, between 170 000 and 300 000 years ago, consistent with a migration out-of-Africa to the middle east and east to eastern Asia during that interval. The leopard may have had to cross the Afro-Arabian landbridge, perhaps by the Egyptian-Sinai-Israeli passageway, following the invasion of many sub-Saharan animal and plant forms into the eastern Mediterranean and Eurasia during late Pliocene and early Pleistocene (Tchernov 1988). The leopard's migration would correspond precisely in time with the postulated migration of modern human populations out-of-Africa similarly deduced from patterning of mitochondrial and nuclear genomic diversity (Hedges *et al.* 1992; Nei & Roychoudhury 1993; Bowcock *et al.* 1994; Goldstein *et al.* 1995; Calafell *et al.* 1998; Ingman *et al.* 2000).

Based on fossil dating of the earliest members of *Panthera*, Hemmer (1976) proposed that first the jaguar-like ancestor of this subgenus spread over Africa, Europe, southern and northern Asia and North America in the middle Lower Pleistocene. Then differentiation into the living species took place as a second stage somewhere in Lower Pleistocene. The ancestral leopard may have gone extinct during faunal turnovers throughout the world except in Africa, and the modern leopard may then have spread out of Africa again. The leopard appears to have taken the same routes that were used by modern human migrations (Hedges 2000).

Conclusions

In the present paper we have confirmed and extended the phylogenetic discrimination of seven phylogeographic groups of leopards to nine revised subspecies, one African, *Panthera pardus pardus*, and eight Asian subspecies, *P. p. saxicolor*, *P. p. fusca*, *P. p. kotiya*, *P. p. melas*, *P. p. delacouri*,

P. p. japonensis, and *P. p. orientalis*. However, we suggest that further analyses involving more extensive sample collection, particularly to examine genetic distinctiveness among African and central Asian leopard populations, are required. We have determined levels of genetic diversity within the leopard subdivisions from analysis of mtDNA sequences and microsatellite data, and estimated an approximate age of modern leopard lineages, based on mtDNA haplotype divergence and a fossil-record calibration. Genetic information provided here, accompanied with ecological and ecosystem approaches, may be useful in setting priorities and developing management strategies for leopard subspecies recognition and conservation.

Acknowledgements

We would like to thank all people listed in the Table 1 who provided the leopard samples. We are grateful to E. Eizirik, T. Sitnikova, J. Martenson, C. Driscoll, M. Menotti-Raymond, W. Murphy, M. Culver, A. Roca, V. David, S. Cevario, A. Snyder, E. Frazer, A. Shoemaker and J. Pecon-Slaterry for useful discussions, suggestions and technical assistance. All tissue samples were collected in full compliance with specific Federal Fish and Wildlife permits [Conservation on International Trade in Endangered Species of Wild Fauna and Flora (CITES); Endangered and Threatened Species] issued to the National Cancer Institute, National Institutes of Health (principal officer S. J. O'Brien) by the U. S. Fish and Wildlife Service of the Department of the Interior.

References

- Avice JC, Ball RM (1990) Principles of genealogical concordance in species concepts and biological taxonomy. *Oxford Surveys in Evolutionary Biology*, **7**, 45–67.
- Bowcock AM, Ruiz-Linares A, Tomfohrde J, Minch E, Kidd JR, Cavalli-Sforza LL (1994) High resolution of human evolutionary trees with polymorphic microsatellites. *Nature*, **368**, 455–457.
- Bull JJ, Huelsenbeck JP, Cunningham CW, Swofford DL, Waddell PJ (1993) Partitioning and combining data in phylogenetic analysis. *Systematic Biology*, **42**, 384–397.
- Calafell F, Shuster A, Speed WC, Kidd JR, Kidd KK (1998) Short tandem repeat polymorphism evolution in humans. *European Journal of Human Genetics*, **6**, 38–49.
- Christen Y (2000) *Le Peuple Leopard*, Editions. Michalon, Paris.
- Culver M, Johnson WE, Pecon-Slaterry J, O'Brien SJ (2000) Genomic ancestry of the American puma (*Puma concolor*). *Journal of Heredity*, **91**, 186–197.
- Driscoll CA (1998) *A characterization of microsatellite loci variation in Panthera leo, Acinonyx jubatus and Felis concolor*. MSc Thesis, Hood College, Frederick MD.
- Eizirik E, Kim J, Crawshaw J, O'Brien SJ, Johnson WE (2000) Phylogeography population history and conservation genetics of jaguars (*Panthera onca*, Mammalia, Felidae). *Molecular Ecology*, **10**, 65–79.
- Ellegren H (1995) Mutation rates at porcine microsatellite loci. *Mammalian Genome*, **6**, 376–377.
- Excoffier L, Smouse PE, Quattro JM (1992) Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics*, **131**, 479–491.
- Felsenstein J (1985a) *Phylogeny inference package (PHYLP)*, Version 3.5. University of Washington, Seattle.
- Felsenstein J (1985b) Confidence limits on phylogenies: an approach using the bootstrap. *Evolution*, **39**, 783–791.
- Goldstein DB, Ruiz Linares A, Cavalli-Sforza LL, Feldman MW (1995) Genetic absolute dating based on microsatellites and the origin of modern humans. *Proceedings of the National Academy of Sciences of the USA*, **92**, 6723–6727.
- Guo SW, Thompson EA (1992) Performing the exact test of Hardy–Weinberg proportion for multiple alleles. *Biometrics*, **48**, 361–372.
- Hasegawa M, Kishino H, Yano T (1985) Dating of the human-ape splitting by a molecular clock of mitochondrial DNA. *Journal of Molecular Evolution*, **22**, 160–174.
- Hedges SB (2000) Human evolution. A start for population genomics. *Nature*, **408**, 652–653.
- Hedges SB, Kumar S, Tamura K, Stoneking M (1992) Human origins and analysis of mitochondrial DNA sequences. *Science*, **255**, 737–739.
- Hemmer H (1976) Fossil History of Living Felidae. In: *The World's Cat* (ed. Eaton RL), pp. 1–14. Carnivore Research Institute, Burke Museum, Seattle.
- Hillis DM, Bull JJ (1993) An empirical test of bootstrapping as a method for assessing confidence in phylogenetic analysis. *Systematic Biology*, **42**, 182–192.
- Hoelzel AR, Halley J, O'Brien SJ *et al.* (1993) Elephant seal genetic variation and the use of simulation models to investigate historical population bottlenecks. *Journal of Heredity*, **84**, 443–449.
- Hoogerwerf A (1970) *Udjung Kulon: the Land of the Last Javan Rhinoceros*. E.J.Brill, Leiden.
- Huelsenbeck JP, Bull JJ, Cunningham CW (1996) Combining data in phylogenetic analysis. *Trends in Ecology and Evolution*, **11**, 152–157.
- Ingman M, Kaessmann H, Paabo S, Gyllenstein U (2000) Mitochondrial genome variation and the origin of modern humans. *Nature*, **408**, 708–713.
- Johnson WE, Culver M, Iriarte JA, Eizirik E, Seymour KL, O'Brien SJ (1998) Tracking the evolution of the elusive Andean mountain cat (*Oreailurus jacobita*) from mitochondrial DNA. *Journal of Heredity*, **89**, 227–232.
- Johnson WE, O'Brien SJ (1997) Phylogenetic reconstruction of the Felidae using 16S rRNA and NADH-5 mitochondrial genes. *Journal of Molecular Evolution*, **44**, S98–S116.
- Kimura M (1980) A simple method for estimation of evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution*, **16**, 111–120.
- Kitchener A (1991) *Natural History of Wild Cats*. Comstock Publishing Associates, Ithaca, New York.
- Kumar S, Tamura K, Nei M (1993) *MEGA, Version 1.01*. The Pennsylvania State University, University Park, PA.
- Lagrot I, Lagrot JF (1999) Leopard in the Arabian Peninsula. *Cat News*, **30**, 21–22.
- Liedloff A (1999) *MANTEL 2*. <http://sci.qut.edu.au/nrs/mantel.htm>.
- Lopez JV, Cevario S, O'Brien SJ (1996) Complete nucleotide sequences of the domestic cat (*Felis catus*) mitochondrial genome and a transposed mtDNA tandem repeat (*Numt*) in the nuclear genome. *Genomics*, **33**, 229–246.
- Menotti-Raymond M, David VA, Lyons LA *et al.* (1999) A genetic linkage map of microsatellites in the domestic cat (*Felis catus*). *Genomics*, **57**, 9–23.
- Minch E, Ruiz-Linares A, Goldstein DB (1995) *MICROSAT*. <http://lotka.stanford.edu/microsat.html>.

- Miquelle D, Arzhanova T, Solkin V (1996) *A Recovery Plan for Conservation of the Far Eastern Leopard*. USAID Russian Far Eastern EPT Project, Vladivostok, Russia.
- Miththapala S, Seidensticker J, O'Brien SJ (1996) Phylogeographic subspecies recognition in leopards (*Panthera pardus*): molecular genetic variation. *Conservation Biology*, **10**, 1115–1132.
- Miththapala S, Seidensticker J, Phillips LG *et al.* (1991) Genetic Variation in Sri Lankan Leopards. *Zoo Biology*, **10**, 139–146.
- Nei M, Roychoudhury AK (1993) Evolutionary relationships of human populations on a global scale. *Molecular Biology and Evolution*, **10**, 927–943.
- Nei M, Tajima F, Tateno Y (1983) Accuracy of estimated phylogenetic trees from molecular data. II. Gene frequency data. *Journal of Molecular Evolution*, **19**, 153–170.
- Nei M, Takezaki N (1996) The root of the phylogenetic tree of human populations. *Molecular Biology and Evolution*, **13**, 170–177.
- Nowell K, Jackson P (1995) New Red List Categories for Wild Cats. *Cat News*, **23**, 21–27.
- Nowell K, Jackson P (1996) *Wild Cats*. IUCN/SSC Cat Specialist Group, Gland, Switzerland.
- O'Brien SJ, Collier G, Benveniste RE (1987) Setting the molecular clock in Felidae: The great cats: *Panthera*. In: *Tigers of the World* (ed. Tilson R), pp. 10–27. Noyes Publications, Park Ridge, N.J.
- O'Brien SJ, Mayr E (1991) Bureaucratic mischief: recognizing endangered species and subspecies. *Science*, **251**, 1187–1188.
- Ota T (1993) *DISPAN: Genetic Distances and Phylogenetic Analysis*. Institute of Molecular Evolutionary Genetics, Pennsylvania State University, University Park, Pennsylvania.
- Pocock RI (1930) The panthers and ounces of Asia. *Journal of Bombay Natural History Society*, **XXXIV**, 63–82, 307–336.
- Pocock RI (1932) The leopards of Africa. *Proceeding of Zoological Society of London*. 543–591.
- Raymond M, Rosset F (1995) *GENEPOP* (Version 1.2): population genetics software for exact test and ecumenicism. *Journal of Heredity*, **86**, 248–249.
- Rice WR (1989) Analysing tables of statistical tests. *Evolution*, **43**, 223–225.
- Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution*, **4**, 406–425.
- Sambrook J, Fritsch E, Maniatis T (1989) *Molecular Cloning: a Laboratory Manual*. Cold Spring Harbor Laboratory Press, New York.
- Schneider S, Kueffer JM, Roessli D, Excoffier L (1997) *ARLEQUIN 1.1*. University of Geneva, Switzerland.
- Seidensticker J, Lumpkin S (1991) *Great Cats*. Merehurst, London.
- Slatkin M (1995) A measure of population subdivision based on microsatellite allele frequencies. *Genetics*, **139**, 457–462.
- Spong G, Johansson M, Bjorklund M (2000) High genetic variation in leopards indicates large and long-term stable effective population size. *Molecular Ecology*, **9**, 1773–1782.
- Swofford DL (1998) *'PAUP* Phylogenetic Analysis Using Parsimony and Other Methods' Computer Program*. Sinauer, Sunderland, MA.
- Takezaki N, Nei M (1996) Genetic distances and reconstruction of phylogenetic trees from microsatellite DNA. *Genetics*, **144**, 389–399.
- Takezaki N, Rzhetsky A, Nei M (1995) Phylogenetic test of the molecular clock and linearized trees. *Molecular Biology and Evolution*, **12**, 823–833.
- Tchernov E (1988) The biogeographical history of the southern Levant. In: *The Zoogeography of Israel* (eds Yom-Tov Y, Tchernov E), pp. 159–250. Dr W Junk Publishers, Dordrecht/Boston/Lancaster.
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997) The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research*, **25**, 4876–4882.
- Turner A, Anton M (1997) *The Big Cats and Their Fossil Relatives*. Columbia University Press, New York.
- Wayne RK, Benveniste RE, Janczewski DN, O'Brien SJ (1993) Molecular and biochemical evolution of the Carnivora. In: *Carnivore Behavior, Ecology, and Evolution* (ed. Gittleman JL), pp. 465–495. Chapman & Hall, London.
- Wentzel J, Stephens JC, Johnson WE (1999) Subspecies of tigers: Molecular assessment using 'voucher specimens' of geographically traceable individuals. In: *Riding the Tiger: Tiger Conservation in Human-Dominated Landscapes* (eds Seidensticker J, Christie S, Jackson P), pp. 40–49. Cambridge University Press, Cambridge.
- Yang Z (1999) *Phylogenetic Analysis by Maximum Likelihood (PAML)*. University College of London, London.
- Yang Z, Kumar S (1996) Approximate methods for estimating the pattern of nucleotide substitution and the variation of substitution rates among sites. *Molecular Biology and Evolution*, **13**, 650–659.

After receiving degrees in Environmental Sciences and Policy from Moscow College and Gene Engineering and Biotechnology from Moscow State University, Olga worked as a geneticist for the Hornocker Wildlife Institute and as a visiting scientist in the Laboratory of Genomic Diversity of the National Cancer Institute. Olga remains actively involved in conservation efforts of the Far Eastern (Amur) leopard and Siberian tiger in her homeland, the Russian Far East.
