

## Rb-Sr AGES OF GRANITOIDS IN THE REHOBOTH-NAUCHAS AREA, SOUTH WEST AFRICA/NAMIBIA

D.L. Reid<sup>1</sup>, S. Malling<sup>2</sup> and H.L. Allsopp<sup>3</sup>

<sup>1</sup> Department of Geochemistry, University of Cape Town, Rondebosch 7700.

<sup>2</sup> Precambrian Research Unit, Department of Geology, University of Cape Town, Rondebosch 7700.

<sup>3</sup> Bernard Price Institute of Geophysics, University of the Witwatersrand, Johannesburg 2000. (Deceased)

### ABSTRACT

Rb-Sr ages have been obtained for several pre-Damaran granitoid suites that intrude various volcano-sedimentary sequences in the Rehoboth region of South West Africa/Namibia. Two major periods of granitoid emplacement are indicated by the results: (1) an older 1 800–1 600 Ma event so far only recognized in a small part of the region near Rehoboth, and (2) a younger 1 200–1 000 Ma event which is widespread throughout the region and represents a major magmatic episode, at least in terms of volume of products involved. The older granitoids include the Naub Diorite ( $1\,725 \pm 52$  Ma) and the Swartmodder Granite ( $1\,639 \pm 25$  Ma), while the Biesiepoort Granite ( $1\,222 \pm 45$  Ma), Piksteel Granodiorite ( $1\,170 \pm 20$  Ma) and Gamsberg Granite ( $1\,079 \pm 25$  Ma) represent members of the voluminous younger granitoids. The Weener Quartz Diorite may also be part of the younger granitoid event (dated at  $1\,207 \pm 170$  Ma), but an older age of  $\sim 1\,800$  Ma cannot be ruled out. Agreement with published zircon U-Pb ages is variable, but detailed comparison is precluded by the incomplete nature of the published zircon data. Initial Sr isotope ratios indicate that the older granitoids may have been derived from pre-existing crust not much older than  $\sim 2\,000$  Ma, and that the younger granitoids showed increasing involvement of older continental crust with time.

### 1. INTRODUCTION

Attempts to elucidate the pre-Damaran geological history of South West Africa have often been based on unravelling the complex stratigraphic relations in the Rehoboth region. As a contribution to the timing of pre-Damaran igneous, sedimentary and metamorphic events in this important region, several granitoid suites have been investigated using the Rb-Sr method. Where possible the samples have been collected from localities showing the least equivocal field relations, which has often resulted in restricting the study to small areas. This has been especially so in the Nauchas area for example (Fig. 1).

The most recent stratigraphic analysis of the pre-Damaran in the Rehoboth region was published in SACS (1980), and has been followed in the compilation of the 1:250 000 geological sheet (Schalk, unpubl.). The stratigraphy presented in Table 1 contains units formally recognized and proposed, but not in the temporal order indicated by the age data presented in this paper.




Fig. 1: General locality map of the Rehoboth region, showing major roads and the areas covered by the detailed locality maps.

TABLE 1: Pre-Damaran stratigraphy of the Rehoboth region. Based on SACS (1980), Borg and Maiden (1986) and Schalk (unpubl.)

VOLCANO-SEDIMENTARY UNITS		INTRUSIVE UNITS
Klein Aub Formation	Sinclair Group	Gamsberg Granite Suite
Doornpoort Formation		
Grauwater Formation		
Dordabis Formation		
Nükopf Formation		
Gaub Valley Formation	Rehoboth Group	Piksteel Granodiorite Suite
Billstein Formation		
Marienhof Formation		
Elim Formation	Pre-Rehoboth Sequences	Naub Diorite
Neuhof Formation		
Moorivier Formation		
		Weener Quartz Diorite
		Alberta Complex

## 2. RESULTS

Analytical data obtained during the whole rock Rb-Sr study appear in Table 2 and the general locality of samples collected is shown in Figs 1–5. Rb and Sr concentrations




Fig. 2: Simplified geologic map of the Biesiepoort–Naub area, showing sample locations for the Naub Diorite (1) and Biesiepoort Granite (4). Geology on this map and on Figs 4 and 5 after Schalk (unpubl.).




Fig. 3: Detailed locality map of the southern part of the Rehoboth townlands, showing sample locations of the Swartmodder Granite (2) and Piksteel Granodiorite (6).




Fig. 4: Geologic map of the Gamsberg–Höhenheim area showing sample locations for the Gamsberg pluton of the Gamsberg Granite Suite.

were determined by XRF at UCT using techniques similar to those described in Duncan *et al.* (1984). Sr isotope ratios were measured by mass spectrometry at the BPI using standard techniques (e.g. Allsopp *et al.*, 1979). A summary of Rb-Sr isochron ages appears in Table 3, together with details of regression procedures adopted. Rb-Sr isochrons have been distinguished from errorchrons using a cut-off value of 2,5 for the MSUM value (York, 1969; Brooks *et al.*, 1972).

TABLE 2: Rb and Sr data obtained during the present study

	Rb	Sr	Rb/Sr	<sup>87</sup> Rb/ <sup>86</sup> Sr	<sup>87</sup> Sr/ <sup>86</sup> Sr
<b>NAUB DIORITE</b>					
SM7630C	26,4	269	0,0981	0,283	0,71108
SM7630G	23,8	217	0,1098	0,317	0,71222
SM7630A	29,1	215	0,135	0,390	0,71364
SM7630	54,0	188	0,288	0,833	0,72500
NaDiBi	49,5	190	0,260	0,753	0,72261
<b>SWARTMODDER GRANITE</b>					
SM1236	115	186	0,618	1,793	0,75026
SM1172	58,3	118	0,494	1,43	0,74160
SM2193	81,1	143	0,566	1,642	0,74600
SM1083	98,3	547	0,19	0,521	0,72030
SM1135	99,2	107	0,925	2,689	0,77230
SM1050	120	116	1,037	3,011	0,77870
<b>ALBERTA MAFIC COMPLEX</b>					
SM7640C	34,7	271	0,128	0,370	0,70942
SM7642	192	632	0,304	0,879	0,72009
SM7642B	0,73	679	0,00108	0,003	0,70190
SM7642C	<0,2	634	—	—	0,70186
<b>BIESIEPOORT GRANITE</b>					
Bi-1	129	295	0,437	1,264	0,72720
Bi-2	158	242	0,654	1,893	0,73780
Bi-3	143	301	0,475	1,375	0,72846
SM76125	140	161	0,874	2,533	0,74910
SM76126	92,0	290	0,317	0,917	0,71996
SM76127	81,0	277	0,292	0,845	0,71923
SM76128	186	105	1,774	5,166	0,79303
<b>WEENER QUARTZ DIORITE</b>					
SM76146	47,9	781	0,060	0,176	0,70720
SM7651A	65,3	761	0,0858	0,247	0,70836
SM7651B	72,0	754	0,0955	0,280	0,70896
SM7651B2	72,7	729	0,0991	0,289	0,70920
SM7651D	69,5	739	0,094	0,275	0,70890
<b>PIKSTEEL GRANODIORITE</b>					
SM76-9	262	107	2,44	7,13	0,82740
SM76-8II	164	116	1,417	4,12	0,77918
SM76-8I	162	111	1,463	4,25	0,78180
SM76-5	202	451	0,449	1,30	0,73124
SM76154	162	134	1,21	3,49	0,76720
<b>GAMSBURG GRANITE</b>					
Nauchas Pluton					
SM064	285	26,9	10,59	32,05	1,19910
SMAG41	308	30,3	10,16	30,43	1,18430
SMAG42	289	34,1	8,476	25,4	1,09110
SM7641	208	279	0,745	2,16	0,74140
Namibgrens Pluton					
SM066	284	40,5	7,022	20,90	1,03000
SM7635	158	217	0,728	2,11	0,73960
SM7636	173	200	0,866	2,51	0,74690
SM76117	160	142	1,08	3,13	0,75800
SM76120	154	155	0,995	2,89	0,75350
Gamsberg Pluton					
SM76125G	126	149	1,18	3,42	0,76192
SM76123	327	16,1	20,34	64,23	1,66310
SM76124	310	11,6	26,63	86,31	1,89630
SM76125a	144	148	0,972	2,884	0,75194
SM76123A	299	12,9	23,29	73,50	1,70310
<b>POST-GAMSBURG GRANITE MAFIC DYKES</b>					
SM76130D2	8,93	360	0,0256	0,072	0,70750
AD21	10,6	191	0,0553	0,160	0,70868
AD22	21,3	342	0,962	0,180	0,70916
NAMD2	9,4	206	0,0457	0,132	0,70834
<b>POST-GAMSBURG GRANITE ACID DYKES</b>					
P2A	17,8	140	0,124	0,369	0,73641
P2B	30,0	123	0,245	0,709	0,74112
SRM16	146	106	1,378	4,005	0,77350
76118	69,1	43,9	1,574	4,595	0,79401
7639	181	220	0,824	2,390	0,77141

TABLE 3: Summary of Rb-Sr ages obtained during the present study

ROCK TYPE	AGE (Ma)	$R_o$	MSUM (N)
Post-Gamsberg Granite Mafic Dykes	1 030 ± 185	0,7064 ± 4	0,71 (4)
Gamsberg Granite (Gamsberg Pluton)	984 ± 55	0,7123 ± 39	21,04 (5)
Gamsberg Granite (Nauchas Pluton)	1 073 ± 25	0,7082 ± 16	3,48 (4)
Gamsberg Granite (Namibgrens Pluton)	1 096 ± 62	0,7075 ± 25	6,87 (5)
Gamsberg Granite (Combined Nauchas/ Namibgrens Plutons)	1 079 ± 25	0,7081 ± 12	4,39 (9)
Piksteel Granodiorite	1 170 ± 20	0,7095 ± 7	2,21 (5)
Weener Quartz Diorite	1 207 ± 170	0,7041 ± 6	0,16 (5)
Biesiepoort Granite	1 222 ± 45	0,7044 ± 9	3,82 (7)
Biesiepoort/Weener	1 238 ± 13	0,70406 ± 13	2,16 (7)
Alberta Mafic Complex	1 442 ± 32	0,70183 ± 14	0,23 (4)
Swartmodder Granite	1 639 ± 25	0,7080 ± 4	1,30 (6)
Naub Diorite	1 725 ± 52	0,7041 ± 3	2,03 (5)

Ages calculated using regression method of York (1969). Error assigned to  $^{87}\text{Rb}/^{86}\text{Sr} = 1,0\%$ ; to  $^{87}\text{Sr}/^{86}\text{Sr} = 0,015\%$ . Regression errors are two sigma where the MSUM < 2,5 and augmented by SQRT (MSUM) when MSUM > 2,5. Decay constant and isotopic ratios after Steiger and Jäger (1977).




Fig. 5: Geologic map of the Nauchas area showing sample locations of the Alberta basic complex (3), Weener Quartz Diorite (5), Nauchas pluton (7) and the Namibgrens pluton of the Gamsberg Granite Suite (8), basic dykes (10) and acid quartz porphyry dykes (11) cutting the Gamsberg Granite. Major faults are labelled with 'f'.

## 2.1 Naub Diorite

The Naub Diorite was sampled where this rock type forms a body straddling the boundary between Naub 274 and Biesiepoort 275 (23°06'S, 17°21'E, Fig. 2). The rock is a mesocratic hornblende-biotite gneiss with an intense, near vertical foliation striking mainly south-east, but which swings nearly east-west, dipping steeply north where it is intruded by the younger Biesiepoort Granite (Schalk, unpubl.). The intense foliation in the Naub Diorite is shared by the Elim Formation metasediments and metavolcanics in this locality, and therefore suggests a pre- to early syntectonic intrusive relationship (Hoffmann, pers. comm., 1988). Contacts are diffuse but the diorite is full of rafts of Elim rocks. Inclusions of hornblende and melagabbro also occur in the Naub Diorite, and may be derived from a large ultramafic body nearby, which has also intruded the Elim Formation. Five samples of

the Naub Diorite define a Rb-Sr isochron (MSUM = 2,03, Fig. 6) with an age of  $1725 \pm 52$  Ma and an initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio of  $0,7041 \pm 3$ .




Fig. 6: Rb-Sr isochron diagram for the Naub Diorite (granodiorite gneiss). Labels on this and all subsequent Rb-Sr diagrams are abbreviated sample numbers.

## 2.2 Swartmodder Granite

The Swartmodder Granite does not feature in the stratigraphic column published in SACS (1980), and appears to have been included within the Piksteel Granodiorite (Schalk, unpubl.). The granite intrudes the Marienhof Formation in the southern portion of the Rehoboth townlands (23°28'S, 17°04'E, Fig. 3), just west of the main highway (N1/4). Petrographically the Swartmodder Granite is leucocratic and shows a variably developed foliation, which makes it difficult to distinguish from other similar granitoids that occur throughout the region. A pluton of the Piksteel Granodiorite Suite has intruded the Swartmodder Granite. Samples of Swartmodder Granite used in this study were obtained from Falconbridge Ltd., Windhoek, who provided fresh unweathered drill core. Six samples define a Rb-Sr isochron (MSUM = 1,30, Fig. 7), with an age of  $1639 \pm 25$  Ma and an initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio of  $0,7080 \pm 4$ .




Fig. 7: Rb-Sr isochron diagram for the Swartmodder Granite.

## 2.3 Alberta mafic complex

The mafic to ultramafic Alberta complex has intruded the Elim Formation, but has in turn been intruded by the Piksteel and Gamsberg granitoids. Fresh material suitable for Rb-Sr dating purposes is difficult to collect and the results reported here are only preliminary. Two anorthosites and two

metagabbros sampled from the complex on Alberta (23° 24'S, 16°20'E, Fig. 5) define a Rb-Sr isochron (MSUM 0,23, Fig. 8), with an age of  $1442 \pm 32$  Ma and an initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio of  $0,70183 \pm 14$ .




Fig. 8: Rb-Sr isochron diagram for the Alberta Complex.

#### 2.4 Biesiepoort Granite

The Biesiepoort Granite was sampled on Biesiepoort 275 (23°06'S, 17°21'E) where it occurs as an adamellite granite, weakly foliated in places, which has intruded the Naub Diorite (Fig. 2). Contacts are knife sharp and cut across the intense foliation in the Naub Diorite (Hoffmann, pers. comm., 1988). Seven samples define a Rb-Sr errorchron (MSUM = 3,82, Fig. 9), with an age of  $1222 \pm 45$  Ma and an initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio of  $0,7044 \pm 9$ .




Fig. 9: Rb-Sr isochron diagram for the Biesiepoort Granite.

#### 2.5 Weener Quartz Diorite

The Weener Quartz Diorite was sampled from a pluton (consisting mainly of tonalite) on Namibgrens 154 (23°37'S, 16°14'W, Fig. 5) just south of an intrusive contact with a younger granitoid (Namibgrens pluton of the Gamsberg Granite Suite). Further south-west this Weener tonalite body is complexly intruded by the Piksteel Granodiorite. The locality on Namibgrens 154 is not the type locality of the Weener Quartz Diorite, which is further north on Weener 193, but the lack of pronounced foliation made the Namibgrens pluton a better candidate for the Rb-Sr study. Numerous and characteristic mafic xenoliths occur in the outcrops sampled and despite attempts to avoid them, some incorporation may have occurred. Five samples define a Rb-Sr isochron (MSUM

= 0,16, Fig. 10), with an age of  $1207 \pm 170$  Ma and an initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio of  $0,7041 \pm 6$ . The samples all have low Rb/Sr ratios, and the range in Rb/Sr is small as well (note that the X-axis of the Rb-Sr diagram in Fig. 10 does not start at the origin), so the error in the age is very high, despite the low MSUM.

Bearing in mind that this result is statistically within error of the Biesiepoort data, combination of the two rock types on a common Rb-Sr isochron (MSUM = 2,16) yields a more precise age of  $1283 \pm 13$  Ma and initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio of  $0,70406 \pm 13$ .




Fig. 10: Rb-Sr isochron diagram for the Weener Quartz Diorite (tonalite from Namibgrens). Note that the spread in data points has been accentuated by not starting the X-axis at the origin, which was necessitated by the very small spread in Rb/Sr ratios.

#### 2.6 Piksteel granodiorite suite

A pluton mapped as part of this granitoid suite occurs in the southern part of the Rehoboth townlands (23°25'S, 17°06'E, Fig. 3). The samples were collected on the Swartmodder River slightly downstream from the Swartmodder Granite locality described earlier. The Rehoboth locality was chosen in preference to the type locality on Piksteel 209 (some 100 km to the south-west) as the field relations were clearer in that a distinct pluton of the Piksteel Suite could be seen to have intruded both the Swartmodder Granite and the Marienhof Formation. Judging from previous descriptions (e.g. De Waal, 1966) the Piksteel granitoids contain a wide variety of rock types, largely migmatitic in character, and large relatively homogeneous plutons are rare. However, subsequent workers (e.g. Schalk, unpubl.) regard the "migmatitic Piksteel rocks" as part of a different unit, the Mooirivier Metamorphic Complex. The Piksteel Granodiorite has therefore been restricted to a recognizable suite of high-level plutons. Five samples from the Rehoboth locality define a Rb-Sr isochron (MSUM = 2,21, Fig. 11), with an age of  $1170 \pm 20$  Ma and an initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio of  $0,7095 \pm 7$ .

#### 2.7 Gamsberg Granite Suite

The extensive development of the Gamsberg Granite Suite argued for the investigation of several separate plutons. The Gamsberg pluton was sampled near the highway between Windhoek and Walvis Bay (R49) on Höhenheim 25 (23°18'S, 16°24'E), and on the farm road to Gamsberg 24 (Fig. 4). Another pluton was sampled on Namibgrens 154 (23°36'S, 16°16'E, Fig. 5), while a third was sampled on Nauchas 14 (23°36'S, 16°24'E, Fig. 5). Several different Gamsberg suite granitoids occur on these latter two farms and in the case of Namibgrens, the sample locality was within the large pluton




Fig. 11: Rb-Sr isochron diagram for the Piksteel Granodiorite.

best exposed around the Drillingskuppen trigonometrical beacon. On Nauchas the sample locality was near the northern boundary with Nauams 177, just south of the road connecting the two farmhouses.

The Gamsberg pluton defined a poor Rb-Sr errorchron (MSUM = 21,04, Fig. 12) and the calculated age (see Table




Fig. 12: Rb-Sr isochron diagram for the Gamsberg pluton of the Gamsberg Granite Suite. The data for this pluton shows considerable scatter, and the isochron obtained for the other two Gamsberg suite plutons (see Fig. 13) is plotted.




Fig. 13: Rb-Sr isochron diagram for the Namibgrens and Nauchas plutons of the Gamsberg Granite Suite.

3) may have little significance. Better results were obtained for the other two plutons, with the Namibgrens data defining a Rb-Sr errorchron (MSUM = 6,87) with an age of  $1096 \pm 62$  Ma and initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio of  $0,7075 \pm 25$ . The Nauchas pluton yielded a Rb-Sr errorchron (MSUM = 3,48) with an age of  $1073 \pm 25$  Ma and initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio of  $0,7082 \pm 16$ . The Namibgrens and Nauchas results agree within error and their combination yields a Rb-Sr errorchron (MSUM = 4,39, Fig. 13) with an age of  $1079 \pm 25$  Ma and initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio of  $0,7081 \pm 12$ .

## 2.8 Post-Gamsberg granite dykes

Dykes cut the Gamsberg Granite Suite and therefore provide a minimum age for this important intrusive event. Basic dykes cutting the Namibgrens pluton ( $23^{\circ}36'S$ ,  $16^{\circ}16'E$ , Fig. 5), define a Rb-Sr isochron (MSUM = 0,71), but the very low Rb/Sr ratios preclude a precise age ( $1030 \pm 185$  Ma, Fig. 14). The initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio is  $0,7064 \pm 4$ . Acid dykes (quartz porphyries) also cut the Namibgrens pluton ( $23^{\circ}38'S$ ,  $16^{\circ}18'E$ , Fig. 5), but Rb-Sr data for these intrusives do not define an isochron. Some plot close to a  $1030$  Ma isochron, but with a substantially higher initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio.




Fig. 14: Rb-Sr isochron diagram for post-Gamsberg mafic and acid dykes.

## 3. DISCUSSION

Each granitoid suite has been studied at a pluton scale, in the sense that sampling of a particular unit was confined to a relatively small area, rather than attempting to obtain a Rb-Sr isochron from apparently similar rock types from widely spaced localities. This approach may overcome problems such as incorrect correlation, variable initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios, differences in degree of deformation and metamorphism, and hence increase the likelihood of obtaining precise isochrons. However, there remains the uncertainty in the relationship between the use of these 'single pluton' results in the identification of major multi-pluton emplacement events on a regional scale. In addition, the complex tectonic and metamorphic history experienced by the Rehoboth region may have disturbed the various radiometric systems. The following discussion of the Rb-Sr results presented here, together with comparison between these new results and published age data, must be regarded in such a light. A summary of Rb-Sr ages obtained during this study is presented in Table 3, while a compilation of published zircon U-Pb ages is given in Table 4.

### 3.1 Older granitoids

Results obtained for the Naub Diorite and Swartmodder Granite indicate Rb-Sr closure about 1800–1640 Ma ago,

TABLE 4: Published zircon U-Pb ages from the Rehoboth-Nauchas region.

LITHOLOGIC UNIT	LOCALITY	SAMPLE	AGE (Ma)	REF
GAMSBURG GRANITE SUITE	Uitdraai Oos 296	PH8	932 ± 50	2, 3
	Höhenheim 24	PH1	1 078 ± 30	2, 3
	Ganeib 61	PH6	1 150 ± 30	3
	Kanaus South 336	KES16	1 064 ± 20	3
	Kangas 371	KES2	1 104 ± 20	3
	Nauchas 14	KES15	1 178 ± 20	3
NÜKOPF FORMATION (Acid Lavas)	Auchas 347	KES138	1 770 ± 35	4
	Lekkerwater Sta		1 130 ± 75	1
	Nauzerus 11	PH2	1 010 ± 30	2, 3
	Uitdraai Oos 296	PH7	1 083 ± 30	3
	Kartatsaus 293	PH5	1 090 ± 90	2, 3
	Piksteel 209	KES117	1 076 ± 25	4
PIKSTEEL GRANODIORITE SUITE	Rehoboth Townlands	KES1	1 476 ± 30	3
BIESIEPOORT GRANITE	Biesiepoort 275	KES179	1 110 ± 30	4
MARIENHOF FORMATION (Acid Lava)	Rehoboth Townlands	KES174	1 232 ± 30	4
KAM KAM GRANITE	Kam Kam 369	KES236	986 ± 20	5
	Swartskap 332	KES177	1 406 ± 35	4
	Rehoboth Townlands	PH9	1 630 ± 50	3
SWARTMODDER GRANITE	Rehoboth Townlands	KES238	1 784 ± 45	6
NEUHOF FORMATION (Acid Lava)	Neuras 330	KES235	1 784 ± 45	6
	Olifantsvloer 453	KES237	1 328 ± 25	5

Ages recalculated to decay constants in Steiger and Jäger (1978).

Sources: 1 Burger and Coertze (1973), 2 Hugo and Schalk (1972), 3 Burger and Coertze (1975), 4 Burger and Coertze (1977), 5 Burger and Walraven (1977), 6 Burger and Walraven (1979).

which contrasts sharply with the majority of Rb-Sr ages obtained in this study, which cluster in the 1200–1000 Ma range. Only the Swartmodder Granite has been investigated by another method, and one zircon U-Pb age obtained for this granitoid is similar to that obtained from the Rb-Sr system. A second sample has a zircon U-Pb age which is slightly older. Uncertainty in the nature and location of the second zircon sample, the problem of only having available the calculated  $^{207}\text{Pb}/^{206}\text{Pb}$  age and the lack of analytical details all contribute to the difficulty of evaluating this discrepancy further. The results do, however, indicate the presence of older granitoids in the Rehoboth region which has hitherto been unrecognized. It is also likely that gneissic granitoids of Swartmodder age may occur further afield, but have been mapped as part of younger suites (e.g. Piksteel).

Supracrustal sequences intruded by the Naub and Swartmodder granitoids (e.g. Elim Formation) must therefore be older than about 1800 Ma. Direct evidence may be in the form of a zircon U-Pb age of  $1784 \pm 45$  Ma obtained from an acid lava within the Neuhof Formation (see Table 4). The results therefore suggest that this particular pre-Damara terrane does indeed contain an early-Proterozoic crustal component ( $>1800$  Ma), as has been found in other regions, such as Fransfontein (Burger *et al.*, 1976), Abbabis (Jacob *et al.*, 1978) and Haib-Vioolsdrif (Reid, 1979a, b).

### 3.2 Weener Quartz Diorite

The Weener Quartz Diorite has been studied by Seifert (1986) at the type locality on Weener 193 and the results contrast quite sharply with those obtained in this study. The Weener locality did not produce a good Rb-Sr isochron, and Seifert (1986) suggested two possible interpretations of the data, which yielded ages of  $1871 \pm 143$  Ma and  $1560 \pm 100$  Ma, depending on which data points were considered aberrant. The two plutons of the Weener Quartz Diorite have significantly different compositions, as shown by their respective Rb and Sr contents (see Fig. 15). Moreover the Rb-Sr isotopic data cannot be combined in any reasonable fashion either. The Weener Quartz Diorite has intruded the Gaub Valley Formation and is itself intruded by the Piksteel

and Gamsberg granitoids. While existing data on the latter granitoids provides a reasonable estimate of a minimum age for the Weener Quartz Diorite, no accurate estimate for the age of the Gaub Valley Formation currently exists.

The numerous mafic xenoliths found within the Weener Quartz Diorite may have introduced compositional effects, and possibly caused variations in initial  $^{87}\text{Sr}/^{86}\text{Sr}$ . As a result no Rb-Sr isochron relationship could be expected to occur amongst the samples collected. This is probably a feature of most granitoid suites, and contributes to the scatter usually found in Rb-Sr isochrons obtained for such rocks. However, the contrast shown between the Weener and Namibgrens localities looks too great to be a function only of inhomogeneous Sr isotope composition. The mafic xenoliths could be regarded as providing a maximum age for the Weener Quartz Diorite, if the source of these xenoliths were older mafic bodies like the Alberta Complex (cf. De Waal, 1966), and the Rb-Sr age for this latter complex was close to the true age. In this case the maximum age for the Weener Quartz Diorite would be  $\sim 1490$  Ma, thereby suggesting that the results reported by Seifert (1986) are anomalous.

Another possible explanation of the above discrepancy is the effect of deformation and metamorphism, since foliation is well developed at the Weener locality but nearly absent at Namibgrens. Both localities are situated very close to contacts with younger granitoids, and so contact metamorphism could have contributed to the contrasting results. However, the Rb-Sr patterns appear the reverse of those that would be predicted, in the sense that the Weener locality should have produced a younger age because of the greater degree of foliation. Contact metamorphic effects are common to both localities and therefore would not be expected to cause the difference.

A final possibility is that the Weener Quartz Diorite really was intruded 1700–1900 Ma ago, as indicated by Seifert (1986), and the result obtained in this study from Namibgrens argues against a correlation. The tonalite in this latter area rather forms a younger post-Weener intrusive, which may represent a mafic variant of either the Piksteel or Gamsberg

suites. The implication that the Weener Quartz Diorite at the type locality may in fact represent an equivalent to the Naub Diorite, clearly requires a more detailed follow-up study.

### 3.3 Biesiepoort Granite

The Biesiepoort Granite at the type locality on Biesiepoort 275 is actually a foliated adamellite, which has intruded previously deformed gneissic Naub Diorite. The initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio for the Biesiepoort Granite of  $0,7044 \pm 9$  is low and bearing in mind the average Rb/Sr ratio of the analysed samples ( $\sim 0,7$ ), a reasonable upper estimate assuming metamorphic resetting is barely outside the isochron error limits ( $\sim 50$  Ma). A zircon U-Pb age of  $1110 \pm 30$  Ma has been obtained, which, while slightly lower than the Rb-Sr age ( $1222 \pm 45$  Ma), suffers from the same problems of evaluation as recognized for the Swartmodder Granite.

### 3.4 Piksteel granodiorite suite

The granodiorite pluton included within the Piksteel suite which crops out on the Rehoboth townlands shows a marked discrepancy between the Rb-Sr and zircon U-Pb ages. The higher zircon age of  $1476 \pm 30$  Ma (cf. Rb-Sr of  $1170 \pm 20$  Ma) may be a function of inheritance, in the sense that the composite zircon sample analysed was in fact made up partly of xenocrystic material. Metamorphic resetting of the Rb-Sr system can also be ruled out in this case as the adjacent Swartmodder Granite has not been affected. Another zircon from the type area on Piksteel 209 yields an age significantly younger ( $1076 \pm 25$  Ma), but perhaps more consistent with the Rb-Sr results, although the localities are very far apart. Also this younger zircon sample apparently suffered severe Pb loss, which implies high discordancy and renders the  $^{207}\text{Pb}/^{206}\text{Pb}$  age difficult to interpret. Bearing in mind the geological complexity of the southern Rehoboth townlands, it is not impossible that the zircon sample may not have been derived from a Piksteel age granitoid, and the old U-Pb age may in fact reflect derivation from an unrecognized Swartmodder age granitoid.

### 3.5 Gamsberg Granite Suite

While there are several zircon U-Pb ages available from the Gamsberg Granite Suite, results by both methods exist only for the Gamsberg pluton. In contrast with the other granitoids, the Gamsberg pluton lacks any foliation where it has been sampled, although further to the east of Höhenheim the rocks become intensely gneissic (De Waal, 1966). Analytical results for the Höhenheim zircon sample were published by Hugo and Schalk (1972) and it can be seen that the degree of discordance is not marked, although the degree of Pb loss this represents is model dependent. In other words, if the discordancy is simply the result of geologically recent Pb loss, then the calculated  $^{207}\text{Pb}/^{206}\text{Pb}$  age can be interpreted as recording magmatic crystallization. However, if the Gamsberg zircon had experienced some Pb loss during Damaran metamorphism (at about 500 Ma for example), then the degree of Pb loss is significantly greater (50 % instead of 20 %), and an upper intercept age of  $\sim 1200$  Ma is not impossible. The comparison is further hindered by the poor Rb-Sr isochron statistics exhibited by the Gamsberg pluton. Nevertheless, the results could indicate intrusion of the Gamsberg pluton somewhere between 1 000 and 1 100 Ma.

Although a zircon U-Pb age is available on a Gamsberg suite granitoid cropping out on Nauchas 14, it is most likely derived from a pluton exposed near the Nauchas village, which is distinct from the pluton sampled in this study. It is therefore not impossible that the Nauchas village pluton (as dated by the zircon age of  $1178 \pm 20$  Ma) was intruded before the northern Nauchas pluton (dated by Rb-Sr at  $1073 \pm 25$  Ma). Other explanations for the Nauchas village zircon

age are possible, but their evaluation is not warranted until a strict comparison can be made.

Seifert (1986) also reported Rb-Sr data for the Gamsberg pluton of the Gamsberg Granite Suite. Again poor isochron relationships were obtained and two alternative interpretations were put forward, with suggested ages of  $1229 \pm 29$  Ma and  $1190 \pm 23$  Ma. Combination of the two sets of data for the Gamsberg pluton does not provide any better isochron statistics, although the very high Rb/Sr samples analysed in this study predictably lower the isochron age to a value within error of those obtained for the Namibgrens and Nauchas plutons of this suite. While confinement of the sampling to one single pluton may not always produce good Rb-Sr isochrons, this practice can, by reducing the likelihood of variable initial  $^{87}\text{Sr}/^{86}\text{Sr}$ , reveal the effects of other possible processes which may cause disturbances. In the case of the Gamsberg pluton, the poor Rb-Sr results may have been caused by subsolidus alteration, possibly due to Damaran metamorphism, as suggested by Seifert (1986).

### 3.6 Geochemistry

Rb and Sr concentrations of the various granitoids are plotted in Fig. 15, where separation on to three diagrams allows clearer comparison. The patterns displayed in Fig. 15 aid in both the evaluation of the Rb-Sr isochrons and the problem of lithostratigraphic correlation.

The Rb-Sr results obtained by Seifert (1986) for the Weener Quartz Diorite have raised the possibility that this rock type could be as old as the Naub Quartz Diorite. Such a correlation is possibly supported on compositional grounds, as inspection of Figs 15a and b shows the Naub and Weener of Seifert (1986) having similar Sr contents, although some of the Naub rocks are much lower in Rb. Neither of these two rock types closely resemble typical calc-alkaline diorites and tonalites, as typified in Fig. 15a using the Vioolsdrif batholith as an example (Reid, 1977). The nearest approach is provided by the Weener rocks from the Namibgrens locality, analysed in this study. Relating the rocks from the two Weener localities by some simple differentiation process (thereby demonstrating a possible correlation) is not easily accomplished, as the required trend diverges markedly from that defined by the Vioolsdrif fields.

Despite the unusual Rb-Sr trend displayed by Weener rocks from the two localities, the Biesiepoort Granite has compositions that plot in a very similar field. Furthermore, the Rb-Sr isotopic results show the Biesiepoort and Weener rocks analysed in this study had practically identical initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios, a feature which would support a genetic relationship. It follows that if the trend displayed by the Weener and Biesiepoort granitoids analysed in this study was produced by some differentiation process, then a relationship between the two Weener localities can not be ruled out simply on compositional grounds. What is indicated is that processes other than that which produced the Vioolsdrif trend may have occurred in the Rehoboth-Nauchas region.

Apart from one exception, the Swartmodder Granite shares the relatively low Sr contents seen in the Naub Diorite, and this distinguishes the former from the Biesiepoort Granite. The Swartmodder Granite can further be distinguished from the less differentiated varieties of the Gamsberg Granite Suite by its lower Rb contents. The Rb-rich nature of the Piksteel rocks also serves to separate this younger granitoid suite from the Swartmodder. Bearing in mind the likelihood of mistaking Swartmodder-age granitoids (with similar petrography) for Piksteel and Gamsberg suite units, the Rb-Sr patterns displayed in Fig. 15 suggest that a regional geochemical study may aid lithostratigraphic mapping.




Fig. 15: Covariation of Rb and Sr for the granitoids analysed in this study. 15a: Granitoids with the oldest Rb-Sr ages, labelled fields enclose data for the Vioolsdrif granitoids (Reid, 1977); 15b: Younger granitoids, including data for the Weener Quartz Diorite from Seifert (1986); 15c: Gamsberg Granite Suite, including data for the Gamsberg pluton from Seifert (1986).

The Gamsberg granitoids are distinguished in Fig. 15 by their relatively low levels of Sr and enrichment in Rb. The tight field with Sr contents between 100–200 ppm constitutes the typical coarse-grained Gamsberg Granite that comprises most of the three plutons studied. The samples showing strong depletion in Sr are local variants that are very leucocratic and sometimes pegmatitic, and appear to be examples of extreme differentiation of the Gamsberg parent magma. The trend defined by the Gamsberg rocks is very similar to the corresponding part of the Vioolsdrif fields, where crystallization of adamellite magmas produced leucogranites extremely depleted in Sr and moderately enriched in Rb. It follows that the Gamsberg trend may represent simple differentiation and therefore not argue against incorporation of the high Rb/Sr samples in the Rb-Sr isochrons.

If all the granitoids with Rb-Sr ages in the 1200–1000 Ma range are inspected in Figs 15b and c, the impression is one of a fairly continuous pattern of increasing Rb/Sr ratio with decreasing emplacement age. In other words, the earliest granitoids (Weener Quartz Diorite) are the least differentiated, with the Biesiepoort Granite and Piksteel Granodiorite forming intermediate members, while the Gamsberg Granite Suite represents the most evolved magmas emplaced at the end of what is a major magmatic

episode. On the other hand, the older granitoids (Naub and Swartmodder) define a distinctly different trend in Fig. 15, thereby suggesting a separate system that could be regarded as having lithostratigraphic significance.

### 3.7 Proterozoic crustal evolution

Initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios for the various granitoids analysed in this study are plotted against Rb-Sr isochron age in Fig. 16. The younger granitoids clearly have a wide variation in initial ratios, with the Biesiepoort and Weener granitoids being derived from a much less radiogenic source than the later Piksteel and Gamsberg magmas. This difference rules out processes like simple crystal fractionation in the production of the compositional range of the granitoids, as represented by Rb and Sr contents discussed in the previous section. Rather the change in average Rb/Sr ratio with time appears to be related to initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio, so that the trends are controlled in some way by source region characteristics. In other words, the Biesiepoort/Weener granitoids represent products from the lower crust and upper mantle, while the slightly younger Piksteel and Gamsberg granitoids represent products from the upper crust. The trend therefore involved increasing contributions from older continental crust with time, a feature which has been already established for the nearby Damara Orogen (Miller, 1983) and the Namaqua Province (Reid and Barton, 1983), but is the reverse to that established for the Hercynian Orogen in Europe (Hamet and Allègre, 1975).




Fig. 16: Plot of initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios against Rb-Sr isochron age for granitoids analysed in this study. Error bars are two sigma estimates from the regression analyses. Arrows define the average rate of increase in  $^{87}\text{Sr}/^{86}\text{Sr}$  with time for each rock type, and the figures in boxes represent the average Rb/Sr ratio calculated from the data in Table 2. Broken line represents the variation in  $^{87}\text{Sr}/^{86}\text{Sr}$  with time for a model mantle having 'bulk earth' composition (O'Nions *et al.*, 1979).

The older granitoids have emplacement ages similar to those already established for other mid-early Proterozoic terranes in South West Africa. Initial Sr isotopic ratios also vary, with the Swartmodder Granite having a relatively high value, suggesting a crustal source region already very radiogenic 1600–1700 Ma ago. The very high average Rb/Sr ratio of the Swartmodder Granite may reflect derivation from a high Rb/Sr source, so that projection back on Fig. 16 to the mantle curve may not yield an age significantly in excess of that for the Naub Diorite. It follows that the maximum age of formation of the crustal protoliths may not exceed ~2000 Ma, a result also reported by Hawkesworth and Marlow (1983) for the Damaran granitoids, although it would



be useful to investigate the Sm-Nd systematics of these old Rehoboth rocks before proceeding further in the assessment.

#### 4. ACKNOWLEDGEMENTS

S.T.M. acknowledges financial support from Rio Tinto Pty. Ltd. for the period 1975–78, during which the field and laboratory work were carried out. D.L.R. acknowledges the continued persistence of R.McG. Miller, P. Joubert and H.L.A., who collectively urged for this rather belated collation and interpretation of the Rb-Sr results, which were discovered in a filing cabinet long after the project had been completed. K. Hoffmann and K. Schalk kindly made unpublished geological data available and B. Eglinton provided a helpful review.

#### 5. REFERENCES

- Allsopp, H.L., Welke, H.J., Burger, A.J., Kröner, A. and Blignault, H.J. 1979. Rb-Sr and U-Pb geochronology of late Precambrian–early Palaeozoic igneous activity in the Richtersveld (South Africa and southern South West Africa). *Trans. geol. Soc. S. Afr.*, **82**, 195–205.
- Borg, G. and Maiden, K.J. 1986. A preliminary appraisal of the tectonic and sedimentological environment of the Sinclair Sequence in the Klein Aub area. *Communs geol. Surv. S.W. Africa/Namibia*, **2**, 65–73.
- Brooks, C., Hart, S.R. and Wendt, I. 1972. Realistic use of two error regression treatment as applied to Rb-Sr data. *Rev. Geophys. Sp. Phys.*, **10**, 551–577.
- Burger, A.J. and Coertze, F.J. 1973. Radiometric age measurements on rocks from Southern Africa to the end of 1971. *Bull. geol. Surv. S. Afr.*, **58**, 46 pp.
- Burger, A.J. and Coertze, F.J. 1975. Age determinations — April 1972 to March 1974. *Annals geol. Surv. S. Afr.*, **10**, 135–142.
- Burger, A.J. and Coertze, F.J. 1977. Summary of age determinations carried out during the period April 1974 to March 1975. *Annals geol. Surv. S. Afr.*, **11**, 317–322.
- Burger, A.J. and Walraven, F. 1977. Summary of age determinations carried out during the period April 1975 to March 1976. *Annals geol. Surv. S. Afr.*, **11**, 323–329.
- Burger, A.J. and Walraven, F. 1979. Summary of age determinations carried out during the period April 1976 to March 1977. *Annals geol. Surv. S. Afr.*, **12**, 199–208.
- Burger, A.J., Clifford, T.N. and Miller, R.McG. 1976. Zircon U-Pb ages of the Franzfontein granitic suite, northern South West Africa. *Precamb. Res.*, **3**, 415–431.
- De Waal, S.A. 1966. *The Alberta Complex, a metamorphosed layered intrusion north of Nauchas, South West Africa, the surrounding granites and repeated folding in the younger Damara system*. D.Sc. Thesis (unpubl.), Univ. Pretoria.
- Duncan, A.J., Erlank, A.J. and Betton, P.J. 1984. Appendix 1: Analytical techniques and database descriptions, 389–395. In: Erlank, A.J. (Ed.), *Petrogenesis of the Volcanic Rocks of the Karoo Province*. Geol. Soc. S. Afr. Spec. Publ., **13**, 395 pp.
- Hamet, J. and Allègre, C.J. 1976. Hercynian orogeny in the Montange-Noire (France): Application of Rb-Sr systematics. *Bull. geol. Soc. Am.*, **87**, 1429–1442.
- Hawkesworth, C.J. and Marlow, A.G. 1983. Isotope evolution of the Damara orogenic belt, 397–407. In: Miller, R.McG. (Ed.), *Evolution of the Damara Orogen of South West Africa/Namibia*. Geol. Soc. S. Afr. Spec. Publ., **11**, 515 pp.
- Hugo, P.J. and Schalk, K.E.L. 1972. The isotopic ages of certain granites and acid lavas in the Rehoboth and Maltahöhe districts, South West Africa. *Annals geol. Surv. S. Afr.*, **9**, 103–105.
- Jacob, R.E., Kröner, A. and Burger, A.J. 1978. Areal extent and first U-Pb age of the pre-Damara Abbabis Complex in the central Damara belt of South West Africa/Namibia. *Geol. Rdsch.*, **67**, 706–718.
- Miller, R.McG. 1983. The Pan-African Damara orogen of South West Africa/Namibia, 431–515. In: Miller, R.McG. (Ed.), *Evolution of the Damara Orogen of South West Africa/Namibia*. Geol. Soc. S. Afr. Spec. Publ., **11**, 515 pp.
- O'Nions, R.K., Carter, S.R., Evensen, N.M. and Hamilton, P.J. 1979. Geochemical and cosmochemical applications of Nd-isotope analysis. *Ann. Rev. Earth Planet. Sc.*, **7**, 11–38.
- Reid, D.L. 1977. Geochemistry of Precambrian igneous rocks in the lower Orange River region, Southern Africa. *Bull. Precamb. Res Unit, Univ. Cape Town*, **22**, 397 pp.
- Reid, D.L. 1979a. Total rock Rb-Sr and U-Th-Pb isotopic study of Precambrian metavolcanic rocks in the lower Orange River region, Southern Africa. *Earth. Planet. Sc. Lett.*, **42**, 368–378.
- Reid, D.L. 1979b. Age relationships within the Mid-Proterozoic Vioolsdrif batholith, lower Orange River region. *Trans. geol. Soc. S. Afr.*, **82**, 305–311.
- Reid, D.L. and Barton, E.S. 1983. Geochemical characterisation of granitoids in the Namaqualand Geotraverse, 67–82. In: Botha, B.J.V. (Ed.), *Namaqualand Metamorphic Complex*. Geol. Soc. S. Afr. Spec. Publ., **10**, 198 pp.
- South African Committee for Stratigraphy (SACS) 1980. *Stratigraphy of South Africa*. Part 1 (Comp. Kent, L.E.). *Lithostratigraphy of the Republic of South Africa, SWA/Namibia, and the Republics of Bophuthatswana, Transkei and Venda*. Handb. geol. Surv. S. Afr., **8**, 690 pp.
- Schalk, K.E.L. *Geological map 2616 — Rehoboth; Scale 1:250 000*. Geol. Surv. S.W. Afr./Namibia. (unpubl.).
- Seifert, N. 1986. Geochronologie am Sudrand des Damara-Orogens, SWA/Namibia. *Schweiz. miner. petrogr. Mitt.*, **66**, 413–451.
- Steiger, R.H. and Jäger, E. 1977. Subcommittee on geochronology: Convention on the use of decay constants in geo- and cosmochronology. *Earth Planet. Sc. Lett.*, **36**, 359–362.
- York, D. 1969. Least squares fitting of a straight line with correlated errors. *Earth Planet. Sc. Lett.*, **5**, 320–324.