

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/366929741>

Response of woody vegetation to bush thinning on freehold farmlands in north-central Namibia

Article in *Scientific Reports* · January 2023

CITATIONS

0

READS

158

1 author:



Mark Appiah

CSIR College of Science and Technology, Ghana

48 PUBLICATIONS 922 CITATIONS

SEE PROFILE



OPEN

Response of woody vegetation to bush thinning on freehold farmlands in north-central Namibia

Matti T. Nghikembua^{1,2}, Laurie L. Marker¹, Bruce Brewer¹, Arvo Leinonen³, Lauri Mehtätalo⁴, Mark Appiah^{2,5,6} & Ari Pappinen²✉

Bush encroachment affects much of the Namibian woodland landscape, causing significant loss of open savannah habitat and farm profits. Thinning of the trees/shrubs is recommended; however, research is required to identify the overall efficacy and effects of this method on the woodland habitat. We aimed to examine the effect of the thinning strategy applied on the vegetation structure of encroaching tree/shrub species, as well as the sighting lines of the habitat. Vegetation surveys were done on three freehold farms in north-central Namibia. The study utilised a combination of a blocked and split-plot study design: each block consisted of a pair of thinned and non-thinned plots with multiple subplots. Thinned plots had been manually thinned, with a post-thinning age of three years or more. Results revealed that tree/shrub abundance differed between species; thinned areas had the least abundance and overall species-treatment interactions were significant. Thinning caused a significant reduction in overall tree/shrub densities, settling within the recommended range for the area. Thinning also significantly reduced the average tree/shrub height, canopy area, medium-sized trees/shrubs, and increased sighting lines. This confirms a bush encroachment mitigation strategy that favours grass cover, and wildlife that rely on longer sighting lines for safety or when hunting.

Namibian farmland has undergone significant ecological changes as a result of anthropogenic influences, resulting in an increase in the density and biomass of native woody species, a phenomenon known as bush encroachment^{1–3}. Approximately 45 million hectares (ha) of the territory has been affected, causing significant declines in the grazing carrying capacity, beef production, suitable habitats for some grazing ungulates, and predator numbers (e.g., cheetah *Acinonyx jubatus*)^{1,3–5}. This phenomenon also has potential to show varied impacts to the local climates according to geographic region. For example, in the humid northern temperate regions (southwest of North America and center of Central Asia), shrub encroachment in grasslands lowered daytime land surface temperatures and albedo, but increased evapotranspiration that is typically associated with heat loss^{6,7}. In contrast, higher land surface temperatures were experienced at nighttime, due to the vegetation shading effects. In the arid and semi-arid northern temperate regions, shrub encroachment led to the increase in daytime and nighttime land surface temperatures, reduction in leaf area index and evapotranspiration, however, albedo was increased⁷. These effects were attributed to higher fraction of bare soil patches in encroached areas that experiences heat fluxes during the day and greater outgoing longwave radiation at night. Also, the reduced evapotranspiration experienced accounted for less heat loss and cooling effect, consequentially causing warming. Shrub encroachment has also been observed to correspond positively to the increase in land surface temperatures and extended growing seasons in China's temperate grasslands region⁸. Effects of this phenomenon towards the local land surface temperatures are unknown on Namibian rangelands. However, bush encroachment may have some warming and cooling effect due to higher evapotranspiration, vegetation shading effects and presence of bare soil patches that could contribute to the absorption and reflection of heat energy.

To counteract the negative effects of this phenomenon, restoration thinning selectively removes the over-abundant trees/shrubs and is recommended since it minimises negative impacts on the vegetation, habitats or biodiversity². Harvesting of trees/shrubs has potential to alter the structure or dynamics of their population as

¹Biomass Technology Centre, Cheetah Conservation Fund, Elandsvregde 367, Otjiwarongo 12001, Namibia. ²School of Forest Sciences, University of Eastern Finland, Joensuu Campus, Yliopistokatu 7, Joensuu, 80101 North Karelia, Finland. ³Bioenergy Consult Arvo Leinonen, Huikkolantie 21, Vaajakoski, 40800 Jyväskylä, Finland. ⁴Natural Resources Institute Finland (LUKE), Yliopistokatu 6, 80101 Joensuu, Finland. ⁵CSIR-Forestry Research Institute of Ghana (CSIR-FORIG), P. O. Box UP 63 KNUST, Kumasi, Ghana. ⁶Faculty of Natural Science and Environmental Management, Department of Natural Resources Management, CSIR College of Science and Technology (CCST), P.O. Box M32, Accra, Ghana. ✉email: ari.pappinen@uef.fi

woody densities and biomass are reduced, and compositions may change due to some species with certain growth characteristics being preferred or avoided. Also, the reduction of densities releases competition for resources (e.g., water, nutrients, sunlight), causing the retained individuals to increase their growth rates, reproduction, and enhancing sapling survival^{9,10}. Trees/shrubs also play a key role in carbon sequestration and the nutrient cycle in savannah ecosystems: the sub-habitats underneath canopies or near dead trees contain substantial concentrations of soil elements, such as nitrogen (N), organic matter (OM), calcium (Ca), phosphorus (P), potassium (K), and carbon (C)^{11–15}. Also, trees/shrubs increase the soil microbial biomass that is important to nutrient cycling, including the mineralization of plant, animal and organic carbon, and leguminous tree/shrub species are also N fixers^{11,16–18}. Severe or uncontrolled bush harvesting may result in the disruption of all these crucial roles played by the vegetation in the savannah environment. However, if done in moderation, it may not disrupt the stability of the ecosystem^{19,20}. Protective measures with regard to biodiversity and the vegetation are crucial as they provide ecological services, which include nutrient cycling, erosion control, water cycling, carbon sequestration, forage for animals, breeding sites, cover against predators or adverse weather conditions, construction materials, and a source of energy^{2,12–15,17,21}.

The farmland matrix serves as an important habitat for livestock farming and biodiversity conservation. As an example, the majority (> 80%) of cheetah and numerous protected and economically important large ungulates, including eland (*Tragelaphus oryx*), oryx (*Oryx gazella*), greater kudu (*Tragelaphus strepsiceros*) and red hartebeest (*Alcelaphus buselaphus* ssp. *caama*), reside on freehold farmlands^{22–24}. The restoration of bush-encroached habitat to open shrubland would increase grazing carrying capacity, and consequentially, would lead to greater farmland profits and boost prey populations, thereby reducing the likelihood for human-wildlife conflict^{1,25}. Hence, for this reason, inventory and monitoring of restoration thinning habitats are increasingly important aspects for sustainable livestock management and biodiversity conservation, and should lead to appropriate management actions.

Absolute population estimates (expressed per unit area) are commonly applied to describe the abundance of species in a specific habitat, evaluate management impacts on species and habitat suitability for wildlife and associations of plant species^{21,26–30}. Density (expressed as the number of individuals per unit area) is one such estimate. Attributes, such as height, canopy diameter or stem diameter often accompany density estimates to quantify biomass volumes and vegetation structure^{9,31–33}. Biomass (wet, dry weight, organic carbon content) contained in trees/shrubs is useful in understanding the trophic structure, carbon sequestration potential, availability of biomass for offtake, or dominance of species in a particular area^{31,34–37}. Such estimates are obtained either directly through destructive sampling, or indirectly using allometric equations that relate oven dry weight with tree/shrub attributes (e.g. height, stem diameter, canopy diameter or spatial canopy volume)^{31,38}.

Indirect acquisition of estimates is efficient due to lower required labour intensity and the time required to obtain samples directly. However, there is a lack of verified models for some species, since current models cannot be applied universally due to differences between species or variations in environmental conditions that could cause over/under estimation of the biomass^{35,36,39}. However, a number of allometric equations, especially suitable for the small, multi-stemmed bush-encroaching species found in semi-arid landscapes in southern Africa, have been published^{31,40}. Hence, use of these allometric models may be appropriate to estimate available bush biomass in our study area since they were developed for almost similar species and region^{39,41}.

The need to restore rangeland productivity and wildlife habitats by thinning the overabundant biomass was the rationale for Cheetah Conservation Fund (CCF) to establish the CCF Bush Project in 2001^{42,43}. To date, a number of studies have reported the impacts of this project on soil fertility, vegetation regeneration, and the response of wildlife^{17,28,44,45}. Our study differs from previous research in that it examines (a) the effects of treatment (thinned vs non-thinned) on the abundance (density, biomass) and structure (height, canopy area, stem size) of bush-encroaching species, and (b) evaluates habitat sighting lines, to gain insight into how effective this method has been in reducing bush encroachment on the north-central freehold farmlands of Namibia.

We focused the research on three questions: (1) how do thinned and non-thinned areas differ in vegetation structure (i.e., height, canopy area, stem size distribution) and abundance (density, woody biomass quantity), (2) in what ways has selective thinning influenced tree/shrub density (expressed as tree equivalents (TE) ha⁻¹) recommended for the area, and (3) in what ways has selective thinning affected sighting lines, a key habitat feature associated with cheetah hunting success. We hypothesized that thinning would reduce the abundance of encroaching species, that the magnitude of the impacts at thinned sites would differ between tree/shrub types, and that thinning would significantly modify the vegetation structure and restore sighting lines for improved detection. This study provides baseline data with regard to the vegetation structure in both encroached and previously thinned areas, which will allow evaluation of the efficacy of restoration thinning and assist with the recommendation of appropriate management actions for the affected areas.

Material and methods

Study area. The study area consisted of three freehold farms; Cheetah View (#317), Boskop (#324) and Elandsvreugde (#367), in north-central Namibia (farm size: 5046–7300 ha); coordinates: 20.42407° S–20.58440° S, 16.86337° E–17.09950° E (Fig. 1). Elevation is approximately 1580 m; the topography of the farms is generally flat and there are no perennial rivers. The farms were developed for livestock and are surrounded by cattle proof fences (~ 1.5 m high) that allow wildlife movement. Semi-permanent waterpoints provide year-round access to water. The region experiences a semi-arid climate with three main seasons: hot-wet (January–April), cold-dry (May–August) and hot-dry (September–December). Mean annual rainfall is 401 mm (± 257.4), which is concentrated mainly between January–April. Mean annual temperature is 19.2 °C (± 2.4), and mean daily maxima is 22.7 °C (± 0.7) in January and 13.4 °C (± 0.7) in July⁴⁶.

The predominant soil types are Eutric Regosols and Chromic Cambisols^{44,47}. Soil texture types consist of sandy loam and loamy sand. The vegetation is broadly classified as thornbush shrubland, dominated by *Dichrostachys*




Figure 1. Location of the study area within the north-central farmland of Namibia. Grey squares indicate non-thinned and open squares indicate thinned plots. Block pairs are numbered 1–13. Map created with QGIS Geographic Information System software, version 3.18.3-Zürich, <https://www.qgis.org>.

cinerea (L.) Wight & Arn, *Senegalia mellifera* (Vahl) Seigler & Ebinger, *Vachellia reficiens* (Wawra & Peyr.) Kyal. & Boatwr., *Terminalia sericea* Burch. ex DC., *Terminalia prunioides* M.A.Lawson, *Combretum imberbe* Wawra, *Grewia flava* DC. and *Grewia flavescens* Juss. species^{47,48}.

Species of interest. Observations focused on five thornbush woody species known to cause encroachment in the area: sickle bush (*D. cinerea* subspecies spp. *africana*), black-thorn acacia (*S. mellifera* subsp. *detinens*), blade thorn (*Senegalia cinerea* (Schinz) Kyal. & Boatwr., red umbrella thorn (*V. reficiens*), and umbrella thorn (*Vachellia tortilis* (Burch.) Kyal. & Boatwr. spp. *heteracantha*). These native trees/shrubs are widespread in the region. Distinguishing features of these species include sharp thorns and spines (straight and hooked), as well as fine compound and bipinnate leaves. These species have economic value since local communities rely on their raw material for fuelwood, charcoal production, as handles for axes/picks, fencing poles, construction materials, as forage for livestock and game, and as an ingredient in the production of animal feeds and traditional medicine^{49–51}.

All five encroaching species were manually thinned using handheld tools (axes and machetes) to reduce their densities by approximately 50%. Thinning of trees/shrubs was random and included all height classes to retain a heterogeneous mix of tree/shrub heights. Smaller tree/shrubs (≤ 4 m. height, ≤ 18 cm stem diameter) were the most selectively thinned (40–60%) due to their higher abundance in the study area. Larger trees/shrubs with ≥ 18 cm stem diameters were avoided. Some vegetation clumps were left undisturbed to provide cover, browse, shelter or breeding sites for wildlife.

Study design and data collection. A combination of a blocked and split-plot study design was used in this study. This included 13 blocks (also called plot pairs), each with two plots with a mean (\pm standard deviation) size of 12.62 (± 8.43) ha, one for each treatment (Fig. 1). For the collection of vegetation data, 4–20 circular subplots with a 6 m radius (area = 113.1 m²) were placed within each plot. A total of 295 subplots were surveyed: 49.5% ($n = 149$) were in the thinned and 50.5% ($n = 146$) in the non-thinned treatment area. The treated plots were established on areas that were manually thinned from 2002 to 2013 with a post-thinning age equal to 3 years or more in 2016, so that multiple seasons followed thinning. A single thinning cycle has been conducted, and tree/shrub densities were reduced by approximately 50%. Since 2005, aftercare treatment to prevent stump regrowth was carried out using Picloram and Tryclopypyr active arboricide (Access®) on freshly cut stumps. Eleven previously thinned plots had been treated with Access® and two plots that had been thinned earlier were not treated. The control treatment plots were established on areas where thinning had not been conducted. These were areas near the thinning treatment, located at a mean distance of approximately 0.71 (± 0.17) km in the same habitat that was affected by bush encroachment, with similar tree/shrub species and management history^{28,45}.

A systematic random plot sampling method was used to designate the vegetation subplots, which were located 100 m apart in a plot. Vegetation data was collected from April to August 2017. We quantified sighting lines at the subplot level by positioning an observer who crouched 0.65 m aboveground, and measured the initial distance at which objects (human) could not be detected by an observer^{4,52}. The starting bearing (direction) was random, three other similar measurements were added sequentially at 90° from the previous bearing. We calibrated the measurements by allowing a second person (preferably wearing non-bright colors) to walk away from the observer until they could not be seen. A one-meter accuracy rangefinder (Bushnell Yardage Pro Scout 6×) was used to take the measurements.

In addition, all encroaching trees/shrubs present in each subplot were assessed as follows: (1) counts of individuals for all target species, (2) tree/shrub height using an extendable marked polyvinyl chloride (PVC) pipe (3 m length), a standard ruler (30 cm) and a measuring tape (30 m) (for tall trees/shrubs (i.e., > 3 m high), the PVC pipe was used to measure heights), (3) tree/shrub condition based on two appearance categories: alive or dead, (4) maximum canopy diameter taken with a 30 m measuring tape from two perpendicular directions (d1, d2), and (5) base (stem) diameter, recorded at 10–15 cm aboveground.

Aboveground biomass quantification. Aboveground biomass of the encroaching species (live individuals) was estimated with the allometric regression equation developed for Namibian thornbush species⁴⁰ that relate biomass with tree/shrub height and stem diameter or the combination of both as predictors: $y = ax^b$, where y = dry woody biomass; a = intercept, x = independent variable/s (tree/shrub height, stem diameter) and b = slope. For each tree/shrub, a combined biomass value was calculated for wood sizes ≤ 2 cm (ideal for bush feed), as well as > 2 cm (wood) (Appendix A1).

Data analysis. A generalized linear mixed-effects model (GLMM) via penalized quasi-likelihood (glmmPQL) (package MASS, Venables and Ripley, 2002) was used to estimate how the number of trees/shrubs per species and stem diameter class were influenced by treatment:

$$E(y_{kji}) = \exp(\mathbf{x}'_{kji}\boldsymbol{\beta} + b_k + c_{kj} + d_{kji}),$$

where $E(y_{kji})$ = mean number of tree/shrub counts; $\mathbf{x}'_{kji}\boldsymbol{\beta}$ includes the effects of fixed factors; tree/shrub species or stem classes, treatment, the tree/shrub species-treatment or stem class-treatment interactions, arboricide application and post-thinning age (expressed as time since thinning minus 7.2 years), where 7.2 is the mean time since thinning over all treated plots; and $b_k + c_{kj} + d_{kji}$ includes the nested random effects for block (plot pair), plot and subplot, respectively. For the non-thinned treatment, the post-thinning age was set to zero. We included aftercare application with arboricide in the estimation, as 11 previously thinned plots were treated and two had not been treated. Hypothesis testing for the GLMM model coefficients were based on Wald χ^2 tests of the fitted final model using R functions Anova and lht of package car (Fox and Weisberg, 2011).

Linear mixed-effects (LME) models (package nlme, Bates et al., 2021) were used to estimate the influence of treatment on the overall tree/shrub density (TE), canopy size, woody biomass availability and sightlines in four separate models of the form:

$$y_{kji} = \mathbf{x}'_{kji}\boldsymbol{\beta} + b_k + c_{kj} + e_{kji},$$

where y_{kji} = mean number of trees/shrubs (TE), canopy size, woody biomass, and sighting lines; $\mathbf{x}'_{kji}\boldsymbol{\beta}$ includes the effects of treatment, post-thinning age and arboricide aftercare as fixed factors; and $b_k + c_{kj}$ includes the nested random effects for block (plot pair) and plot, with e_{kji} the subplot-level residual error, respectively. Further, the LME model was also used to estimate how the vertical structure (height) of the areas was affected by treatment:

$$y_{kjil} = \mathbf{x}'_{kjil}\boldsymbol{\beta} + b_k + c_{kj} + d_{kji} + e_{kjil},$$

where y_{kjil} = mean tree/shrub height of the subplot; $\mathbf{x}'_{kjil}\boldsymbol{\beta}$ includes the effects of fixed factors; treatment, post-thinning age and aftercare application; and $b_k + c_{kj} + d_{kji}$ includes the nested random effects for block (plot pair), plot and subplot, with e_{kjil} the tree-level residual error, respectively. The fit of all final models was evaluated graphically by exploring the trends and homoscedasticity of the Pearson and deviance residuals; normal q–q plots of random effects were used to evaluate the assumption of the normality of random effects⁵⁵. For all LME models, the non-constant residual variance was modelled by applying weighting (variance Power) following the inspections of the residuals in the initially fitted models:

$\text{var}(\epsilon_{kji}) = \sigma^2 |y_{kji}|^{2\delta}$, for tree/shrub densities (TE), canopy area, woody biomass, and sightlines and $\text{var}(\epsilon_{kjil}) = \sigma^2 |y_{kjil}|^{2\delta}$ for vertical structure (height), where y_{kji} and y_{kjil} indicates group-level predictions for the models. Conditional F-tests on the fixed effects were carried out to determine significant predictors. All analyses were carried out with R version 4.0.4⁵⁶ at the $P \leq 0.05\%$ significance level.

Approval of bush thinning and field research. Permission to conduct field research on the farms was acquired from the landowner (Cheetah Conservation Fund); all harvest operations, ecological research and monitoring program were done in accordance with the laws of the Namibian Ministry of Environment Tourism and Forestry (METF) (research permits: 2151/2016, 2324/2017) and certified by the Forest Stewardship Council (FSC) for sustainable forestry (certificate: FSC-C004580).

Results

Number of tree/shrub observations and structural characteristics. A total of 4223 trees/shrubs from five species were observed of which most (92.5%) were alive and few (7.5%) were dead (Table 1). Live trees/shrubs were recorded in almost all subplots (97.97%), although none were recorded in six subplots (thinned = 3, non-thinned = 3). Dead trees/shrubs were recorded in 56.9% of subplots, with most of the observations in the non-thinned area (Table 1).

Tree/shrub height was measured amongst all 3908 live trees/shrubs (Table 1). Overall, red umbrella thorn and blade thorn were, on average, the tallest (> 1.5 m) species. Canopy diameter was measured for 81.2% (n = 3174) of the live trees/shrubs (Table 1). On average, canopy size per individual tree/shrub was greatest in red umbrella thorn, followed by black-thorn acacia, blade thorn and umbrella thorn (Table 1). Sickie bush had the smallest average canopy size of all species. Stem diameter was measured for 90.3% (n = 3529) of the live trees/shrubs (Table 1). On average, red umbrella thorn had the largest stem diameter (> 7 cm) per individual tree/shrub. Other species with large stem diameters (4–5 cm) were black-thorn acacia, blade thorn and umbrella thorn. Sickie bush had the smallest average stem diameter of all species (Table 1).

Number of observed individuals and species-treatment interactions. The baseline category (β_0) of the GLMM model (Tables 2 and 3, Fig. 2) showed that the mean number of live sickie bush specimens counted

Variable/common name	Thinned			Non-thinned			Total Abundance
	Abundance	Mean	SD	Abundance	Mean	SD	
Overall (alive and dead)							
Sickle bush	887	–	–	1034	–	–	1921
Black-thorn acacia	640	–	–	290	–	–	930
Blade thorn	318	–	–	340	–	–	658
Red umbrella thorn	115	–	–	388	–	–	503
Umbrella thorn	190	–	–	21	–	–	211
Total	2150	–	–	2073	–	–	4223
Height (m)/ all live individuals							
Sickle bush	848	0.94	0.74	841	1.52	0.91	1689
Black-thorn acacia	619	0.55	0.78	269	2.44	1.67	888
Blade thorn	310	1.35	0.99	329	1.96	1.16	639
Red umbrella thorn	111	1.48	1.49	370	2.66	1.49	481
Umbrella thorn	190	1.27	0.95	21	2.26	1.77	211
Total	2078	0.94	0.92	1830	1.97	1.32	3908
Canopy diameter (m)							
Sickle bush	797	0.93	0.81	667	1.53	1.08	1464
Black-thorn acacia	411	0.75	1.1	201	2.79	2.23	612
Blade thorn	284	1.51	1.27	268	2.1	1.3	552
Red umbrella thorn	101	1.94	1.97	247	2.84	2.2	348
Umbrella thorn	182	1.73	1.17	16	2.31	2.13	198
Total	1775	1.12	1.16	1399	2.06	1.68	3174
Stem diameter (cm)							
Sickle bush	755	2.44	2.25	748	2.72	3.23	1503
Black-thorn acacia	515	2.55	4.53	243	8.24	7.16	758
Blade thorn	291	3.89	3.9	308	4.6	4.7	599
Red umbrella thorn	107	5.54	7.19	362	7.98	6.9	469
Umbrella thorn	182	4.23	4.79	18	9.74	10.98	200
Total	1850	3.06	4.02	1679	5.07	5.8	3529
Dead trees/shrubs							
Sickle bush	39	–	–	193	–	–	232
Black-thorn acacia	21	–	–	21	–	–	42
Blade thorn	8	–	–	11	–	–	19
Red umbrella thorn	4	–	–	18	–	–	22
Umbrella thorn	–	–	–	–	–	–	–
Total	72	–	–	243	–	–	315

Table 1. Observations of trees/shrubs by species and measured structural characteristics in the thinned and non-thinned treatments. The number of trees/shrubs counted are indicated as abundances; the height, canopy and stem diameters are expressed as averages (mean) with associated standard deviation (SD) values for each species in the non-thinned and thinned areas.

Variable	Parameter (β)	Value	SE	Exp(β)
Fixed effects				
(Intercept)	β_0	1.777	0.103	5.912
Black-thorn acacia	β_1	- 1.142	0.166	0.319
Blade thorn	β_2	- 0.941	0.154	0.39
Red umbrella thorn	β_3	- 0.823	0.148	0.439
Umbrella thorn	β_4	- 3.692	0.523	0.025
Treatment thinned	β_5	- 0.582	0.297	0.559
Post-thinning age	β_6	0.107	0.036	1.113
Arbicide aftercare applied	β_7	0.571	0.307	1.77
Black-thorn acacia: thinned	β_8	0.826	0.208	2.285
Blade thorn: thinned	β_9	- 0.067	0.220	0.936
Red umbrella thorn: thinned	β_{10}	- 1.211	0.281	0.298
Umbrella thorn: thinned	β_{11}	2.195	0.557	8.984
Random effects				
		SD		
Subplot	d_{kji}	0.0001		
Plot	c_{kj}	0.131		
Block	b_k	0.182		

Table 2. Generalized linear mixed-effects model (GLMM) showing the effects of fixed factors (treatment, post-thinning age, and arbicide aftercare) on the abundance of encroaching trees/shrubs.

Treatment/species	Model coefficients (β)	Sum (β)	Mean counts exp(sum(β))	\pm 95% confidence level
Non-thinned				
Sickle bush	$\beta_0 + \beta_5(0)$	1.777	5.912	1.224
Black-thorn acacia	$\beta_0 + \beta_1(1) + \beta_5(0) + \beta_8(0)$	0.635	1.887	1.692
Blade thorn	$\beta_0 + \beta_2(1) + \beta_5(0) + \beta_9(0)$	0.836	2.307	1.653
Red umbrella thorn	$\beta_0 + \beta_3(1) + \beta_5(0) + \beta_{10}(0)$	0.954	2.595	1.633
Umbrella thorn	$\beta_0 + \beta_4(1) + \beta_5(0) + \beta_{11}(0)$	- 1.915	0.147	3.401
Thinned				
Sickle bush	$\beta_0 + \beta_5(1)$	1.195	3.305	2.367
Black-thorn acacia	$\beta_0 + \beta_1(1) + \beta_5(1) + \beta_8(1)$	0.879	2.41	4.910
Blade thorn	$\beta_0 + \beta_2(1) + \beta_5(1) + \beta_9(1)$	0.188	1.207	4.914
Red umbrella thorn	$\beta_0 + \beta_3(1) + \beta_5(1) + \beta_{10}(1)$	- 0.839	0.432	5.468
Umbrella thorn	$\beta_0 + \beta_4(1) + \beta_5(1) + \beta_{11}(1)$	- 0.302	0.74	19.511

Table 3. Expected number of tree/shrubs counted per subplot (area = 113.1 m²) in the thinned and non-thinned treatment areas estimated from the generalized linear mixed-effects model (GLMM) model coefficients.




Figure 2. Expected number of tree/shrub densities (plants ha⁻¹) in the thinned and non-thinned treatment areas estimated from the generalized linear mixed-effects model (GLMM) coefficients. Error bars indicate \pm 95% confidence level of the variability in tree/shrub densities.

in the non-thinned area was 5.91 individuals per subplot (equates to 522.74 plants ha⁻¹). In the same treatment, all other encroacher species were less abundant than the baseline category (β_0); mean counts of black-thorn acacia, blade thorn, red umbrella thorn, and umbrella thorn were 68.1%, 61%, 56.1%, and 97.5% different, respectively, from the live sickle bush count. The overall number of individuals counted in the non-thinned area was 12.85 (1136 plants ha⁻¹).

The parameter estimate β_5 refers to the mean count of sickle bush specimens in the thinned treatment area (Tables 2 and 3). The number of sickle bush specimens was 55.9% of the expected amount in the non-thinned area (non-thinned = 5.91, thinned = 3.31). For the other species, less individuals were also recorded in the thinned area, with a difference of 47.7% for blade thorn and 83.4% for red umbrella thorn compared to the non-thinned area. In contrast, the mean counts in the thinned area were greater by 21.7% for black-thorn acacia and by 80.1% for umbrella thorn than amounts compared to the non-thinned area. The overall number of individuals counted in the thinned area was 8.09 (716 individuals ha⁻¹) (Tables 2 and 3, Fig. 2).

Wald chi-square tests revealed statistically significant differences in the mean counts between tree/shrub types (Table 4). Differences in the mean tree/shrub counts between treatments was statistically significant (P -value < 0.001). The post-hoc tests of treatments by tree/shrub type showed significant differences between treatments for red umbrella thorn (P -value < 0.001) (negative response to the thinning treatment), and umbrella thorn (P -value = 0.039) (positive response to the thinning treatment). The response to thinning was also positive for black-thorn acacia, and negative for sickle bush and blade thorn, although statistically insignificant. Post-thinning age was significantly and positively related to the number of tree/shrub counts. Thus, mean tree/shrub count increased by 11% concomitant with every annual increase in post-thinning age. The effect of arboricide aftercare treatment was positive on mean tree/shrub count, but was statistically insignificant.

Effect of treatment on the tree/shrub equivalent (TE) densities recommended for the area and approximate woody biomass availability. The overall densities (TE) for encroaching species measured in the non-thinned area (β_0) was, on average, 1464.05 TE ha⁻¹ (\pm 95% CI = 269.93) (Table 5a and Fig. 3a). The negative value for the effect of thinning treatment (β_1) indicates that the overall mean was significantly less (by 693.09 TE ha⁻¹; P -value = 0.001) than the baseline value (β_0). This led to a mean estimate of 770.95 TE ha⁻¹ (95% CI 128.49), which was 47.3% less than the amounts in the non-thinned area. Our results indicate that mean TE densities were 59.01% greater in the non-thinned area and 13.4% greater in the thinned area than the minimum recommended density for a 400 mm rainfall area (i.e., 600 TE ha⁻¹).

Variable	χ^2	df	Pr(> Chisq)
Tree/shrub types	114.137	4	0.000
Post-thinning age	8.798	1	0.003
Aftercare	3.477	1	0.062
Effect of treatment by tree/shrub type			
Overall	63.606	5	0.000
Sickle bush	3.862	1	0.247
Black-thorn acacia	0.575	1	1
Blade thorn	3.845	1	0.249
Red umbrella thorn	23.026	1	0.000
Umbrella thorn	7.068	1	0.039

Table 4. Hypothesis tests of the estimated generalized linear mixed-effects model (GLMM) coefficients showing overall treatment effects and differences in the abundance of encroacher species in the thinned and non-thinned treatment areas. P -values for the effect of treatment by tree/shrub type were adjusted for multiple testing using the Bonferroni method.

	Parameter (β)	(a) Density (TE ha ⁻¹)			(b) Wood biomass (kg ha ⁻¹)		
		Estimate	SE	P -value	Estimate	SE	P -value
Fixed effects							
(Intercept)	β_0	1464.05	137.1	<0.001	15,387.73	1630.468	<0.001
Treatment-thinned	β_1	- 693.095	151.841	0.001	- 10,167.4	1785.401	<0.001
Post-thinning age	β_2				639.96	227.849	0.017
Random effects							
		SD			SD		
Plot	c_{kj}	0.207			7.466		
Block	b_k	0.004			3.847		

Table 5. Linear mixed-effects (LME) models showing the effects of fixed factors (treatment, post-thinning age) on (a) tree/shrub densities expressed as TE ha⁻¹, and (b) woody biomass of encroaching trees/shrubs.




Figure 3. Expected (a) densities (TE ha^{-1}), and (b) dry wood biomass (kg ha^{-1}) of encroaching trees/shrubs in the thinned and non-thinned treatment areas estimated from the linear mixed-effects (LME) model coefficients. Error bars indicate $\pm 95\%$ confidence level of the variability in tree/shrub densities (TE ha^{-1}) and woody biomass (kg ha^{-1}) of the non-thinned and thinned areas.

Mean woody biomass of encroaching species in the non-thinned area (β_0) was 15.4 tonnes ha^{-1} ($\pm 95\%$ CI 3.2) (Table 5b and Fig. 3b). The effect of thinning treatment (β_1) resulted in a mean difference of 10.2 tonnes ha^{-1} , which was significantly less (P -value < 0.001) than the biomass in the non-thinned treatment (β_0). Hence, the overall woody biomass in the thinned area was, on average, 5.2 tonnes ha^{-1} ($\pm 95\%$ CI 1.4) and 66.1% less than the biomass in the non-thinned treatment. The effect of post-thinning age had a significant (P -value = 0.017) and positive trend on the mean woody biomass estimates, which corresponded to a 0.64 tonne ha^{-1} change in biomass concomitant with an annual increase in post-thinning age.

Effect of treatment on the vegetation structure. *Stem sizes.* The number of trees/shrubs in stem diameter class ≤ 6 cm in the non-thinned area (β_0) was, on average, 7.74 plants per subplot (684.12 plants ha^{-1}) (Tables 6 and 7, Fig. 4). The negative values for estimates in the other stem diameter classes showed that counts were less abundant than in the ≤ 6 cm diameter class: mean counts for the 6–18 cm and ≥ 18 cm classes differed by 65.2% and 93.4% times, respectively. The parameter estimate (β_3) refers to the mean counts of trees/shrubs in stem diameter class ≤ 6 cm in the thinned treatment. The low value of parameter (β_3) can be directly interpreted as the average relative change (1.2%) from the counts in the ≤ 6 cm diameter size class in the non-thinned area (non-thinned = 7.74, thinned = 7.65) (Tables 6 and 7). In comparison to the non-thinned area, mean counts in the thinned area were 74% and 68.3% lower for the 6–18 cm and ≥ 18 cm diameter classes, respectively (Tables 6 and 7, Fig. 4). Overall, our predictions showed that the majority (79.1%) of the measured stem diameters were in the ≤ 6 cm diameter class.

Wald chi-square tests revealed statistically significant differences in the average counts among the different stem classes (P -value < 0.001) (Table 8). Differences in the mean tree/shrub counts per stem size between

Variable	Parameter (β)	Value	SE	exp (β)
Fixed part				
(Intercept)	β_0	2.046	0.133	7.737
6–18 cm	β_1	– 1.057	0.078	0.348
≥ 18 cm	β_2	– 2.723	0.160	0.066
Treatment thinned	β_3	– 0.012	0.441	0.988
Post-thinning age	β_4	0.087	0.055	1.091
Arbicide aftercare applied	β_5	0.250	0.494	1.284
6–18 cm: thinned	β_6	– 1.336	0.140	0.263
≥ 18 cm: thinned	β_7	– 1.138	0.284	0.321
Random part				
		SD		
Subplot	d_{kji}	0.316		
Plot	c_{kj}	0.315		
Block	b_k	0.309		

Table 6. Generalized linear mixed-effects model (GLMM) showing the effects of fixed factors (treatment, post-thinning age, and arbicide aftercare) on the abundance of encroaching trees/shrubs measured in stem classes, 0–6, 6–18 and ≥ 18 cm.

Treatment/stem diameter class	Model coefficients (β)	Sum (β)	Mean counts ($\exp(\text{sum}(\beta))$)	\pm 95% confidence level
Non-thinned				
≤ 6 cm	$\beta_0 + \beta_3(0)$	2.046	7.737	1.297
6–18 cm	$\beta_0 + \beta_1(1) + \beta_3(0) + \beta_6(0)$	0.989	2.69	1.512
> 18 cm	$\beta_0 + \beta_2(1) + \beta_3(0) + \beta_7(0)$	- 0.677	0.508	1.775
Thinned				
≤ 6 cm	$\beta_0 + \beta_3(1)$	2.034	7.648	3.454
6–18 cm	$\beta_0 + \beta_1(1) + \beta_3(1) + \beta_6(1)$	- 0.358	0.699	5.297
> 18 cm	$\beta_0 + \beta_2(1) + \beta_3(1) + \beta_7(1)$	- 1.827	0.161	8.239

Table 7. Expected tree/shrub density per subplot (area: 113.1 m²) in the non-thinned and thinned treatment areas estimated from the generalized linear mixed-effects model (GLMM) coefficients in stem classes, 0–6 cm., 6–18 cm. and ≥ 18 cm.




Figure 4. Expected tree/shrub density per hectare in the non-thinned and thinned treatment areas estimated from the generalized linear mixed-effects model (GLMM) model coefficients in stem classes, 0–6 cm, 6–18 cm and ≥ 18 cm. Error bars indicate \pm 95% confidence level of the variability in tree/shrub densities of the three measured stem classes of the non-thinned and thinned areas.

treatments were significant (P -value < 0.001); the post-hoc tests of treatment by stem diameter size showed significant differences between treatments in the 6–18 cm stem diameter class (P -value = 0.009), with a negative response to the thinning treatment. A negative response to thinning was also observed in the ≤ 6 cm and ≥ 18 cm stem diameter classes, but were statistically insignificant. Both post-thinning age and aftercare did not have any significant effect on the number of tree/shrubs present in each stem diameter class.

Vegetation structure (vertical, horizontal) and habitat sighting lines. The mean height of trees/shrubs in the non-thinned area (β_0) was 1.9 m (\pm 95% CI 2.1) (Table 9a and Fig. 5a). The effect of thinning treatment (β_1) resulted in a significant (P -value < 0.001) reduction in height (1 m) compared to the non-thinned area (β_0). Thus, for the thinned area a mean height of 0.9 m (\pm 95% CI 0.21) was predicted.

The overall canopy area measured per subplot (area: 113.1 m²) in the non-thinned area (β_0) was, on average, 50.27 m² (\pm 95% CI 10.51) (Table 9b and Fig. 5b). Thinning treatment (β_1) resulted in a significant (P -value = 0.001) reduction in canopy area (28.64 m²) in relation to the non-thinned (β_0) area. As such, mean

Variable	χ^2	Df	P -value
Stem diameter class	422.357	2	< 0.001
Post-thinning age	2.511	1	0.113
Aftercare	0.258	1	0.611
Effects of treatment by stem diameter class			
Overall	103.17	3	< 0.001
≤ 6 cm	0.001	1	1
6–18 cm	8.762	1	0.009
≥ 18 cm	4.934	1	0.079

Table 8. Hypothesis tests of the estimated generalized linear mixed-effects model (GLMM) model coefficients showing overall treatment effects and differences in the abundance of encroacher species in stem classes 0–6 cm, 6–18 cm and ≥ 18 cm. P -values for the effect of treatment by tree/shrub counts in the different stem classes were adjusted for multiple testing using the Bonferroni method.

	Parameter (β)	(a) Heights			(b) Canopy area			(c) Sighting lines		
		Estimate	Std. error	<i>P</i> -value	Estimate	Std. error	<i>P</i> -value	Estimate	Std. error	<i>P</i> -value
Fixed effects										
(Intercept)	β_0	1.9	0.108	<0.001	50.266	5.339	<0.001	24.773	2.776	<0.001
Treatment-thinned	β_1	-0.996	0.15	<0.001	-28.642	6.914	0.001	32.618	3.733	<0.001
Post-thinning age	β_2							-4.937	0.902	<0.001
Random effects										
		SD			SD			SD		
Subplot	d_{kji}	0.427								
Plot	c_{ij}	0.347			14.546			0.014		
Block	b_k	0.001			0.014			9.015		

Table 9. Linear mixed effects (LME) models showing the effects of fixed factors (treatment, post-thinning age) on the vegetation structure (height, canopy area) of encroaching trees/shrubs, as well as on sighting lines in the thinned and non-thinned treatment areas.




Figure 5. Expected mean (a) height (m), and (b) canopy area (m^2) of bush-encroaching trees/shrubs, as well as (c) sighting lines (m) in the thinned and non-thinned treatment areas estimated from the linear mixed effects (LME) model coefficients. Error bars indicate $\pm 95\%$ confidence level of the variability in tree/shrub height, canopy area and sighting lines of the non-thinned and thinned areas.

canopy area in the thinned area was $21.62 m^2 (\pm 95\% CI 8.65)$ per subplot, which was equivalent to a 57% reduction in woody canopy cover of encroaching species compared to the non-thinned area. Correspondingly, the woody canopy cover per subplot in the non-thinned area equates to $44.4\% (\pm 95\% CI 9.30)$; for the thinned area $19.1\% (\pm 95\% CI 7.65)$. The estimates showed greater random variability in the average tree/shrub heights and canopy areas amongst the plots than between blocks (plot pairs).

For habitat sighting lines, a total of 1173 measurements were taken in 293 subplots, of which 50.6% were in the thinned and 49.5% were in the non-thinned areas. The mean sighting line measured in the non-thinned area was $24.8 m (\pm 95\% CI 8.73 m)$ (Table 9c and Fig. 5c). The effect of thinning treatment (β_1) resulted in a greater mean sighting line of $57.4 m (\pm 95\% CI 5.47 m)$: a statistically significant (P -value < 0.001) difference of 32.62 m. Post-thinning age (β_2) had a significant and negative effect on the sighting lines. Thus, mean sighting lines would decrease by 4.94 m concomitant with every annual increase in post-thinning age.

Discussion

Number of individuals and observed species-treatment interactions. Our results indicate that sickle bush had the greatest mean abundance, followed by black-thorn acacia, blade-thorn and red umbrella thorn. Non-thinned plots had the greatest abundance of trees/shrubs, and overall tree/shrub-treatment interactions were significant. In non-thinned areas, the abundance of the main encroacher species, sickle bush and black-thorn acacia (Tables 2 and 3) were lower than the average estimates for encroached habitat in this region^{3,57}. We expect those regional estimates were influenced by factors that included rainfall, soil type, fire history, number of sample sites and management of the area, which could cause inaccuracies in extrapolations over larger areas. Our data provides detailed information regarding the extent of bush encroachment, as well as the effects of thinning at the localised level.

With regard to red umbrella thorn, a negative and significant response to thinning was observed; umbrella thorn had a positive and significant response to thinning. Contrary to our expectations, the most abundant species, sickle bush, was not reduced significantly by thinning, possibly due to the avoidance of small-sized stems⁵⁸, and this may explain the substantial reduction in red umbrella thorn as it was structurally larger (Table 1). The

significantly greater abundance of umbrella thorn in the thinned area shows that thinning dramatically accelerated regeneration of this species, possibly due to reduced competition and also because the retained trees/shrubs may have increased their reproductive output^{33,59}. These results are consistent with those reported by Nghikembua et al., (2021) who found that the occurrence probabilities for red umbrella thorn were significantly less, and were greater for umbrella thorn in the same thinned plots.

The response to thinning was not significant for other species, which may be partly due to the herbicide application that prevented regeneration via coppicing. Also, as species abundance at the mean 7.2-year mark time since thinning were still poor, this may indicate that a longer regeneration period is required for these species. Inventories of bush control operations performed elsewhere have shown that in the absence of aftercare or timely aftercare applications, re-encroachment can occur within a short period (<5 years) due to coppicing stumps and root suckering^{2,21}. Moreover, in these areas, regardless of chemical aftercare, mechanically cleared sites showed greater levels of re-encroachment, possibly due to soil disturbance, delayed timing of the chemical aftercare, and greater probability of leaving stumps untreated.

A slight increase in black-thorn acacia counts by 21.7% was observed in the thinned plots. According to Joubert et al., (2013), significantly higher seed production and sapling establishment for this species were positively related with exceptional high rainfall seasons (at least two consecutive years). Also, survival rates of seedlings were five times greater away from trees (in open gaps) than under trees. It is likely that recruitment from seed occurred in our study area during 2008–2012 when exceptional, above average, annual rainfall (>444 mm) was recorded⁴⁵.

Tree/shrub equivalent densities (TE) recommended for the area and approximate woody biomass availability.

The mean overall TE density for the non-thinned area showed a moderate level of encroachment and exceeded twice the recommended threshold for the area. The estimated density was lower than in other areas of the country where ~6,000 TE ha⁻¹ has been observed⁶⁰. Also, overall woody biomass (Table 5b) in the non-thinned areas was much less than the regional average (36.2 tonnes ha⁻¹)²¹. However, the estimates in our study correspond closely with previous studies conducted on the same farms^{61,62}. Possible explanations for reduced woody biomass volumes include differences in species composition, soil type, climatic factors, management history, as well as methodologies applied in biomass estimation^{35,36,39}. The sickle bush specimens that were dominant in our study area were structurally smaller (Table 1), which would also affect the total woody biomass value. Thinning treatment was effective in causing a significant reduction in tree/shrub densities, well within the recommended levels (600–750 TE ha⁻¹) for the area. Correspondingly, woody biomass was significantly reduced. The mean difference of 10.2 tonnes ha⁻¹ dry woody biomass between treatments suggests that this amount may have been removed during the thinning operations. Based on these estimates, we expect that our study area of 17,422.44 ha contains ~268,091.8 tonnes (equates to 0.268 Mt) of woody biomass. The Namibian Ministry of Environment, Forestry and Tourism (MEFT) Guidelines highlight the importance of avoiding complete clearance of trees/shrubs by retaining a minimum density consistent with the rainfall received in the area². During the thinning operations, harvesters are trained to identify target and protected species, to select harvestable size classes, and to leave small clumps of undisturbed vegetation to provide browsing and cover for biodiversity.

Effect of treatment on vegetation structure and sighting lines. Thinning was effective in modifying the vegetation structure as shown by the significant differences between the treatments in vegetation structure and the abundance of small-medium sized trees/shrubs (6–18 cm stem class). It was evident that following thinning, the overall mean height, canopy size and abundance of trees/shrubs were significantly less than in the non-thinned areas. Moreover, thinned areas exhibited longer sighting lines, which were significantly different to the non-thinned areas (Table 9). The obvious explanation for these results is the considerable reduction in the overall tree/shrub abundance (Tables 2 and 3) in the thinned plots, with the number of individuals in the 1–3 m and >3 m height classes significantly less than in the non-thinned areas⁴⁵.

The mean sighting line (57.4 m) in the thinned areas in this study is less than those measured in sparsely vegetated areas preferred by cheetahs reported in other studies^{4,52} in the same region. The shorter mean sightline in the thinned areas in our study could be due partly to the fact that thinning was carried out conservatively in this area with the aim to retain specific tree/shrub densities rather than the densities that would result from natural disturbances (e.g. fire, natural dieback) or when bush removal was carried out for different management objectives (e.g. clearing for a grass fields). Also, our predictions show that sighting lines would decline with progressing post-thinning age and may take ~14 years to equalise between treatments. One possible explanation for this occurrence is the rapid natural regeneration present in the same study area due to re-encroachment by young cohorts (0–1 m height class)⁴⁵. Another possible explanation for reduced sighting lines is the accelerated growth of structural features (e.g. canopy, stems) due to the absence of competition amongst the retained vegetation^{10,33,59,63,64}.

A reduction in tree/shrub densities would be advantageous as it would restore the soil moisture balance, reduce the suppression effects of woody plants on the herbaceous layer, thereby increasing the likelihood for greater grazing capacity and ease of movement as animals manoeuvre through the habitat^{1,3,4,12,21,65–67}. This may explain why thinned areas were strongly frequented by wildlife in the same plots in another study²⁸.

Conclusion and management recommendations

Our results indicate that thinning was effective in reducing the overall abundance of trees/shrubs, which was still evident at the average 7.2-year mark since thinning and when herbicide aftercare was applied. Of the measured trees/shrubs, red umbrella thorn was significantly reduced, and umbrella thorn significantly increased in the

thinned areas. This suggests that thinning has the potential to change the floristic composition of species within an ecosystem, as preferred species are reduced (e.g. red umbrella thorn) and low-density species are promoted (e.g. umbrella thorn). The effect of thinning treatment was not significant for sickle bush, blade thorn and black-thorn acacia. However, the difference in expected counts for these species may indicate the true thinning treatment effect, which is just too small to be significant with our sample size.

This study has revealed that trees/shrubs can be selectively harvested, well within the recommended densities (600–750 TE ha⁻¹) for the area. The average difference in woody biomass between the treatments (10.4 tonnes ha⁻¹) suggests that this amount could be harvested within the recommended tree/shrub density range for the area. Thinning also modified the vegetation structure; tall, dense shrubland with limited sighting lines in the non-thinned areas was restored to short and low-cover shrubland with greater sighting lines. We anticipate that a low woody canopy cover and tree/shrub abundance would restore the soil moisture balance, promote perennial grass cover, and increase the grazing capacity. Also, the sparsely vegetated thinned areas would be an ideal habitat for wildlife that rely on longer sighting lines when hunting or evading predators.

This study revealed important findings regarding the effects of treatment on the encroaching woody vegetation and success of restoration thinning. However, it is not without any limitations. Firstly, there is uncertainty as to how climate change induced weather patterns (e.g., sporadic rainfall, increased drought frequencies) could influence the regeneration rate of woody species and grass cover, which could determine the restoration success and its duration. For example, enhanced plant growth, seed production and recruitment of saplings are expected during years of exceptional rainfall⁶⁸. Over the long term, and without post thinning management, this could further promote bush encroachment, reduce sighting lines and grazing capacity. Equally, during drought years, poor plant growth, less grass cover and more overgrazing are expected and could further promote woody seedling establishment in subsequent above average rainfall years due to the loss of the herbaceous layer³. The uncertainties of these weather patterns demand better adaptive rangeland management strategies consistent with climate change including regulating livestock stocking rates, rotation grazing, and post thinning management of previously thinned areas. Long term monitoring into the patterns of the vegetation structure in relation to treatment (thinned, non-thinned) is necessary, particularly in response to climate change induced weather patterns.

Secondly, the results were limited to the study period, number of plots observed and a single thinning methodology; therefore, longer-term data, replicating the study over larger areas (multiple farms) and stratifying different levels of thinning are required to fully evaluate the effects on the vegetation structure. For future research and assessment, we recommend: (1) long term monitoring of regrowth to reveal trends and rates of regeneration by species, (2) thinning of established saplings in restored areas to prevent loss of longer sighting lines and re-encroachment, (3) development of allometric equations to predict available biomass, and (4) ongoing restoration to control bush encroachment to increase rangeland productivity and suitable habitats for species that prefer an open savannah vegetation structure.

Data availability

The datasets used in deriving the results during the current study are available from the corresponding author on reasonable request.

Received: 11 April 2022; Accepted: 19 December 2022

Published online: 06 January 2023

References

1. MAWF. *National Rangeland Management Policy & Strategy. Restoring Namibia's Rangelands* (2012).
2. SAIEA. *Strategic environmental assessment of large-scale bush thinning and value-addition activities in Namibia*. (2016).
3. de Klerk, J. N. *Bush Encroachment in Namibia. Report on Phase 1 of the Bush Encroachment Research, Monitoring and Management Project*. (2004).
4. Muntifering, J. R. *et al.* Managing the matrix for large carnivores: A novel approach and perspective from cheetah (*Acinonyx jubatus*) habitat suitability modelling. *Anim. Conserv.* **9**, 103–112 (2006).
5. Marker, L. L., Dickman, A. J., Mills, M. G. L., Joo, R. M. & Macdonald, D. W. Spatial ecology of cheetahs on north-central Namibian farmlands. *J. Zool.* **274**, 226–238 (2008).
6. Wang, J. *et al.* Impacts of juniper woody plant encroachment into grasslands on local climate. *Agric. For. Meteorol.* **307**, 108508 (2021).
7. Shen, X. *et al.* Effect of shrub encroachment on land surface temperature in semi-arid areas of temperate regions of the Northern Hemisphere. *Agric. For. Meteorol.* **320**, 108943 (2022).
8. Shen, X. *et al.* Vegetation greening, extended growing seasons, and temperature feedbacks in warming temperate grasslands of China. *J. Clim.* **35**, 5103–5117 (2022).
9. Martins, A. R. O. & Shackleton, C. M. Population structure and harvesting selection of two palm species (*Hyphaene coriacea* and *Phoenix reclinata*) in Zitundo area, southern Mozambique. *For. Ecol. Manage.* **398**, 64–74 (2017).
10. Brown, G. W., Murphy, A., Fanson, B. & Tolsma, A. The influence of different restoration thinning treatments on tree growth in a depleted forest system. *For. Ecol. Manage.* **437**, 10–16 (2019).
11. Belsky, A. Influences of trees on savanna productivity: Tests of shade, nutrients and grass–tree competition. *Ecology* **75**, 922–932 (1994).
12. Hagos, M. G. & Smit, G. N. Soil enrichment by *Acacia mellifera* subsp. *detinens* on nutrient poor sandy soil in a semi-arid southern African savanna. *J. Arid Environ.* **61**, 47–59 (2005).
13. Ludwig, F., Kroon, H. D., Berendse, F. & Prins, H. H. T. The influence of savanna trees on nutrient, water and light availability and the understorey vegetation. *Plant Ecol.* **170**, 93–105 (2004).
14. Ridolfi, L., Laio, F., D'Odorico, P. & D'Odorico, P. Fertility island formation and evolution in dryland ecosystems. *Ecol. Soc.* **13**, 13 (2008).
15. Wiegand, K., Ward, D. & Saltz, D. Multi-scale patterns and bush encroachment in an arid savanna with a shallow soil layer. *J. Veg. Sci.* **16**, 311–320 (2005).
16. Burke, A. Savanna trees in Namibia - Factors controlling their distribution at the arid end of the spectrum. *Flora Morphol. Distrib. Funct. Ecol. Plants* **201**, 181–201 (2006).

17. Buyer, J. S., Schmidt-Küntzel, A., Nghikembua, M., Maul, J. E. & Marker, L. Soil microbial communities following bush removal in a Namibian savanna. *Soil* **2**, 101–110 (2016).
18. Dwivedi, V. & Soni, P. A review on the role of soil microbial biomass in eco-restoration of degraded ecosystem with special reference to mining areas. *J. Appl. Nat. Sci.* 151–158 (2011). <https://doi.org/10.31018/jans.v3i1.173>.
19. Smit, G. N. An approach to tree thinning to structure southern African savannas for long-term restoration from bush encroachment. *J. Environ. Manage.* **71**, 179–191 (2004).
20. MAWE. Forestry and environmental authorisation process for bush harvesting projects. 34 (2017).
21. Smit, G. N., de Klerk, J. N., Schneider, M. B. & van Eck, J. Detailed assessment of the biomass resource and potential yield in a selected bush encroached area of Namibia. 141 (2015).
22. Marker, L. *et al.* and Distribution. in *Cheetahs: Biology and Conservation: Biodiversity of the World: Conservation from Genes to Landscapes* (eds. Marker, L., Boast, L. k. & Schmidt-Küntzel, A.) 9–20 (John Fedor, 2018). <https://doi.org/10.1016/B978-0-12-804088-1.00004-6>.
23. NAPHA. Namibia Professional Hunting Association. <http://www.napha-namibia.com/conservation/hunttable-species/carnivora/> (2015).
24. SWA. *Nature Conservation Ordinance, 1975 (No. 4 of 1975)*. vol. 1975 (1975).
25. Marker, L. *et al.* The status of Key pre species and the Consequences of Prey Loss for Cheetah Conservation in North and West Africa. in *Cheetahs: Biology and Conservation: Biodiversity of the World: Conservation from Genes to Landscapes*. (eds. Marker, L., Boast, L. k. & Schmidt-Küntzel, A.) 151–161 (John Fedor, 2018). <https://doi.org/10.1016/B978-0-12-804088-1.00004-6>.
26. Kiruki, H. M., van der Zanden, E. H., Gikuma-Njuru, P. & Verburg, P. H. The effect of charcoal production and other land uses on diversity, structure and regeneration of woodlands in a semi-arid area in Kenya. *For. Ecol. Manage.* **391**, 282–295 (2017).
27. Harmse, C. J., Kellner, K. & Dreber, N. Restoring productive rangelands: A comparative assessment of selective and non-selective chemical bush control in a semi-arid Kalahari savanna. *J. Arid Environ.* **135**, 39–49 (2016).
28. Nghikembua, M. T. *et al.* Response of wildlife to bush thinning on the north central freehold farmlands of Namibia. *For. Ecol. Manage.* **473**, 118330 (2020).
29. Soto-Shoender, J. R., McCleery, R. A., Monadjem, A. & Gwinn, D. C. The importance of grass cover for mammalian diversity and habitat associations in a bush encroached savanna. *Biol. Conserv.* **221**, 127–136 (2018).
30. Strohbach, B. J. Environmental information service, Namibia for the Ministry of Environment and Tourism, the Namibian Chamber of Environment and the Namibia University of Contribution to the knowledge of southern African Lepismatidae. *Namibian J. Environ.* **8327**, 14–33 (2017).
31. Smit, N. BECVOL 3: An expansion of the aboveground biomass quantification model for trees and shrubs to include the wood component. *Afr. J. Range Forage Sci.* **31**, 179–186 (2014).
32. Zimmermann, I. Causes and Consequences of Fenceline Contrasts in Namibia. (Free State, Bloemfontein, South Africa, 2009).
33. Dwyer, J. M. & Mason, R. Plant community responses to thinning in densely regenerating Acacia harpophylla forest. *Restor. Ecol.* **26**, 97–105 (2018).
34. Thomas, S. C. & Martin, A. R. Carbon content of tree tissues: A synthesis. *Forests* **3**, 332–352 (2012).
35. Chave, J. *et al.* Improved allometric models to estimate the aboveground biomass of tropical trees. *Glob. Chang. Biol.* **20**, 3177–3190 (2014).
36. Djomo, A. N. & Chimi, C. D. Tree allometric equations for estimation of above, below and total biomass in a tropical moist forest: Case study with application to remote sensing. *For. Ecol. Manage.* **391**, 184–193 (2017).
37. Ganamé, M., Bayen, P., Dimobe, K., Ouédraogo, I. & Thiombiano, A. Aboveground biomass allocation, additive biomass and carbon sequestration models for Pterocarpus erinaceus Poir. in Burkina Faso. *Heliyon* **6**, (2020).
38. Picard, N., Saint-André, L. & Henry, M. *Manual for building tree volume and biomass allometric equations: From field measurement to prediction*. Food and Agricultural Organization of the United Nations, Rome, and Centre de Coopération Internationale en Recherche Agronomique pour le Développement. (2012).
39. Feyisa, K. *et al.* Allometric equations for predicting above-ground biomass of selected woody species to estimate carbon in East African rangelands. *Agrofor. Syst.* **92**, 599–621 (2018).
40. Boys, J. M. & Smit, N. G. *Development of an Excel Based Bush Biomass Quantification Tool*. (2020).
41. Ngomanda, A. *et al.* Site-specific versus pantropical allometric equations: Which option to estimate the biomass of a moist central African forest?. *For. Ecol. Manage.* **312**, 1–9 (2014).
42. CCF. Cheetah Conservation Fund Bush PTY (Ltd). <https://bushblok.com/management-plan> (2019).
43. Wykstra, M. *et al.* Improved and Alternative Livelihoods: Links Between Poverty Alleviation, Biodiversity, and Cheetah Conservation. in *Cheetahs: Biology and Conservation: Biodiversity of the World: Conservation from Genes to Landscapes* (eds. Marker, L., Boast, L. k. & Schmidt-Küntzel, A.) 223–237 (John Fedor, 2018).
44. Zimmermann, I. *et al.* The influence of two levels of debushing in Namibia's Thornbush Savanna on overall soil fertility, measured through bioassays. *Namibian J. Environ.* **1**, 52–59 (2017).
45. Nghikembua, M. T. *et al.* Restoration thinning reduces bush encroachment on freehold farmlands in north-central Namibia. *For. An Int. J. For. Res.* 1–14 (2021). <https://doi.org/10.1093/forestry/cpab009>.
46. Fick, S. E. & Hijmans, R. J. WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *Int. J. Climatol.* **37**, 4302–4315 (2017).
47. Mendelsohn, J., Jarvis, A., Roberts, C. & Robertson, T. *Atlas of Namibia: A portrait of the land and its people*. (2003).
48. Curtis, B. A. & Mannheimer, C. A. *Tree Atlas of Namibia; National Botanical Research Institute (NBRI)*. (National Botanical Research Institute, 2005).
49. Mannheimer, C. & Curtis, B. *Le Roux and Muller's Field Guide to the Trees & Shrubs of Namibia*. (Macmillan Education Namibia (PTY) LTD, 2009).
50. Honsbein, D., Shiningavamwe, K., Iikela, J. & de la Puerta Fernandez, Maria, L. *Animal Feed from Namibian Encroacher Bush*. (2017).
51. Coates Palgrave, K. *Trees of Southern Africa*. (Struik publishers, 1993).
52. Nghikembua, M., Harris, J., Tregenza, T. & Marker, L. Spatial and temporal habitat use by GPS collared male cheetahs in modified bushland habitat. *Open J. For.* **06**, 269–280 (2016).
53. Venables, W. N. & Ripley, B. D. *Modern Applied Statistics with S*. (Springer, 2002).
54. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* **67**, 1–48 (2015).
55. Mehtatalo, L. & Lappi, J. *Forest biometrics with examples in R*. (Taylor & Francis Inc, 2020).
56. R Core Team. R: A language and environment for statistical computing (2021).
57. Birch, C. & Middleton, A. *Economics of Land Degradation Related To Bush Encroachment in Namibia*. (2017).
58. Neke, K. S., Owen-Smith, N. & Witkowski, E. T. F. Comparative resprouting response of Savanna woody plant species following harvesting: the value of persistence. *For. Ecol. Manage.* **232**, 114–123 (2006).
59. Smit, N. Response of Colophospermum mopane to different intensities of tree thinning in the Mopane Bushveld of southern Africa. *Afr. J. Range Forage Sci.* **31**, 173–177 (2014).
60. DAS. *Bush Control Manual*. (John Meinert Printing, 2017).
61. Leinonen, A. *Wood chip production technology and costs for fuel in Namibia*. VTT Tiedotteita - Valtion Teknillinen Tutkimuskeskus (2007).

62. Chakanga, M. *A preliminary analysis of the Economic Plots Research Data of the CCF Bush Project*. (2003).
63. West, P. W. Thinning. in *Growing Plantation Forests* vol. 9783319018 115–129 (Springer International Publishing, 2014).
64. Dwyer, J. M., Fensham, R. & Buckley, Y. M. Restoration thinning accelerates structural development and carbon sequestration in an endangered Australian ecosystem. *J. Appl. Ecol.* **47**, 681–691 (2010).
65. Groengroeft, A., de Blécourt, M., Classen, N., Landschreiber, L. & Eschenbach, A. Acacia trees modify soil water dynamics and the potential groundwater recharge in savanna ecosystems. in *Climate change and adaptive land management in southern Africa - assessments, changes, challenges, and solutions* (ed. eds. Revermann, R. et al.) 177–186 (Klaus Hess Publishers, 2018).
66. Richter, C. G. F., Snyman, H. A. & Smit, G. N. The influence of tree density on the grass layer of three semi-arid savanna types of southern africa. *Afr. J. Range Forage Sci.* **18**, 103–109 (2001).
67. Stafford, W. *et al.* The economics of landscape restoration: Benefits of controlling bush encroachment and invasive plant species in South Africa and Namibia. *Ecosyst. Serv.* **27**, 193–202 (2017).
68. Joubert, D. F., Smit, G. N. & Hoffman, M. T. The influence of rainfall, competition and predation on seed production, germination and establishment of an encroaching Acacia in an arid Namibian savanna. *J. Arid Environ.* **91**, 7–13 (2013).

Acknowledgements

We thank Cheetah Conservation Fund (CCF) and CCF Bush (Pty Ltd), who provided all the logistical support and access to the farmland for research; CCF staff: Johan Britz and Paul Visser for transport logistics arrangements, Richard Siririka and David Shipingana for field work and data entry; CCF volunteers; Adolf Honibeb, Leinhold Kashikola, Werner Iilundwa and Ebenezer Tsibeb for help with the fieldwork; MAWF for providing research and harvest work permits.

Author contributions

M.T.N. is the primary author, responsible for project conceptualisation, data collection, data analysis and writing. The experimental design was planned together with A.P., L.M. and M.A. Statistical data analysis was supported by L.M. Individual contributions were as follows: L.M., project conceptualisation, critical review, editing. B.B., project conceptualisation, critical review, editing. A.L., critical comments and editing of the papers. M.A., advice, guidance, editing of the papers. L.M., statistical analysis and editing of the papers. A.P.: guidance and advice regarding the study design of this research. All authors reviewed the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-022-26639-4>.

Correspondence and requests for materials should be addressed to A.P.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023