



Received: 14 February 2026
Accepted: 27 March 2026
Published: 10 April 2026
Edited by: Frowin Becker

A teratological case of Monteiro's hornbill (*Tockus monteiri*) with beak deformity in Namibia (Aves: Bucerotiformes: Bucerotidae)

Filippo Ceccolini

Via Europa, Rassina (Arezzo), Italy

Correspondence: ceccolinif@virgilio.it

ABSTRACT This note describes a case of beak deformity in a specimen of Monteiro's hornbill (*Tockus monteiri* Hartlaub, 1865) observed in north-central Namibia. The deformity consisted of an elongated upper mandible. Although this teratology presumably makes it more difficult to find food and maintain its feathers, the observed bird had reached adult size and appeared to be healthy.

KEYWORDS Avian Keratin Disorder; beak abnormality; elongated beak; Monteiro's hornbill; Namibia; rhamphotheca

INTRODUCTION

The rhamphotheca, the outer cornified layer of a bird's beak, plays an important role in several aspects of its life, such as feeding (Navalón et al. 2019), feather maintenance and parasite control (Clayton et al. 2005), social behaviour (Rogers & Kaplan 2000), and sometimes even navigation (Wiltschko & Wiltschko 2013). Deformities of this structure (or teratology) have been recorded in wild specimens of several bird species around the world (e.g. Pomeroy 1962, Craves 1994) and may be either permanent or temporary (Pomeroy 1962). Several causes can induce deformities, either genetic mutations or environmental factors, such as nutritional deficiencies, injuries, contact with chemical pollutants and viral infection (see Pomeroy 1962, Craves 1994, Veltri & Klem 2005, Handel & Van Hemert 2014, Zylberberg et al. 2016, 2018, 2021).

Many cases of beak deformity are recorded as isolated incidents (e.g. Gallo-Ortiz 2011, Rezende 2013, dos Santos et al. 2018, Hodges et al. 2019,

Crozariol 2020, Íthalo et al. 2021, de Moura et al. 2022, Tinajero 2023, Tenez 2025), but some can be traced back to epizootic phenomena. One remarkable case is that of wild birds in Alaska, where since the 1990s beak deformities have been recorded in more than 2 500 specimens of 30 bird species (Handel et al. 2010, Van Hemert & Handel 2010). An especially high frequency was documented in Black-capped Chickadee (*Poecile atricapillus*) populations (Handel et al. 2010, Van Hemert et al. 2012, Zylberberg et al. 2016, 2018, 2021), in which an epizootic has been suggested to cause crossed or elongated beaks. This has been termed Avian Keratin Disorder (AKD), and it has afflicted an average of 6.5% of the adult population annually over the course of a decade (Handel et al. 2010).

In related literature most cases of beak teratology are described in American birds (e.g. Handel et al. 2010, Van Hemert & Handel 2010, Van Hemert et al. 2012, 2025, Bianchini & Arenas 2014, 2018, Gorosito et al. 2016, dos Santos et al. 2018, Valdebenito et al. 2018, Zylberberg et al. 2016, 2018,



Figure 1 Map of Namibia with the observation site indicated in red.

2021, Hodges et al. 2019, Purificação, 2019, Smith et al. 2019, Íthalo et al. 2021) and few cases have been reported from the other continents, with some notable exceptions, such as Harrison (2011) for the

United Kingdom. In this note a case of beak abnormality is described, concerning a specimen of Monteiro's hornbill (*Tockus monteiri* Hartlaub, 1865) in Namibia - a species which is native to dry thorn savanna in south-western Africa (Simmons 1997, Barnes et al. 2024).

OBSERVATION

A teratological specimen of Monteiro's hornbill was observed on the morning of 22 August 2025 (at around 06h45, local time) at the Eagle Tented Lodge and Spa, located within Kunene Region, south of Etosha National Park, Namibia (Figure 1). The coordinates of the location of observation are 19.3853° S, 15.6870°E.

This specimen showed a notable beak alteration, with the upper mandible substantially elongated (Figure 2a). It was observed for some minutes on a branch a few meters away from a conspecific (Figure 2b). Despite this deformity, the bird appeared to be healthy, with well-preened plumage, and similar in size to other conspecifics.




Figure 2 a) The Monteiro's hornbill with the elongated upper mandible; b) A conspecific from the same location with a normal upper mandible (photos: F. Ceccolini).

DISCUSSION

Attention to AKD cases has grown in recent years, as demonstrated by the global citizen science project on birds with deformed beaks, which was launched by iNaturalist in 2019 (iNaturalist 2026). However, there are still only few cases of beak teratologies reported in Africa. In addition to the documented presence of a *Circovirus* that causes Psittacine Beak and Feather Disease (Pbfd) in parrots and other birds and that can induce beak deformities and other pathologies (Heath et al. 2004, Downs et al. 2015, Regnard et al. 2015), the known records come mainly from few countries including Algeria (Belkacem et al. 2025) and South Africa (Jones et al. 2015).

I found no other cases of beak deformity of Monteiro's hornbill in the literature, nor on iNaturalist (2026). In this note I have described, to the best of my knowledge, the first occurrence of a beak abnormality in this species, which can serve as a basis for possible future observations. Based on my observation, it was not possible to establish the etiology (genetic or viral) of the keratin disorder described. Only future observations and more detailed analyses (through clinical observation, molecular testing, and histopathology) of similar records in other Monteiro's hornbill specimens of the region could help to draw possible links epizootic origins. Previous studies have shown that avian keratin disorder frequently results in severe health consequences (often lethal) for affected birds (Van Hemert et al. 2012).

REFERENCES

- Barnes K, Stevenson T, Fanshawe J (2024) *Birds of Greater Southern Africa*. Helm, Dublin, Ireland.
- Belkacem M, Boulaouad BM, Mairif M, Djetti T, Harzallah B (2025) Documentation of avian deformities in Algeria. *Avocetta* 49: 1–16. <https://doi.org/10.30456/avo.28753>
- Bianchini M, Arenas C (2014) Registros documentados de aves con picos deformados en Argentina. *Nótulas Faunísticas* 233: 1–11.
- Bianchini M, Arenas C (2018) Registros documentados de aves con picos deformados hallados en la Argentina. *Nuestras Aves* 59: 12–13. <https://doi.org/10.56178/na.vi59.290>
- Clayton DH, Moyer BR, Bush SE, Jones TG, Gardiner DW, Rhodes BB, Goller F (2005) Adaptive significance of avian beak morphology for ectoparasite control. *Proceedings of the Royal Society B* 272(1565): 811–817. <https://doi.org/10.1098/rspb.2004.3036>
- Craves JA (1994) Passerines with deformed bills. *North American Bird Bander* 19: 14–18.
- Crozariol MA (2020) Deformidade no bico do cardeal-do-nordeste *Paroaria dominicana* (Passeriformes: Thraupidae). *Atualidades Ornitológicas* 213: 26–27.
- de Moura AS, de Macedo AS, Machado FS, Fontes MAL (2022) Primeiro registro de deformidade de bico em “Pipira-preta” (*Tachyphonus rufus*) Passeriformes: Thraupidae. *Brazilian Journal of Science* 1(9): 56–60.
- dos Santos LES, Wagener TLS, Almeida RS (2018) Registro de deformação no bico de saracura-do-mato, *Aramides saracura* (Gruiformes: Rallidae). *Atualidades Ornitológica* 201: 26.
- Downs CT, Brown M, Hart L, Symes CT (2015) Review of documented beak and feather disease virus cases in wild Cape parrots in South Africa during the last 20 years. *Journal of Ornithology* 156(4): 867–875. <https://doi.org/10.1007/s10336-015-1258-6>
- Handel CM, Van Hemert C (2014) Environmental contaminants and chromosomal damage associated with beak deformities in a resident North American passerine. *Environmental Toxicology and Chemistry* 34(2) (2015): 314–327. <https://doi.org/10.1002/etc.2799>
- Handel CM, Pajot LM, Matsuoka SM, Van Hemert C, Terenzi J, Talbot SL, Mulcahy DM, Meteyer CU, Trust KA (2010) Epizootic of beak deformities among wild birds in Alaska: an emerging disease in North America? *The Auk* 127(4): 882–898. <https://doi.org/10.1525/auk.2010.10111>
- Harrison T. 2011. Beak deformities of garden birds. *British Birds* 104: 538–541.
- Heath L, Martin DP, Warburton L, Perrin M, Horsfield W, Kingsley C, Rybicki EP, Williamson A-L (2004) Evidence of unique genotypes of beak and feather disease virus in southern Africa. *Journal of Virology* 78(17): 9277–9284. <https://doi.org/10.1128/jvi.78.17.9277-9284.2004>
- Hodges CJ, Poorboy DM, Weber BM, Thompson CF (2019) Beak abnormality hinders provisioning ability and reduces body condition of a female House Wren (*Troglodytes aedon*). *Wilson Journal of Ornithology* 131(1): 128–134. <https://doi.org/10.1676/18-50>
- iNaturalist (2026) Birds with deformed beaks. <https://www.inaturalist.org/projects/birds-with-deformed-beaks> [last accessed on 5 February 2026]
- Íthalo FR, Petters J, Cristina BCL (2021) Deformidade de bico em Tucanuçu (*Ramphastos toco*) da região de Chaco húmido no Paraguai: Relato de Caso. *Compendio de ciencias veterinarias* 11(1): 27–31. <https://doi.org/10.18004/compend.cienc.vet.2021.11.01.27>
- Jones CW, Risi MM, Kuntz W, Ryan PG, Steinfurth A, Bond AL (2015) Bill deformities in penguins (Spheniscidae): a global review *Marine Ornithology* 43(2), 207–209. <http://doi.org/10.5038/2074-1235.43.2.1132>
- Navalón G, Bright JA, Marugán-Lobón J, Rayfield EJ (2019) The evolutionary relationship among beak shape, mechanical advantage, and feeding ecology in modern birds. *Evolution* 73(3): 422–435. <https://doi.org/>

- 10.1111/evo.13655
- Gallo-Ortiz G (2011) Deformidade de bico em um indivíduo de papa-taoca-do-sul, *Pyriglena leucoptera* (Passeriformes: Thamnophilidae), na Mata Atlântica do Estado de São Paulo, Brasil. *Atualidades Ornitológicas* 164: 20-21.
- Gorosito CA, Gonda H, Cueto VR (2016) Beak deformities in north Patagonian birds. *Ornitología Neotropical* 27: 289–295. <https://doi.org/10.58843/ornneo.v27i0>
- Pomeroy DE (1962) Birds with abnormal bills. *British Birds* 56: 49–72.
- Purificação KN (2019) A case of beak deformity in the Shiny Cowbird *Molothrus bonariensis* and a review on beak deformities in wild birds in Brazil. *Revista Brasileira de Ornitologia* 27(3): 212–217. <https://doi.org/10.1007/BF03544473>
- Regnard GL, Boyes RS, Martin RO, Hitzeroth II, Rybicki EP (2015) Beak and feather disease viruses circulating in Cape parrots (*Poicephalus robustus*) in South Africa. *Archives of Virology* 160(1): 47–54. <https://doi.org/10.1007/s00705-014-2226-9>
- Rezende MA (2013) Um caso de deformidade de bico em Tucanuçu *Ramphastos toco* em Minas Gerais, sudeste do Brasil. *Atualidades Ornitológicas* 174: 23.
- Rogers LJ, Kaplan GT (2000) *Songs, Roars and Rituals: Communication in birds, mammals and other animals*. Harvard University Press, Boston, MA.
- Simmons RE (1997) Monteiro's Hornbill - Monteirose Neushoringvoël - *Tockus monteiri*. In: Harrison JA, Allan DG, Underhill LG, Herremans M, Tree AJ, Parker V, Brown CJ (eds) *The Atlas of Southern African Birds including Botswana, Lesotho, Namibia, South Africa, Swaziland and Zimbabwe - Volumes 1 (Non-Passerines)*. P. 706-707. Birdlife South Africa & Avian Demography Unit, South Africa.
- Smith P, Ríos SD, Doldán L, Ruíz R (2019) Bill deformations in five species of Paraguayan birds. *Revista Chilena de Ornitología* 25(1): 3–8.
- Tenez ED (2025) Reporte de caso: Eufonia Olivácea (*Euphonia gouldi*) con pico deforme en Guatemala. *Revista Científica (Instituto de Investigaciones Químicas y Biológicas. Facultad de Ciencias Químicas y Farmacia. Universidad de San Carlos de Guatemala)* 33(1): 20250721. [12 pp.] <https://doi.org/10.54495/Rev.Cientifica.v33i1.496>
- Tinajero R (2023) Hallazgo inédito de carboneros cresta negra (*Baeolophus atricristatus*) con picos deformes. *Huitzil Revista Mexicana de Ornitología* 24(1): e-646 [5 pp.]. <https://doi.org/10.28947/hrmo.2022.23.2.669>
- Valdebenito JO, Grandón-Ojeda A, Pantoja-Maggi V, Novoa FJ, González-Acuña D (2018) Report on beak abnormalities of some birds of Patagonia. *The Wilson Journal of Ornithology* 130(4): 1014–1019. <https://doi.org/10.1676/17-00039.1>
- Van Hemert C, Handel CM (2010) Beak deformities in Northwestern Crows: evidence of a multispecies epizootic. *The Auk* 127(4): 746–751. <https://doi.org/10.1525/auk.2010.10132>
- Van Hemert C, Handel CM, O'Hara TM (2012) Evidence of accelerated beak growth associated with avian keratin disorder in Black-capped Chickadees (*Poecile atricapillus*). *Journal of Wildlife Diseases* 48(3): 686–694. <https://doi.org/10.7589/0090-3558-48.3.686>
- Van Hemert C, Handel CM, Cottrell S, Gerik DE, Bildfell RJ (2025) Beak deformities in *Buteo jamaicensis* (Red-tailed Hawk) signal possible emergence of avian keratin disorder among raptors. *Ornithology* 142(1): ukaf023 [11 pp.]. <https://doi.org/10.1093/ornithology/ukaf023>
- Veltri CJ, Klem D (2005) Comparison of fatal bird injuries from collisions with towers and windows. *Journal of Field Ornithology* 76(2): 127–133. <https://doi.org/10.1648/0273-8570-76.2.127>
- Wiltschko R, Wiltschko WJ (2013) The magnetite-based receptors in the beak of birds and their role in avian navigation. *Journal of Comparative Physiology A* 199: 89–98. <https://doi.org/10.1007/s00359-012-0769-3>
- Zylberberg M, Van Hemert C, Dumbacher JP, Handel CM, Tihand T, DeRisi JL (2016) Novel picornavirus associated with avian keratin disorder in Alaskan birds. *mBio* 7(4): e00874-16 [10 pp.]. <https://doi.org/10.1128/mBio.00874-16>
- Zylberberg M, Van Hemert C, Handel CM, DeRisi JL (2018) Avian keratin disorder of Alaska black-capped chickadees is associated with *Poecivirus* infection. *Virology Journal* 15: 100 [9 pp.]. <https://doi.org/10.1186/s12985-018-1008-5>
- Zylberberg M, Van Hemert C, Handel CM, Liu RM, DeRisi JL (2021) *Poecivirus* is present in individuals with beak deformities in seven species of North American birds. *The Journal of Wildlife Diseases* 57(2): 273–281. <https://doi.org/10.7589/JWD-D-20-00017>